



南方科技大学
SOUTHERN UNIVERSITY OF SCIENCE AND TECHNOLOGY

Applied stochastic processes

Review for midterm

Jie Xiong

Department of Mathematics
Southern University of Science and Technology
Shenzhen, China, 518055

Math 208, Spring 2023

Review for midterm

1. If X has pdf f_X and g is 1-to-1, then $Y = g(X)$ has pdf

$$f_Y(y) = f_X(g^{-1}(y)) \left| \frac{d}{dy} g^{-1}(y) \right|.$$

Review for midterm

1. If X has pdf f_X and g is 1-to-1, then $Y = g(X)$ has pdf

$$f_Y(y) = f_X(g^{-1}(y)) \left| \frac{d}{dy} g^{-1}(y) \right|.$$

2. If $X = (X_1, X_2)$ has joint pdf $f_X(x)$ and $g: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is 1-to-1, then $(Y_1, Y_2) = Y = g(X)$ has joint pdf f_Y given by

$$f_Y(y) = f_X(g^{-1}(y)) |J|^{-1}$$

where J is the Jacobian determinant

$$J = \begin{vmatrix} \frac{\partial g_1}{\partial x_1} & \frac{\partial g_1}{\partial x_2} \\ \frac{\partial g_2}{\partial x_1} & \frac{\partial g_2}{\partial x_2} \end{vmatrix}$$



3. The moment generating function (m.g.f.) of X is

$$\phi(t) = \mathbb{E}(e^{tX}) = \begin{cases} \sum_x e^{tx} p(x) & \text{if } X \text{ discrete} \end{cases}$$



3. The moment generating function (m.g.f.) of X is

$$\phi(t) = \mathbb{E}(e^{tX}) = \begin{cases} \sum_x e^{tx} p(x) & \text{if } X \text{ discrete} \\ \int_{-\infty}^{\infty} e^{ix} f(x) dx & \text{if } X \text{ continuous.} \end{cases}$$



3. The moment generating function (m.g.f.) of X is

$$\phi(t) = \mathbb{E}(e^{tX}) = \begin{cases} \sum_x e^{tx} p(x) & \text{if } X \text{ discrete} \\ \int_{-\infty}^{\infty} e^{ix} f(x) dx & \text{if } X \text{ continuous.} \end{cases}$$



$$\mathbb{E}(X^n) = \phi^{(n)}(0).$$



3. The moment generating function (m.g.f.) of X is

$$\phi(t) = \mathbb{E}(e^{tX}) = \begin{cases} \sum_x e^{tx} p(x) & \text{if } X \text{ discrete} \\ \int_{-\infty}^{\infty} e^{ix} f(x) dx & \text{if } X \text{ continuous.} \end{cases}$$



$$\mathbb{E}(X^n) = \phi^{(n)}(0).$$

- If X and Y are independent and a, b are constants, then

$$\phi_{aX+bY}(t) = \phi_X(at)\phi_Y(bt).$$

3. The moment generating function (m.g.f.) of X is

$$\phi(t) = \mathbb{E}(e^{tX}) = \begin{cases} \sum_x e^{tx} p(x) & \text{if } X \text{ discrete} \\ \int_{-\infty}^{\infty} e^{ix} f(x) dx & \text{if } X \text{ continuous.} \end{cases}$$



$$\mathbb{E}(X^n) = \phi^{(n)}(0).$$

- If X and Y are independent and a, b are constants, then

$$\phi_{aX+bY}(t) = \phi_X(at)\phi_Y(bt).$$

4. A stochastic process is a family of random variables $\{X_t : t \in \mathbb{T}\}$ with time set \mathbb{T} and state S .

5. For discrete random variables, then

$$p_{X|Y}(x|y) = \frac{p(x, y)}{p_Y(y)},$$

is the conditional pmf of X given $Y = y$.

5. For discrete random variables, then

$$p_{X|Y}(x|y) = \frac{p(x, y)}{p_Y(y)},$$

is the conditional pmf of X given $Y = y$.

- The conditional expectation of X given $Y = y$:

$$E(X|Y = y) = \sum_x xp_{X|Y}(x|y).$$

5. For discrete random variables, then

$$p_{X|Y}(x|y) = \frac{p(x, y)}{p_Y(y)},$$

is the conditional pmf of X given $Y = y$.

- The conditional expectation of X given $Y = y$:

$$E(X|Y = y) = \sum_x xp_{X|Y}(x|y).$$

- The conditional expectation of X given Y is

$$E(X|Y) = \sum_x xp_{X|Y}(x|Y).$$

5. For discrete random variables, then

$$p_{X|Y}(x|y) = \frac{p(x, y)}{p_Y(y)},$$

is the conditional pmf of X given $Y = y$.

- The conditional expectation of X given $Y = y$:

$$E(X|Y = y) = \sum_x xp_{X|Y}(x|y).$$

- The conditional expectation of X given Y is

$$E(X|Y) = \sum_x xp_{X|Y}(x|Y).$$

- If X and Y are independent, then $E(X|Y) = E(X)$.



6. For continuous random variables, then, the conditional pdf of X given $Y = y$ is

$$f_{X|Y}(x|y) = \frac{f(x, y)}{f_Y(y)}.$$



6. For continuous random variables, then, the conditional pdf of X given $Y = y$ is

$$f_{X|Y}(x|y) = \frac{f(x, y)}{f_Y(y)}.$$

- The conditional expectation of X given $Y = y$ is

$$E(X|Y = y) = \int_{-\infty}^{\infty} x f_{X|Y}(x|y) dx.$$



6. For continuous random variables, then, the conditional pdf of X given $Y = y$ is

$$f_{X|Y}(x|y) = \frac{f(x, y)}{f_Y(y)}.$$

- The conditional expectation of X given $Y = y$ is

$$E(X|Y = y) = \int_{-\infty}^{\infty} x f_{X|Y}(x|y) dx.$$

- The conditional expectation of X given Y is

$$E(X|Y) = \int_{-\infty}^{\infty} x f_{X|Y}(x|Y) dx.$$



7. Calculating expectation by conditioning

$$E(X) = E(E(X|Y)).$$

7. Calculating expectation by conditioning

$$E(X) = E(E(X|Y)).$$

8. Calculating variance by conditioning

$$V(X) = E(V(X|Y)) + V(E(X|Y)).$$

7. Calculating expectation by conditioning

$$E(X) = E(E(X|Y)).$$

8. Calculating variance by conditioning

$$V(X) = E(V(X|Y)) + V(E(X|Y)).$$

9. Calculating probability by conditioning

$$P(A) = E(P(A|Y)).$$



10. $\{X_n\}$ is a *Markov chain* if

$$\mathbb{P}(X_{n+1} = j | X_n = i, X_{n-1} = i_{n-1}, \dots, X_0 = i_0) = p(i, j),$$

which is the *transition matrix* of the chain.



10. $\{X_n\}$ is a *Markov chain* if

$$\mathbb{P}(X_{n+1} = j | X_n = i, X_{n-1} = i_{n-1}, \dots, X_0 = i_0) = p(i, j),$$

which is the *transition matrix* of the chain.

- The m -step transition matrix is P^m , namely, $P^{(m)} = P^m$.



南方科技大学
SOUTHERN UNIVERSITY OF SCIENCE AND TECHNOLOGY

11. Time to y .



11. Time to y .

- $T_y = \inf\{n \geq 1 : X_n = y\}$ is first time return to (or reach) y .



11. Time to y .

- $T_y = \inf\{n \geq 1 : X_n = y\}$ is first time return to (or reach) y .



$$\rho_{yy} = \mathbb{P}_y(T_y < \infty)$$

is the prob. of returning to initial y eventually.



11. Time to y .

- $T_y = \inf\{n \geq 1 : X_n = y\}$ is first time return to (or reach) y .



$$\rho_{yy} = \mathbb{P}_y(T_y < \infty)$$

is the prob. of returning to initial y eventually.



$$T_y^1 = T_y, \quad T_y^k = \inf\{n > T_y^{k-1} : X_n = y\}.$$



11. Time to y .

- $T_y = \inf\{n \geq 1 : X_n = y\}$ is first time return to (or reach) y .



$$\rho_{yy} = \mathbb{P}_y(T_y < \infty)$$

is the prob. of returning to initial y eventually.



$$T_y^1 = T_y, \quad T_y^k = \inf\{n > T_y^{k-1} : X_n = y\}.$$



$$\mathbb{P}_y(T_y^k < \infty) = \rho_{yy}^k.$$



12. Status of states:

- 1 y is *transient* if $\rho_{yy} < 1$.



12. Status of states:

- ① y is *transient* if $\rho_{yy} < 1$.
- ② y is *recurrent* if $\rho_{yy} = 1$.



12. Status of states:

- ① y is *transient* if $\rho_{yy} < 1$.
- ② y is *recurrent* if $\rho_{yy} = 1$.

- If y is transient, then

$$\mathbb{P}_y(X_n = y, i.o.) = 0.$$

12. Status of states:

- ① y is *transient* if $\rho_{yy} < 1$.
- ② y is *recurrent* if $\rho_{yy} = 1$.

- If y is transient, then

$$\mathbb{P}_y(X_n = y, i.o.) = 0.$$

- If y is recurrent, then

$$\mathbb{P}_y(X_n = y, i.o.) = 1.$$



13. x communicates with y , write $x \rightarrow y$, if

$$\rho_{xy} \equiv \mathbb{P}_x(T_y < \infty) > 0.$$



13. x communicates with y , write $x \rightarrow y$, if

$$\rho_{xy} \equiv \mathbb{P}_x(T_y < \infty) > 0.$$

- If $x \rightarrow y$ and $y \rightarrow z$, then $x \rightarrow z$.



13. x communicates with y , write $x \rightarrow y$, if

$$\rho_{xy} \equiv \mathbb{P}_x(T_y < \infty) > 0.$$

- If $x \rightarrow y$ and $y \rightarrow z$, then $x \rightarrow z$.
- If $\rho_{xy} > 0$ and $\rho_{yx} < 1$, then x is transient.



13. x communicates with y , write $x \rightarrow y$, if

$$\rho_{xy} \equiv \mathbb{P}_x(T_y < \infty) > 0.$$

- If $x \rightarrow y$ and $y \rightarrow z$, then $x \rightarrow z$.
- If $\rho_{xy} > 0$ and $\rho_{yx} < 1$, then x is transient.
- If x is recurrent and $\rho_{xy} > 0$, then $\rho_{yx} = 1$.



14. Decomposition



14. Decomposition

- A set A is *closed* if it is impossible to get out.



14. Decomposition

- A set A is *closed* if it is impossible to get out.
- A set B is *irreducible* if $\forall i, j \in B, i \rightarrow j$.



14. Decomposition

- A set A is *closed* if it is impossible to get out.
- A set B is *irreducible* if $\forall i, j \in B, i \rightarrow j$.
- If C is a finite closed irreducible set, then all states in C are recurrent.



14. Decomposition

- A set A is *closed* if it is impossible to get out.
- A set B is *irreducible* if $\forall i, j \in B, i \rightarrow j$.
- If C is a finite closed irreducible set, then all states in C are recurrent.
- If the state space S is finite, then S can be written as a disjoint union

$$S = T \cup R_1 \cup \cdots \cup R_k,$$

where T is a set of transient states and $R_i, 1 \leq i \leq k$, are closed irreducible sets of recurrent states.



15.

$$N(y) = \# \text{ of visits to } y.$$



15.

$N(y)$ = # of visits to y .



$$\mathbb{E}_x N(y) = \frac{\rho_{xy}}{1 - \rho_{yy}} = \sum_{n=1}^{\infty} p^n(x, y).$$



15.

$N(y) = \#$ of visits to y .



$$\mathbb{E}_x N(y) = \frac{\rho_{xy}}{1 - \rho_{yy}} = \sum_{n=1}^{\infty} p^n(x, y).$$

- y is recurrent iff

$$\sum_{n=1}^{\infty} p^n(y, y) = \mathbb{E}_y N(y) = \infty.$$



15.

$N(y) = \#$ of visits to y .



$$\mathbb{E}_x N(y) = \frac{\rho_{xy}}{1 - \rho_{yy}} = \sum_{n=1}^{\infty} p^n(x, y).$$

- y is recurrent iff

$$\sum_{n=1}^{\infty} p^n(y, y) = \mathbb{E}_y N(y) = \infty.$$

- If x is recurrent and $x \rightarrow y$, then y is recurrent.



15.

$N(y) = \#$ of visits to y .



$$\mathbb{E}_x N(y) = \frac{\rho_{xy}}{1 - \rho_{yy}} = \sum_{n=1}^{\infty} p^n(x, y).$$

- y is recurrent iff

$$\sum_{n=1}^{\infty} p^n(y, y) = \mathbb{E}_y N(y) = \infty.$$

- If x is recurrent and $x \rightarrow y$, then y is recurrent.
- In a finite closed set there has to be at least one recurrent state.



16. Vector π is a *stationary distribution* if $\pi P = \pi$.



16. Vector π is a *stationary distribution* if $\pi P = \pi$.

- Suppose that the $k \times k$ transition matrix is irreducible. Then, there is a unique solution to $\pi P = \pi$ with $\sum_{i=1}^k \pi_i = 1$. Further, $\pi_i > 0, \forall i$.



17. Transition matrix P is *double stochastic* if its each column sums to 1, i.e.

$$\sum_x p(x, y) = 1, \quad \forall y.$$



17. Transition matrix P is *double stochastic* if its each column sums to 1, i.e.

$$\sum_x p(x, y) = 1, \quad \forall y.$$

- If P is double stochastic with N states, then $\pi(x) = \frac{1}{N}$, $\forall x$ is a stationary distribution.



18. A distribution π satisfies the *detailed balance condition* if

$$\pi(x)p(x, y) = \pi(y)p(y, x), \quad \forall x, y \in S.$$



18. A distribution π satisfies the *detailed balance condition* if

$$\pi(x)p(x, y) = \pi(y)p(y, x), \quad \forall x, y \in S.$$

- If π satisfies the DBC, then π is a stationary distribution.



19. Let X_n be the MC with transition probability $p(i, j)$ and starting from the stationary distribution.



19. Let X_n be the MC with transition probability $p(i, j)$ and starting from the stationary distribution.

Let n be fixed and

$$Y_m = X_{n-m}, \quad 0 \leq m \leq n.$$

19. Let X_n be the MC with transition probability $p(i, j)$ and starting from the stationary distribution.

Let n be fixed and

$$Y_m = X_{n-m}, \quad 0 \leq m \leq n.$$

Then, (Y_m) is a MC with transition probability

$$\hat{p}(i, j) = \frac{\pi(j)p(j, i)}{\pi(i)}.$$

19. Let X_n be the MC with transition probability $p(i, j)$ and starting from the stationary distribution.

Let n be fixed and

$$Y_m = X_{n-m}, \quad 0 \leq m \leq n.$$

Then, (Y_m) is a MC with transition probability

$$\hat{p}(i, j) = \frac{\pi(j)p(j, i)}{\pi(i)}.$$

- The MC is *reversible* if $\hat{p}(i, j) = p(i, j)$, $\forall i, j \in S$.



19. Let X_n be the MC with transition probability $p(i, j)$ and starting from the stationary distribution.

Let n be fixed and

$$Y_m = X_{n-m}, \quad 0 \leq m \leq n.$$

Then, (Y_m) is a MC with transition probability

$$\hat{p}(i, j) = \frac{\pi(j)p(j, i)}{\pi(i)}.$$

- The MC is *reversible* if $\hat{p}(i, j) = p(i, j)$, $\forall i, j \in S$.
- (X_n) is reversible iff DBC holds.



20. Limit behavior



20. Limit behavior

Under S , all states y satisfying $\pi(y) > 0$ are recurrent.



20. Limit behavior

Under S , all states y satisfying $\pi(y) > 0$ are recurrent.



21. Let

$$N_n(y) = \sum_{i=1}^n 1_{X_i=y}$$

be the # of visits to y before time n .



21. Let

$$N_n(y) = \sum_{i=1}^n 1_{X_i=y}$$

be the # of visits to y before time n .

- Suppose (I) and (R) hold. Then,

$$\lim_{n \rightarrow \infty} \frac{N_n(y)}{n} = \frac{1}{\mathbb{E}_y T_y}, \quad a.s.$$



21. Let

$$N_n(y) = \sum_{i=1}^n 1_{X_i=y}$$

be the # of visits to y before time n .

- Suppose (I) and (R) hold. Then,

$$\lim_{n \rightarrow \infty} \frac{N_n(y)}{n} = \frac{1}{\mathbb{E}_y T_y}, \quad a.s.$$

- Under (I) and (S), we have

$$\pi(y) = \frac{1}{\mathbb{E}_y T_y},$$



22. Under (I), (S) and $\sum_x |f(x)|\pi(x) < \infty$, we have

$$\frac{1}{n} \sum_{m=1}^n f(X_m) \rightarrow \sum_x f(x)\pi(x), \quad a.s.$$

22. Under (I), (S) and $\sum_x |f(x)|\pi(x) < \infty$, we have

$$\frac{1}{n} \sum_{m=1}^n f(X_m) \rightarrow \sum_x f(x)\pi(x), \quad a.s.$$

Namely, long term average equals spatial average, i.e., ergodicity.



23. A recurrent state x is *positive recurrent* if $\mathbb{E}_x T_x < \infty$;



23. A recurrent state x is *positive recurrent* if $\mathbb{E}_x T_x < \infty$; it is *null recurrent* if $\mathbb{E}_x T_x = \infty$.



23. A recurrent state x is *positive recurrent* if $\mathbb{E}_x T_x < \infty$; it is *null recurrent* if $\mathbb{E}_x T_x = \infty$.

For any irreducible chain the following are equivalent:



23. A recurrent state x is *positive recurrent* if $\mathbb{E}_x T_x < \infty$; it is *null recurrent* if $\mathbb{E}_x T_x = \infty$.

For any irreducible chain the following are equivalent:

- 1 There is a positive recurrent state.



23. A recurrent state x is *positive recurrent* if $\mathbb{E}_x T_x < \infty$; it is *null recurrent* if $\mathbb{E}_x T_x = \infty$.

For any irreducible chain the following are equivalent:

- 1 There is a positive recurrent state.
- 2 There is a stationary distribution π .



23. A recurrent state x is *positive recurrent* if $\mathbb{E}_x T_x < \infty$; it is *null recurrent* if $\mathbb{E}_x T_x = \infty$.

For any irreducible chain the following are equivalent:

- ① There is a positive recurrent state.
- ② There is a stationary distribution π .
- ③ All states are positive recurrent.



24. Branching process: $X_n = \#$ of individuals in n th generation.



24. Branching process: $X_n = \#$ of individuals in n th generation.
 $Y_i = \#$ of children of i th individual.



24. Branching process: $X_n = \#$ of individuals in n th generation.
 $Y_i = \#$ of children of i th individual.

$$\mathbb{P}(Y_i = k) = p_k, \quad k = 0, 1, 2, \dots$$



24. Branching process: $X_n = \#$ of individuals in n th generation.
 $Y_i = \#$ of children of i th individual.

$$\mathbb{P}(Y_i = k) = p_k, \quad k = 0, 1, 2, \dots.$$

Let

$$\phi(\theta) = \sum_{k=0}^{\infty} p_k \theta^k.$$



24. Branching process: $X_n = \#$ of individuals in n th generation.
 $Y_i = \#$ of children of i th individual.

$$\mathbb{P}(Y_i = k) = p_k, \quad k = 0, 1, 2, \dots.$$

Let

$$\phi(\theta) = \sum_{k=0}^{\infty} p_k \theta^k.$$

Let $\rho = \mathbb{P}_1(\text{extinction})$.

24. Branching process: $X_n = \#$ of individuals in n th generation.
 $Y_i = \#$ of children of i th individual.

$$\mathbb{P}(Y_i = k) = p_k, \quad k = 0, 1, 2, \dots.$$

Let

$$\phi(\theta) = \sum_{k=0}^{\infty} p_k \theta^k.$$

Let $\rho = \mathbb{P}_1(\text{extinction})$. ρ is the smallest root of
 $x = \phi(x)$, $0 \leq x \leq 1$.