



南方科技大学
SOUTHERN UNIVERSITY OF SCIENCE AND TECHNOLOGY

Applied stochastic processes

Chapter 3

Jie Xiong

Department of Mathematics
Southern University of Science and Technology
Shenzhen, China, 518055

Math 208, Spring 2021



Chapter 3. Markov chain: Basics



Chapter 3. Markov chain: Basics

3.1. Definitions and examples



Chapter 3. Markov chain: Basics

3.1. Definitions and examples

Example (Gambler's ruin)

In a game, you win \$1 with prob. $p = .4$ or lose \$1 w/ prob. $1 - p = .6$. Suppose you will stop if your fortune reaches $\$N$ (or $\$0$ when you have to).



南方科技大学
SOUTHERN UNIVERSITY OF SCIENCE AND TECHNOLOGY

Let

$$X_n =$$

Let

$X_n =$ your money after n plays.

Let

$X_n =$ your money after n plays.

Then,

$$\mathbb{P}(X_{n+1} = i + 1 | X_n = i, X_{n-1} = i_{n-1}, \dots, X_0 = i_0) =$$

Let

$X_n =$ your money after n plays.

Then,

$$\mathbb{P}(X_{n+1} = i + 1 | X_n = i, X_{n-1} = i_{n-1}, \dots, X_0 = i_0) = .4,$$

Let

$X_n =$ your money after n plays.

Then,

$$\mathbb{P}(X_{n+1} = i + 1 | X_n = i, X_{n-1} = i_{n-1}, \dots, X_0 = i_0) = .4,$$

and

$$\mathbb{P}(X_{n+1} = i - 1 | X_n = i, X_{n-1} = i_{n-1}, \dots, X_0 = i_0) =$$

Let

$X_n =$ your money after n plays.

Then,

$$\mathbb{P}(X_{n+1} = i + 1 | X_n = i, X_{n-1} = i_{n-1}, \dots, X_0 = i_0) = .4,$$

and

$$\mathbb{P}(X_{n+1} = i - 1 | X_n = i, X_{n-1} = i_{n-1}, \dots, X_0 = i_0) = .6.$$

Let

$X_n =$ your money after n plays.

Then,

$$\mathbb{P}(X_{n+1} = i + 1 | X_n = i, X_{n-1} = i_{n-1}, \dots, X_0 = i_0) = .4,$$

and

$$\mathbb{P}(X_{n+1} = i - 1 | X_n = i, X_{n-1} = i_{n-1}, \dots, X_0 = i_0) = .6.$$

Namely, given X_n , the state X_{n+1} is indep. of the past X_0, X_1, \dots, X_{n-1} .



Definition

$\{X_n\}$ is a *Markov chain* if

$$\mathbb{P}(X_{n+1} = j | X_n = i, X_{n-1} = i_{n-1}, \dots, X_0 = i_0) = p(i, j),$$

which is the *transition matrix* of the chain.



Definition

$\{X_n\}$ is a *Markov chain* if

$$\mathbb{P}(X_{n+1} = j | X_n = i, X_{n-1} = i_{n-1}, \dots, X_0 = i_0) = p(i, j),$$

which is the *transition matrix* of the chain.

Remark

$$\mathbb{P}(X_{n+1} = j | X_n = i) = p(i, j).$$



Example 1 (Conti.)

$$p(i, i + 1) = .4,$$



Example 1 (Conti.)

$$p(i, i + 1) = .4, \quad p(i, i - 1) = .6,$$



Example 1 (Conti.)

$$p(i, i + 1) = .4, \quad p(i, i - 1) = .6, \quad 0 < i < N,$$



Example 1 (Conti.)

$$p(i, i + 1) = .4, \quad p(i, i - 1) = .6, \quad 0 < i < N,$$

$$p(0, 0) = p(N, N) = 1.$$

If $N = 3$, transition matrix is 4×4 :



Example 1 (Conti.)

$$p(i, i+1) = .4, \quad p(i, i-1) = .6, \quad 0 < i < N,$$

$$p(0, 0) = p(N, N) = 1.$$

If $N = 3$, transition matrix is 4×4 :

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ .6 & 0 & .4 & 0 \\ 0 & .6 & 0 & .4 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$



Example 1 (Conti.)

$$p(i, i+1) = .4, \quad p(i, i-1) = .6, \quad 0 < i < N,$$

$$p(0, 0) = p(N, N) = 1.$$

If $N = 3$, transition matrix is 4×4 :

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ .6 & 0 & .4 & 0 \\ 0 & .6 & 0 & .4 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

We can also write



Example 1 (Conti.)

$$p(i, i+1) = .4, \quad p(i, i-1) = .6, \quad 0 < i < N,$$

$$p(0, 0) = p(N, N) = 1.$$

If $N = 3$, transition matrix is 4×4 :

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ .6 & 0 & .4 & 0 \\ 0 & .6 & 0 & .4 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

We can also write

$$0 \leftarrow 1 \xleftrightarrow{\quad} 2 \rightarrow 3.$$



Example (Ehrenfest chain)

N ball total in two urns side by side. Pick one ball at random to move to the other urn.



Example (Ehrenfest chain)

N ball total in two urns side by side. Pick one ball at random to move to the other urn.

$X_n = \#$ of balls in the left urn after n picks.



Example (Ehrenfest chain)

N ball total in two urns side by side. Pick one ball at random to move to the other urn.

$X_n = \#$ of balls in the left urn after n picks.

Then

$$\mathbb{P}(X_{n+1} = i + 1 | X_n = i, X_{n-1} = i_{n-1}, \dots, X_0 = i_0) =$$

Example (Ehrenfest chain)

N ball total in two urns side by side. Pick one ball at random to move to the other urn.

$X_n = \#$ of balls in the left urn after n picks.

Then

$$\mathbb{P}(X_{n+1} = i + 1 | X_n = i, X_{n-1} = i_{n-1}, \dots, X_0 = i_0) = \frac{N - i}{N}.$$



Example (Ehrenfest chain)

N ball total in two urns side by side. Pick one ball at random to move to the other urn.

$X_n = \#$ of balls in the left urn after n picks.

Then

$$\mathbb{P}(X_{n+1} = i + 1 | X_n = i, X_{n-1} = i_{n-1}, \dots, X_0 = i_0) = \frac{N - i}{N}.$$

So,

$$p(i, i + 1) = \frac{N - i}{N},$$



Example (Ehrenfest chain)

N ball total in two urns side by side. Pick one ball at random to move to the other urn.

$X_n = \#$ of balls in the left urn after n picks.

Then

$$\mathbb{P}(X_{n+1} = i + 1 | X_n = i, X_{n-1} = i_{n-1}, \dots, X_0 = i_0) = \frac{N - i}{N}.$$

So,

$$p(i, i + 1) = \frac{N - i}{N}, \quad p(i, i - 1) = \frac{i}{N},$$



Example (Ehrenfest chain)

N ball total in two urns side by side. Pick one ball at random to move to the other urn.

$X_n = \#$ of balls in the left urn after n picks.

Then

$$\mathbb{P}(X_{n+1} = i + 1 | X_n = i, X_{n-1} = i_{n-1}, \dots, X_0 = i_0) = \frac{N - i}{N}.$$

So,

$$p(i, i + 1) = \frac{N - i}{N}, \quad p(i, i - 1) = \frac{i}{N}, \quad 0 \leq i \leq N.$$



Example (Ehrenfest chain)

N ball total in two urns side by side. Pick one ball at random to move to the other urn.

$X_n = \#$ of balls in the left urn after n picks.

Then

$$\mathbb{P}(X_{n+1} = i + 1 | X_n = i, X_{n-1} = i_{n-1}, \dots, X_0 = i_0) = \frac{N - i}{N}.$$

So,

$$p(i, i + 1) = \frac{N - i}{N}, \quad p(i, i - 1) = \frac{i}{N}, \quad 0 \leq i \leq N.$$

$$p(i, j) = 0, \quad \text{otherwise.}$$



南方科技大学
SOUTHERN UNIVERSITY OF SCIENCE AND TECHNOLOGY

Facts:



Facts: i) $p(i, j) \geq 0$.



Facts: i) $p(i, j) \geq 0$.
ii)

$$\sum_j p(i, j) = 1.$$



Facts: i) $p(i, j) \geq 0$.
ii)

$$\sum_j p(i, j) = 1. \quad (\text{After } X_n = i, X_{n+1} \text{ will be at a state } j).$$



Example (Inventory chain)

Let $s < S$ and D_n the demand of certain product for day n .



Example (Inventory chain)

Let $s < S$ and D_n the demand of certain product for day n .

If the inventory is s or below, we order enough to bring it back to S .



Example (Inventory chain)

Let $s < S$ and D_n the demand of certain product for day n .

If the inventory is s or below, we order enough to bring it back to S . Let X_n be the inventory at the end of day n .



Example (Inventory chain)

Let $s < S$ and D_n the demand of certain product for day n .
If the inventory is s or below, we order enough to bring it back to S . Let X_n be the inventory at the end of day n . Suppose D_1, D_2, \dots are i.i.d.



Example (Inventory chain)

Let $s < S$ and D_n the demand of certain product for day n .
If the inventory is s or below, we order enough to bring it back to S . Let X_n be the inventory at the end of day n . Suppose D_1, D_2, \dots are i.i.d.

$$X_{n+1} =$$



Example (Inventory chain)

Let $s < S$ and D_n the demand of certain product for day n .
If the inventory is s or below, we order enough to bring it back to S . Let X_n be the inventory at the end of day n . Suppose D_1, D_2, \dots are i.i.d.

$$X_{n+1} = \begin{cases} (X_n - D_{n+1})^+ & \text{if } X_n > s \\ S & \text{if } X_n \leq s \end{cases}$$



Example (Inventory chain)

Let $s < S$ and D_n the demand of certain product for day n .
If the inventory is s or below, we order enough to bring it back to S . Let X_n be the inventory at the end of day n . Suppose D_1, D_2, \dots are i.i.d.

$$X_{n+1} = \begin{cases} (X_n - D_{n+1})^+ & \text{if } X_n > s \\ (S - D_{n+1})^+ & \text{if } X_n \leq s. \end{cases}$$



Example (Inventory chain)

Let $s < S$ and D_n the demand of certain product for day n . If the inventory is s or below, we order enough to bring it back to S . Let X_n be the inventory at the end of day n . Suppose D_1, D_2, \dots are i.i.d.

$$X_{n+1} = \begin{cases} (X_n - D_{n+1})^+ & \text{if } X_n > s \\ (S - D_{n+1})^+ & \text{if } X_n \leq s. \end{cases}$$

Then, $\{X_n\}$ is a Markov chain.



Example (Inventory chain)

Let $s < S$ and D_n the demand of certain product for day n .
If the inventory is s or below, we order enough to bring it back to S . Let X_n be the inventory at the end of day n . Suppose D_1, D_2, \dots are i.i.d.

$$X_{n+1} = \begin{cases} (X_n - D_{n+1})^+ & \text{if } X_n > s \\ (S - D_{n+1})^+ & \text{if } X_n \leq s. \end{cases}$$

Then, $\{X_n\}$ is a Markov chain.

Suppose $s = 1, S = 5$



Example (Inventory chain)

Let $s < S$ and D_n the demand of certain product for day n .
If the inventory is s or below, we order enough to bring it back to S . Let X_n be the inventory at the end of day n . Suppose D_1, D_2, \dots are i.i.d.

$$X_{n+1} = \begin{cases} (X_n - D_{n+1})^+ & \text{if } X_n > s \\ (S - D_{n+1})^+ & \text{if } X_n \leq s. \end{cases}$$

Then, $\{X_n\}$ is a Markov chain.

Suppose $s = 1$, $S = 5$ and the prob. function of D_1 is

x	0	1	2	3
p(x)	.3	.4	.2	.1



Example (Inventory chain)

Let $s < S$ and D_n the demand of certain product for day n .
If the inventory is s or below, we order enough to bring it back to S . Let X_n be the inventory at the end of day n . Suppose D_1, D_2, \dots are i.i.d.

$$X_{n+1} = \begin{cases} (X_n - D_{n+1})^+ & \text{if } X_n > s \\ (S - D_{n+1})^+ & \text{if } X_n \leq s. \end{cases}$$

Then, $\{X_n\}$ is a Markov chain.

Suppose $s = 1, S = 5$ and the prob. function of D_1 is

x	0	1	2	3
p(x)	.3	.4	.2	.1

Find the transition matrix.



南方科技大学
SOUTHERN UNIVERSITY OF SCIENCE AND TECHNOLOGY

Solution: Do it on board!



Solution: Do it on board! If $X_n = 0, 1$ or 5 , then
 $X_{n+1} = 5 - D_{n+1} = 5, 4, 3, 2$.

$$P = \begin{bmatrix} 0 & 0 & .1 & .2 & .4 & .3 \\ 0 & 0 & .1 & .2 & .4 & .3 \\ & & & & & \\ & & & & & \\ 0 & 0 & .1 & .2 & .4 & .3 \end{bmatrix}$$



If $X_n = 2$, then $X_{n+1} = (2 - D_{n+1})^+ = 2, 1, 0, 0$.

$$P = \begin{bmatrix} 0 & 0 & .1 & .2 & .4 & .3 \\ 0 & 0 & .1 & .2 & .4 & .3 \\ .3 & .4 & .3 & 0 & 0 & 0 \\ .1 & .2 & .4 & .3 & 0 & 0 \\ 0 & .1 & .2 & .4 & .3 & 0 \\ 0 & 0 & .1 & .2 & .4 & .3 \end{bmatrix}$$





Example (Branching process)



Example (Branching process)

In the population, each individual gives birth (indep. of others) to k children with prob. p_k , $k = 0, 1, 2, \dots$.



Example (Branching process)

In the population, each individual gives birth (indep. of others) to k children with prob. p_k , $k = 0, 1, 2, \dots$. Let X_n be the # of individuals in n th generation.



Example (Branching process)

In the population, each individual gives birth (indep. of others) to k children with prob. p_k , $k = 0, 1, 2, \dots$. Let X_n be the # of individuals in n th generation. Then, $\{X_n\}$ is a MC.



Example (Branching process)

In the population, each individual gives birth (indep. of others) to k children with prob. p_k , $k = 0, 1, 2, \dots$. Let X_n be the # of individuals in n th generation. Then, $\{X_n\}$ is a MC.

If $X_n = i$,



Example (Branching process)

In the population, each individual gives birth (indep. of others) to k children with prob. p_k , $k = 0, 1, 2, \dots$. Let X_n be the # of individuals in n th generation. Then, $\{X_n\}$ is a MC.

If $X_n = i$, let Y_k , $k = 1, 2, \dots, i$ be the # of children by k th individual.

Example (Branching process)

In the population, each individual gives birth (indep. of others) to k children with prob. p_k , $k = 0, 1, 2, \dots$. Let X_n be the # of individuals in n th generation. Then, $\{X_n\}$ is a MC.

If $X_n = i$, let Y_k , $k = 1, 2, \dots, i$ be the # of children by k th individual. Then,

$$X_{n+1} = Y_1 + \dots + Y_i.$$



Example (Branching process)

In the population, each individual gives birth (indep. of others) to k children with prob. p_k , $k = 0, 1, 2, \dots$. Let X_n be the # of individuals in n th generation. Then, $\{X_n\}$ is a MC.

If $X_n = i$, let Y_k , $k = 1, 2, \dots, i$ be the # of children by k th individual. Then,

$$X_{n+1} = Y_1 + \dots + Y_i.$$

So,

$$p(i, j) = \mathbb{P}(Y_1 + \dots + Y_i = j), \quad i > 0, \quad j \geq 0.$$



Example (Branching process)

In the population, each individual gives birth (indep. of others) to k children with prob. p_k , $k = 0, 1, 2, \dots$. Let X_n be the # of individuals in n th generation. Then, $\{X_n\}$ is a MC.

If $X_n = i$, let Y_k , $k = 1, 2, \dots, i$ be the # of children by k th individual. Then,

$$X_{n+1} = Y_1 + \dots + Y_i.$$

So,

$$p(i, j) = \mathbb{P}(Y_1 + \dots + Y_i = j), \quad i > 0, \quad j \geq 0.$$

Q: What is the extinction probability?



Example (Wright-Fisher model)



Example (Wright-Fisher model)

Population of N genes that can be of two types: A or a .



Example (Wright-Fisher model)

Population of N genes that can be of two types: A or a .
Population at time $n + 1$ is obtained by drawing with replacement from the population at time n .



Example (Wright-Fisher model)

Population of N genes that can be of two types: A or a .
Population at time $n + 1$ is obtained by drawing with replacement from the population at time n . Let X_n be # of A alleles at time n .



Example (Wright-Fisher model)

Population of N genes that can be of two types: A or a .
Population at time $n + 1$ is obtained by drawing with replacement from the population at time n . Let X_n be # of A alleles at time n . Then, $\{X_n\}$ is a MC.



Example (Wright-Fisher model)

Population of N genes that can be of two types: A or a .
Population at time $n + 1$ is obtained by drawing with replacement from the population at time n . Let X_n be # of A alleles at time n . Then, $\{X_n\}$ is a MC.

$$p(i, j) = P(X_{n+1} = j | X_n = i)$$



Example (Wright-Fisher model)

Population of N genes that can be of two types: A or a . Population at time $n + 1$ is obtained by drawing with replacement from the population at time n . Let X_n be # of A alleles at time n . Then, $\{X_n\}$ is a MC.

$$\begin{aligned} p(i, j) &= P(X_{n+1} = j | X_n = i) \\ &= \binom{N}{j} \left(\frac{i}{N}\right)^j \left(1 - \frac{i}{N}\right)^{N-j} \end{aligned}$$



Example (Wright-Fisher model)

Population of N genes that can be of two types: A or a .
Population at time $n + 1$ is obtained by drawing with replacement from the population at time n . Let X_n be # of A alleles at time n . Then, $\{X_n\}$ is a MC.

$$\begin{aligned} p(i, j) &= P(X_{n+1} = j | X_n = i) \\ &= \binom{N}{j} \left(\frac{i}{N}\right)^j \left(1 - \frac{i}{N}\right)^{N-j} \\ &\quad (\text{binomial with } p = \frac{i}{N} \text{ of } N \text{ trials}) \end{aligned}$$



Example (Wright-Fisher model)

Population of N genes that can be of two types: A or a . Population at time $n + 1$ is obtained by drawing with replacement from the population at time n . Let X_n be # of A alleles at time n . Then, $\{X_n\}$ is a MC.

$$\begin{aligned} p(i, j) &= P(X_{n+1} = j | X_n = i) \\ &= \binom{N}{j} \left(\frac{i}{N}\right)^j \left(1 - \frac{i}{N}\right)^{N-j} \\ &\quad (\text{binomial with } p = \frac{i}{N} \text{ of } N \text{ trials}) \end{aligned}$$

Q: What is the probability that the population fixates in the all A state?



Example (Wright-Fisher model)

Population of N genes that can be of two types: A or a . Population at time $n + 1$ is obtained by drawing with replacement from the population at time n . Let X_n be # of A alleles at time n . Then, $\{X_n\}$ is a MC.

$$\begin{aligned} p(i, j) &= P(X_{n+1} = j | X_n = i) \\ &= \binom{N}{j} \left(\frac{i}{N}\right)^j \left(1 - \frac{i}{N}\right)^{N-j} \\ &\quad (\text{binomial with } p = \frac{i}{N} \text{ of } N \text{ trials}) \end{aligned}$$

Q: What is the probability that the population fixates in the all A state? i.e. $X_n = N$?



南方科技大学
SOUTHERN UNIVERSITY OF SCIENCE AND TECHNOLOGY

We can also introduce mutation into the model:



We can also introduce mutation into the model: A is mutated to a with prob. u , and



We can also introduce mutation into the model: A is mutated to a with prob. u , and a mutated to A w/ prob. v .



We can also introduce mutation into the model: A is mutated to a with prob. u , and a mutated to A w/ prob. v . Then, if $X_n = i$, the prob. of getting A in one draw is



We can also introduce mutation into the model: A is mutated to a with prob. u , and a mutated to A w/ prob. v . Then, if $X_n = i$, the prob. of getting A in one draw is

$$\rho_i = \frac{i}{N}(1 - u) + \left(1 - \frac{i}{N}\right)v.$$



We can also introduce mutation into the model: A is mutated to a with prob. u , and a mutated to A w/ prob. v . Then, if $X_n = i$, the prob. of getting A in one draw is

$$\rho_i = \frac{i}{N}(1 - u) + \left(1 - \frac{i}{N}\right)v.$$

So,

$$p(i, j) = \binom{N}{j} \rho_i^j (1 - \rho_i)^{N-j}.$$



南方科技大学
SOUTHERN UNIVERSITY OF SCIENCE AND TECHNOLOGY

3.2 C-K equation



3.2 C-K equation

In last section, we introduced one-step transition matrix

$$p(i, j) = \mathbb{P}(X_{n+1} = j | X_n = i).$$

3.2 C-K equation

In last section, we introduced one-step transition matrix

$$p(i, j) = \mathbb{P}(X_{n+1} = j | X_n = i).$$

How about m -step transition probabilities? $m > 1$.

3.2 C-K equation

In last section, we introduced one-step transition matrix

$$p(i, j) = \mathbb{P}(X_{n+1} = j | X_n = i).$$

How about m -step transition probabilities? $m > 1$.

$$p^{(m)}(i, j) = \mathbb{P}(X_{n+m} = j | X_n = i).$$



Example (7)

Let $\{X_n\}$ be a MC (social mobility) with states $\{1, 2, 3\}$ (poor, middle, rich) and transition matrix

$$P = \begin{bmatrix} .7 & .2 & .1 \\ .3 & .5 & .2 \\ .2 & .4 & .4 \end{bmatrix}$$

Find $p^{(2)}(2, 1)$.



Example (7)

Let $\{X_n\}$ be a MC (social mobility) with states $\{1, 2, 3\}$ (poor, middle, rich) and transition matrix

$$P = \begin{bmatrix} .7 & .2 & .1 \\ .3 & .5 & .2 \\ .2 & .4 & .4 \end{bmatrix}$$

Find $p^{(2)}(2, 1)$.

Solution: Do it on board!



Example (7)

Let $\{X_n\}$ be a MC (social mobility) with states $\{1, 2, 3\}$ (poor, middle, rich) and transition matrix

$$P = \begin{bmatrix} .7 & .2 & .1 \\ .3 & .5 & .2 \\ .2 & .4 & .4 \end{bmatrix}$$

Find $p^{(2)}(2, 1)$.

Solution: Do it on board! Recall $\mathbb{P}(AB) = \mathbb{P}(A|B)\mathbb{P}(B)$. So



$$\begin{aligned} p^{(2)}(2, 1) &= \mathbb{P}(X_2 = 1 | X_0 = 2) \\ &= \sum_{j=1}^3 \mathbb{P}(X_2 = 1, X_1 = j | X_0 = 2) \\ &= \sum_{j=1}^3 \mathbb{P}(X_2 = 1 | X_1 = j, X_0 = 2) \mathbb{P}(X_1 = j | X_0 = 2) \\ &= \sum_{j=1}^3 \mathbb{P}(X_2 = 1 | X_1 = j) \mathbb{P}(X_1 = j | X_0 = 2) \\ &= \sum_{j=1}^3 p(2, j) p(j, 1) \\ &= .3 \times .7 + .5 \times .3 + .2 \times .2 = .4 \end{aligned}$$





Remark

If the transition matrix is P , then the 2-step transition matrix is P^2 , namely,

$$p^{(2)}(i, j) = \sum_k p(i, k)p(k, j).$$



Remark

If the transition matrix is P , then the 2-step transition matrix is P^2 , namely,

$$p^{(2)}(i, j) = \sum_k p(i, k)p(k, j).$$

Theorem (C-K equation)

The m -step transition matrix is P^m , namely, $P^{(m)} = P^m$.



Example 7 (Conti.) Find the 2-step transition matrix.



Example 7 (Conti.) Find the 2-step transition matrix.
Solution:



Example 7 (Conti.) Find the 2-step transition matrix.
Solution:

$$P^2 = \begin{bmatrix} .7 & .2 & .1 \\ .3 & .5 & .2 \\ .2 & .4 & .4 \end{bmatrix} \begin{bmatrix} .7 & .2 & .1 \\ .3 & .5 & .2 \\ .2 & .4 & .4 \end{bmatrix}$$



Example 7 (Conti.) Find the 2-step transition matrix.

Solution:

$$P^2 = \begin{bmatrix} .7 & .2 & .1 \\ .3 & .5 & .2 \\ .2 & .4 & .4 \end{bmatrix} \begin{bmatrix} .7 & .2 & .1 \\ .3 & .5 & .2 \\ .2 & .4 & .4 \end{bmatrix} = \begin{bmatrix} .57$$



Example 7 (Conti.) Find the 2-step transition matrix.

Solution:

$$P^2 = \begin{bmatrix} .7 & .2 & .1 \\ .3 & .5 & .2 \\ .2 & .4 & .4 \end{bmatrix} \begin{bmatrix} .7 & .2 & .1 \\ .3 & .5 & .2 \\ .2 & .4 & .4 \end{bmatrix} = \begin{bmatrix} .57 & .28 & .15 \\ .4 & .39 & .21 \\ .34 & .4 & .26 \end{bmatrix}.$$



3.3. Classification of states

3.3. Classification of states

Notation:

$$\mathbb{P}_x(A) = \mathbb{P}(A|X_0 = x) \quad \text{conditional prob. given } X_0 = x.$$

3.3. Classification of states

Notation:

$\mathbb{P}_x(A) = \mathbb{P}(A|X_0 = x)$ conditional prob. given $X_0 = x$.

\mathbb{E}_x = expectation using \mathbb{P}_x

3.3. Classification of states

Notation:

$$\mathbb{P}_x(A) = \mathbb{P}(A|X_0 = x) \quad \text{conditional prob. given } X_0 = x.$$

$$\mathbb{E}_x = \text{expectation using } \mathbb{P}_x$$

$$T_y = \inf\{n \geq 1 : X_n = y\} \quad \text{first time return to (or reach) } y$$

3.3. Classification of states

Notation:

$$\mathbb{P}_x(A) = \mathbb{P}(A|X_0 = x) \quad \text{conditional prob. given } X_0 = x.$$

\mathbb{E}_x = expectation using \mathbb{P}_x

$$T_y = \inf\{n \geq 1 : X_n = y\} \quad \text{first time return to (or reach) } y$$

Let

$$\rho_{xy} = \mathbb{P}_x(T_y < \infty)$$

3.3. Classification of states

Notation:

$$\mathbb{P}_x(A) = \mathbb{P}(A|X_0 = x) \quad \text{conditional prob. given } X_0 = x.$$

\mathbb{E}_x = expectation using \mathbb{P}_x

$$T_y = \inf\{n \geq 1 : X_n = y\} \quad \text{first time return to (or reach) } y$$

Let

$$\rho_{xy} = \mathbb{P}_x(T_y < \infty)$$

be the prob. of reaching (returning) y eventually while starting at x .

3.3. Classification of states

Notation:

$$\mathbb{P}_x(A) = \mathbb{P}(A|X_0 = x) \quad \text{conditional prob. given } X_0 = x.$$

\mathbb{E}_x = expectation using \mathbb{P}_x

$$T_y = \inf\{n \geq 1 : X_n = y\} \quad \text{first time return to (or reach) } y$$

Let

$$\rho_{xy} = \mathbb{P}_x(T_y < \infty)$$

be the prob. of reaching (returning) y eventually while starting at x .

Note the book use notation: $f_{i,j}$ or f_i .



Note that, the prob. of the chain returning to y twice is ρ_{yy}^2 because



Note that, the prob. of the chain returning to y twice is ρ_{yy}^2 because once it returns to y , the chain starts at y again (Markov property).



Note that, the prob. of the chain returning to y twice is ρ_{yy}^2 because once it returns to y , the chain starts at y again (Markov property).

Let T_y^k be the k th time MC returns to y ,



Note that, the prob. of the chain returning to y twice is ρ_{yy}^2 because once it returns to y , the chain starts at y again (Markov property).

Let T_y^k be the k th time MC returns to y , namely,

$$T_y^1 = T_y, \quad T_y^k = \inf\{n > T_y^{k-1} : X_n = y\}.$$



Note that, the prob. of the chain returning to y twice is ρ_{yy}^2 because once it returns to y , the chain starts at y again (Markov property).

Let T_y^k be the k th time MC returns to y , namely,

$$T_y^1 = T_y, \quad T_y^k = \inf\{n > T_y^{k-1} : X_n = y\}.$$

Then,

$$\mathbb{P}_y(T_y^k < \infty) = \rho_{yy}^k.$$



Definition

i) y is *transient* if $\rho_{yy} < 1$.



Definition

- i) y is *transient* if $\rho_{yy} < 1$.
- ii) y is *recurrent* if $\rho_{yy} = 1$.



南方科技大学
SOUTHERN UNIVERSITY OF SCIENCE AND TECHNOLOGY

Notation: i.o. stands for “infinitely often”.



Notation: i.o. stands for “infinitely often”.

Proposition

i) If y is transient, then

$$\mathbb{P}_y(X_n = y, \text{ i.o.}) = 0.$$



Notation: i.o. stands for “infinitely often”.

Proposition

i) If y is transient, then

$$\mathbb{P}_y(X_n = y, \text{ i.o.}) = 0.$$

ii) If y is recurrent, then

$$\mathbb{P}_y(X_n = y, \text{ i.o.}) = 1.$$



Notation: i.o. stands for “infinitely often”.

Proposition

i) If y is transient, then

$$\mathbb{P}_y(X_n = y, \text{ i.o.}) = 0.$$

ii) If y is recurrent, then

$$\mathbb{P}_y(X_n = y, \text{ i.o.}) = 1.$$

Proof: Do it on board!

i)

$$\begin{aligned}\mathbb{P}_y(X_n = y, i.o.) &= \mathbb{P}_y(T_y^k < \infty, \forall k) \\ &\leq \mathbb{P}_y(T_y^n < \infty) \quad \forall n.\end{aligned}$$

Thus,

$$\mathbb{P}_y(X_n = y, i.o.) \leq \rho_{yy}^n \rightarrow 0.$$

ii)

$$\begin{aligned}1 - \mathbb{P}_y(X_n = y, i.o.) &= \mathbb{P}_y(\exists k, T_y^k = \infty) \\ &\leq \sum_k \mathbb{P}_y(T_y^k = \infty) \\ &\leq \sum_k (1 - \rho_{yy}^k) = 0.\end{aligned}$$





Example (Gambler's ruin $N = 4$)

$$P = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ .6 & 0 & .4 & 0 & 0 \\ 0 & .6 & 0 & .4 & 0 \\ 0 & 0 & .6 & 0 & .4 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$



Example (Gambler's ruin $N = 4$)

$$P = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ .6 & 0 & .4 & 0 & 0 \\ 0 & .6 & 0 & .4 & 0 \\ 0 & 0 & .6 & 0 & .4 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Classify all 5 states.



南方科技大学
SOUTHERN UNIVERSITY OF SCIENCE AND TECHNOLOGY

Solution: Do it on board!

Solution: Do it on board!

As 0 and 4 are absorbing state, they are clearly recurrent. e.g.

$$\rho_{00} = \mathbb{P}_0(T_0 < \infty) = \mathbb{P}_0(1 < \infty) = 1.$$

Claim: 1, 2, 3 are transient. e.g.

$$\mathbb{P}_1(T_1 = \infty) \geq p(1, 0) = .6$$

so,

$$\rho_{11} = \mathbb{P}_1(T_1 < \infty) \leq 1 - .6 = .4 < 1.$$

The MC gets stuck in either 0 or 4 eventually.





Example (Social mobility)

$$P = \begin{bmatrix} .7 & .2 & .1 \\ .3 & .5 & .2 \\ .2 & .4 & .4 \end{bmatrix}$$



Example (Social mobility)

$$P = \begin{bmatrix} .7 & .2 & .1 \\ .3 & .5 & .2 \\ .2 & .4 & .4 \end{bmatrix}$$

All 3 states are recurrent.



Example (Social mobility)

$$P = \begin{bmatrix} .7 & .2 & .1 \\ .3 & .5 & .2 \\ .2 & .4 & .4 \end{bmatrix}$$

All 3 states are recurrent.

Proof: Do it later!



南方科技大学
SOUTHERN UNIVERSITY OF SCIENCE AND TECHNOLOGY

Now, we will give some general results about state classification.



Now, we will give some general results about state classification.

Definition

We say that x communicates with y , write $x \rightarrow y$, if

$$\rho_{xy} \equiv \mathbb{P}_x(T_y < \infty) > 0.$$



Lemma

If $x \rightarrow y$ and $y \rightarrow z$, then $x \rightarrow z$.



Lemma

If $x \rightarrow y$ and $y \rightarrow z$, then $x \rightarrow z$.

Proof: Do it on board!



Lemma

If $x \rightarrow y$ and $y \rightarrow z$, then $x \rightarrow z$.

Proof: Do it on board!

Since $x \rightarrow y$, $\exists m$ s.t. $p^m(x, y) > 0$. Similarly, $\exists n$, $p^n(y, z) > 0$.

Then,

$$p^{m+n}(x, z) \geq p^m(x, y)p^n(y, z) > 0.$$

Hence, $x \rightarrow z$. □



Theorem (14)

If $\rho_{xy} > 0$ and $\rho_{yx} < 1$, then x is transient.



Theorem (14)

If $\rho_{xy} > 0$ and $\rho_{yx} < 1$, then x is transient.

Proof: Do it on board!



Theorem (14)

If $\rho_{xy} > 0$ and $\rho_{yx} < 1$, then x is transient.

Proof: Do it on board!

Let k be the smallest s.t. $p^k(x, y) > 0$. Then,

$$0 < p^k(x, y)(1 - \rho_{yx}) \leq \mathbb{P}_x(T_x = \infty) = 1 - \rho_{xx}.$$

So, $\rho_{xx} < 1$ and hence, x is transient. □



Theorem (14)

If $\rho_{xy} > 0$ and $\rho_{yx} < 1$, then x is transient.

Proof: Do it on board!

Let k be the smallest s.t. $p^k(x, y) > 0$. Then,

$$0 < p^k(x, y)(1 - \rho_{yx}) \leq \mathbb{P}_x(T_x = \infty) = 1 - \rho_{xx}.$$

So, $\rho_{xx} < 1$ and hence, x is transient. □

Corollary (1.5)

If x recurrent and $\rho_{xy} > 0$, then $\rho_{yx} = 1$.



Example

A seven-state chain

$$P = \begin{bmatrix} .7 & 0 & 0 & 0 & .3 & 0 & 0 \\ .1 & .2 & .3 & .4 & 0 & 0 & 0 \\ 0 & 0 & .5 & .3 & .2 & 0 & 0 \\ 0 & 0 & 0 & .5 & 0 & .5 & 0 \\ .6 & 0 & 0 & 0 & .4 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & .2 & .8 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \end{bmatrix}.$$



Example

A seven-state chain

$$P = \begin{bmatrix} .7 & 0 & 0 & 0 & .3 & 0 & 0 \\ .1 & .2 & .3 & .4 & 0 & 0 & 0 \\ 0 & 0 & .5 & .3 & .2 & 0 & 0 \\ 0 & 0 & 0 & .5 & 0 & .5 & 0 \\ .6 & 0 & 0 & 0 & .4 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & .2 & .8 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \end{bmatrix}.$$

Classify all states.



Example

A seven-state chain


$$P = \begin{bmatrix} .7 & 0 & 0 & 0 & .3 & 0 & 0 \\ .1 & .2 & .3 & .4 & 0 & 0 & 0 \\ 0 & 0 & .5 & .3 & .2 & 0 & 0 \\ 0 & 0 & 0 & .5 & 0 & .5 & 0 \\ .6 & 0 & 0 & 0 & .4 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & .2 & .8 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \end{bmatrix}.$$

Classify all states.

Solution: Do it on board!



To identify recurrent and transient states, we draw a graph of 1-step communication.



$\rho_{21} > 0$ but $\rho_{12} = 0 < 1$, so 2 is transient (by Theorem 14). Similarly, 3 is transient.

We will prove all other states are recurrent later.





Definition

A set A is *closed* if it is impossible to get out.



Definition

A set A is *closed* if it is impossible to get out.

Example (Conti.) $\{1, 5\}$,



Definition

A set A is *closed* if it is impossible to get out.

Example (Conti.) $\{1, 5\}, \quad \{4, 6, 7\},$



Definition

A set A is *closed* if it is impossible to get out.

Example (Conti.) $\{1, 5\}$, $\{4, 6, 7\}$, $\{1, 5, 4, 6, 7\}$,



Definition

A set A is *closed* if it is impossible to get out.

Example (Conti.) $\{1, 5\}$, $\{4, 6, 7\}$, $\{1, 5, 4, 6, 7\}$,
 $\{1, 2, 3, 4, 5, 6, 7\}$ are all closed.



Definition

A set B is *irreducible* if $\forall i, j \in B, i \rightarrow j$.



Definition

A set B is *irreducible* if $\forall i, j \in B, i \rightarrow j$.

Example (Conti.) $\{1, 5\}, \{4, 6, 7\}$ are irreducible.



Definition

A set B is *irreducible* if $\forall i, j \in B, i \rightarrow j$.

Example (Conti.) $\{1, 5\}, \{4, 6, 7\}$ are irreducible. $\{1, 5, 4, 6, 7\},$
 $\{1, 2, 3, 4, 5, 6, 7\}$ are not.



Theorem

If C is a finite closed irreducible set, then all states in C are recurrent.



Theorem

If C is a finite closed irreducible set, then all states in C are recurrent.

Proof:



Theorem

If C is a finite closed irreducible set, then all states in C are recurrent.

Proof: Later.



Theorem

If C is a finite closed irreducible set, then all states in C are recurrent.

Proof: Later.

Example (conti.) Seven-state, $\{1, 5\}$ is closed,



Theorem

If C is a finite closed irreducible set, then all states in C are recurrent.

Proof: Later.

Example (conti.) Seven-state, $\{1, 5\}$ is closed, irreducible.
So,



Theorem

If C is a finite closed irreducible set, then all states in C are recurrent.

Proof: Later.

Example (conti.) Seven-state, $\{1, 5\}$ is closed, irreducible.
So, 1 and 5 are recurrent.



Theorem

If the state space S is finite, then S can be written as a disjoint union

$$S = T \cup R_1 \cup \cdots \cup R_k,$$

where T is a set of transient states and R_i , $1 \leq i \leq k$, are closed irreducible sets of recurrent states.



Theorem

If the state space S is finite, then S can be written as a disjoint union

$$S = T \cup R_1 \cup \cdots \cup R_k,$$

where T is a set of transient states and R_i , $1 \leq i \leq k$, are closed irreducible sets of recurrent states.

Proof: Do it on board!



Proof: Let T be the set of all transient states. Then, $S \setminus T$ contains recurrent states. Let $x \in S \setminus T$ and

$$R_1 = \{y \in S \setminus T : x \rightarrow y\}.$$

$\forall y \in R_1$, since $\rho_{xy} > 0$ and x recurrent, $\rho_{yx} = 1 > 0$. Hence, $y \rightarrow x$.

For any $z \in R_1$, then $y \rightarrow x \rightarrow z$. Thus, R_1 is irreducible. It is clear that R_1 is closed.

Replacing $S \setminus T$ by $S \setminus (T \cup R_1)$, we can construct R_2 .

Eventually, we obtain the desired decomposition. □



Recall

$$T_y^k = \inf\{n > T_y^{k-1} : X_n = y\} \text{ and } \rho_{xy} = \mathbb{P}_x(T_y < \infty).$$

Recall

$$T_y^k = \inf\{n > T_y^{k-1} : X_n = y\} \text{ and } \rho_{xy} = \mathbb{P}_x(T_y < \infty).$$

Then

$$\mathbb{P}_x(T_y^k < \infty) = \rho_{xy}\rho_{yy}^{k-1}.$$

Recall

$$T_y^k = \inf\{n > T_y^{k-1} : X_n = y\} \text{ and } \rho_{xy} = \mathbb{P}_x(T_y < \infty).$$

Then

$$\mathbb{P}_x(T_y^k < \infty) = \rho_{xy}\rho_{yy}^{k-1}.$$

Let

$$N(y) = \# \text{ of visits to } y \text{ at } n \geq 1.$$



Lemma (23)

$$\mathbb{E}_x N(y) = \frac{\rho_{xy}}{1 - \rho_{yy}}.$$



Lemma (23)

$$\mathbb{E}_x N(y) = \frac{\rho_{xy}}{1 - \rho_{yy}}.$$

Proof: Do it on board!



Lemma (23)

$$\mathbb{E}_x N(y) = \frac{\rho_{xy}}{1 - \rho_{yy}}.$$

Proof: Do it on board! Note that for \mathbb{Z}_+ -valued r.v. X ,

$$\mathbb{E}X = \sum_{k=1}^{\infty} \mathbb{P}(X \geq k).$$

Thus,



$$\begin{aligned}\mathbb{E}_x N(y) &= \sum_{k=1}^{\infty} \mathbb{P}_x(N(y) \geq k) \\ &= \sum_{k=1}^{\infty} \mathbb{P}_x(T_y^k < \infty) \\ &= \sum_{k=1}^{\infty} \rho_{xy} \rho_{yy}^{k-1} \\ &= \frac{\rho_{xy}}{1 - \rho_{yy}}.\end{aligned}$$





Lemma

$$\mathbb{E}_x N(y) = \sum_{n=1}^{\infty} p^n(x, y).$$



Lemma

$$\mathbb{E}_x N(y) = \sum_{n=1}^{\infty} p^n(x, y).$$

Proof: Do it on board!



Lemma

$$\mathbb{E}_x N(y) = \sum_{n=1}^{\infty} p^n(x, y).$$

Proof: Do it on board!

As

$$N(y) = \sum_{n=1}^{\infty} 1_{X_n=y},$$

we have

$$\mathbb{E}_x N(y) = \mathbb{E}_x \sum_{n=1}^{\infty} 1_{X_n=y} = \sum_{n=1}^{\infty} \mathbb{P}_x(X_n = y) = \sum_{n=1}^{\infty} p^n(x, y).$$





Theorem

y is recurrent iff

$$\sum_{n=1}^{\infty} p^n(y, y) = \mathbb{E}_y N(y) = \infty.$$



Theorem

y is recurrent iff

$$\sum_{n=1}^{\infty} p^n(y, y) = \mathbb{E}_y N(y) = \infty.$$

Proof: Do it on board!



Theorem

y is recurrent iff

$$\sum_{n=1}^{\infty} p^n(y, y) = \mathbb{E}_y N(y) = \infty.$$

Proof: Do it on board! “ \Rightarrow ” Since $X_n = y$ i.o., $N(y) = \infty$.
Thus, $\mathbb{E}_y N(y) = \infty$.
“ \Leftarrow ” By Lemma 20 ,

$$\frac{\rho_{yy}}{1 - \rho_{yy}} = \mathbb{E}_y N(y) = \infty.$$

$$\rho_{yy} = 1.$$





Lemma

If x is recurrent and $x \rightarrow y$, then y is recurrent.



Lemma

If x is recurrent and $x \rightarrow y$, then y is recurrent.

Proof: Do it on board!



Lemma

If x is recurrent and $x \rightarrow y$, then y is recurrent.

Proof: Do it on board! By Cor. 1.4, $\rho_{yx} = 1 > 0$. Let j, ℓ be s.t.

$$p^j(y, x) > 0 \text{ and } p^\ell(x, y) > 0.$$

Then,

$$\begin{aligned} \sum_{n=1}^{\infty} p^n(y, y) &\geq \sum_{k=0}^{\infty} p^{j+k+\ell}(y, y) \\ &\geq \sum_{k=0}^{\infty} p^j(y, x) p^k(x, x) p^\ell(x, y) \\ &= p^j(y, x) p^\ell(x, y) \mathbb{E}_x N(x) = \infty. \end{aligned}$$

So, y is recurrent. □



Lemma

In a finite closed set there has to be at least one recurrent state.



Lemma

In a finite closed set there has to be at least one recurrent state.

Proof: Do it on board!



Lemma

In a finite closed set there has to be at least one recurrent state.

Proof: Do it on board! Suppose all states in the closed set C are transient. Then, $\mathbb{E}_x N(y) < \infty, \forall x, y \in C$. Thus

$$\begin{aligned} \infty &> \sum_{y \in C} \mathbb{E}_x N(y) = \sum_{y \in C} \sum_{n=1}^{\infty} p^n(x, y) \\ &= \sum_{n=1}^{\infty} \sum_{y \in C} p^n(x, y) = \sum_{n=1}^{\infty} 1 = \infty. \end{aligned}$$





Recall

Theorem (18)

If C is a finite closed irreducible set, then all states in C are recurrent.



Recall

Theorem (18)

If C is a finite closed irreducible set, then all states in C are recurrent.

Proof of Theorem 18: Do it on board!



Recall

Theorem (18)

If C is a finite closed irreducible set, then all states in C are recurrent.

Proof of Theorem 18: Do it on board! Let $x \in C$ be recurrent. Let $y \in C$. Since C is irreducible, $x \rightarrow y$. Thus, y is recurrent. Thus, all states in C are recurrent. \square

HW: Ch4, 5, 8, 13, 14.