

Step-1

Consider the following 3 by 3 Vandermonde determinant:

$$\det \begin{bmatrix} 1 & a & a^2 \\ 1 & b & b^2 \\ 1 & c & c^2 \end{bmatrix} = (b-a)(c-a)(c-b)$$

The objective is to use the row operations to verify that the 3 by 3 Vandermonde determinant.

Use row operations and properties of the determinant,

$$\begin{vmatrix} 1 & a & a^2 \\ 1 & b & b^2 \\ 1 & c & c^2 \end{vmatrix} \xrightarrow{\substack{R_2 \rightarrow R_2 - R_1 \\ R_3 \rightarrow R_3 - R_1}} \begin{vmatrix} 1 & a & a^2 \\ 0 & b-a & b^2-a^2 \\ 0 & c-a & c^2-a^2 \end{vmatrix}$$

By linearity of the determinant, we can pull out a factor of $(b-a)$ from row 2 and a factor of $(c-a)$ from row 3.

$$\begin{vmatrix} 1 & a & a^2 \\ 0 & b-a & b^2-a^2 \\ 0 & c-a & c^2-a^2 \end{vmatrix} = (b-a)(c-a) \begin{vmatrix} 1 & a & a^2 \\ 0 & 1 & b+a \\ 0 & 1 & c+a \end{vmatrix}$$

Step-2

Use the cofactor formula or alternatively use row transformations.

From the cofactor formula, the determinant of the matrix is,

$$\begin{aligned} \begin{vmatrix} 1 & a & a^2 \\ 0 & b-a & b^2-a^2 \\ 0 & c-a & c^2-a^2 \end{vmatrix} &= (b-a)(c-a) \begin{vmatrix} 1 & a & a^2 \\ 0 & 1 & b+a \\ 0 & 1 & c+a \end{vmatrix} \\ &= (b-a)(c-a)(1) \begin{vmatrix} 1 & b+a \\ 1 & c+a \end{vmatrix} \\ &= (b-a)(c-a)(c+a-(b+a)) \\ &= (b-a)(c-a)(c-b). \end{aligned}$$