Step-1

Given that
$$b = \begin{bmatrix} 0.217 \\ 0.254 \end{bmatrix}$$
, $A = \begin{bmatrix} 0.780 & 0.563 \\ 0.913 & 0.659 \end{bmatrix}$, $y = \begin{bmatrix} 0.341 \\ -0.087 \end{bmatrix}$, $z = \begin{bmatrix} 0.999 \\ -1.0 \end{bmatrix}$

We have to compute b - Ay and b - Az

Step-2

Now

$$Ay = \begin{bmatrix} 0.780 & 0.563 \\ 0.913 & 0.659 \end{bmatrix} \begin{bmatrix} 0.341 \\ -0.087 \end{bmatrix}$$
$$= \begin{bmatrix} 0.780(0.341) + (0.563)(-0.087) \\ 0.913(0.341) + (0.659)(-0.087) \end{bmatrix}$$
$$= \begin{bmatrix} 0.21699 \\ 0.254 \end{bmatrix}$$

Therefore,

$$b - Ay = \begin{bmatrix} 0.217 \\ 0.254 \end{bmatrix} - \begin{bmatrix} 0.216999 \\ 0.254 \end{bmatrix}$$
$$= \begin{bmatrix} 0.000001 \\ 0 \end{bmatrix}_{\hat{\mathbf{a}} \in [\hat{\mathbf{a}} \in [1])}$$
$$b - Ay = \begin{bmatrix} 0.000001 \\ 0 \end{bmatrix}$$

Hence

Step-3

Now

$$Az = \begin{bmatrix} 0.780 & 0.563 \\ 0.913 & 0.659 \end{bmatrix} \begin{bmatrix} 0.999 \\ -1.0 \end{bmatrix}$$
$$= \begin{bmatrix} 0.780(0.999) + (0.563)(-1.0) \\ 0.913(0.999) + (0.659)(-1.0) \end{bmatrix}$$
$$= \begin{bmatrix} 0.216220 \\ 0.253087 \end{bmatrix}$$

Step-4

Therefore,

$$b - Az = \begin{bmatrix} 0.217 \\ 0.254 \end{bmatrix} - \begin{bmatrix} 0.21622 \\ 0.253087 \end{bmatrix}$$
$$= \begin{bmatrix} 0.00078 \\ 0.000913 \end{bmatrix} \hat{\mathbf{a}} \in \hat{\mathbf{a}} \in [1]$$
$$b - Az = \begin{bmatrix} 0.00078 \\ 0.000913 \end{bmatrix}$$

Hence

Step-5

If Ax = b is a non homogeneous system of equations in two variables whose solutions are y and z, then we follow that Ay = b, Az = b.

Consequently, b - Ay and b - Az

In view of (1) and (2), we see that y is the better solution than z.

Step-6

On the other hand, comparing the solutions y and z to the true solution $x = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$, z is the better solution than y.

The corresponding residuals $x - y = \delta y = \begin{bmatrix} 0.659 \\ -0.913 \end{bmatrix}$ and $x - z = \delta z = \begin{bmatrix} 0.001 \\ 0 \end{bmatrix}$ confirms this.