

## Step-1

The number  $\lambda$  is an eigen value of  $A$  if and only if  $A - \lambda I$  is singular

In other words,  $\det(A - \lambda I) = 0$

This is the characteristic equation. Each  $\lambda$  is associated with eigen vector  $x$  such that

$$Ax = \lambda x.$$

## Step-2

In view of this definition, let us consider  $A = \begin{bmatrix} 3 & 4 \\ 4 & -3 \end{bmatrix}$

The characteristic equation of this matrix is  $\det(A - \lambda I) = \begin{vmatrix} 3-\lambda & 4 \\ 4 & -3-\lambda \end{vmatrix} = 0$

$$\Rightarrow \lambda^2 - 9 - 16 = 0$$

$\Rightarrow \lambda_1 = -5, \lambda_2 = 5$  are the roots of the characteristic equation or the eigen values of the given matrix.

## Step-3

The eigen vector corresponding to the eigen value  $\lambda_1 = -5$  satisfies  $(A - \lambda_1 I)x_1 = 0$

$$\Rightarrow \begin{bmatrix} 8 & 4 \\ 4 & 2 \end{bmatrix} \begin{bmatrix} t_1 \\ t_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Using the row operations, we reduce this as

$$R_2 \rightarrow 2R_2 - R_1, R_1 / 4 \Rightarrow \begin{bmatrix} 2 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} t_1 \\ t_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\Rightarrow 2t_1 + t_2 = 0$$

$$t_2 = -2t_1$$

When  $t_1 = 1$ , we get  $t_2 = -2$  and thus, the solution set is  $x_1 = \begin{bmatrix} 1 \\ -2 \end{bmatrix}$  is the eigen vector corresponding to  $\lambda_1 = -5$

## Step-4

Similarly, using  $\lambda_2 = 5$ , we get  $(A - \lambda_2 I)x_2 = 0$  as  $\begin{bmatrix} -2 & 4 \\ 4 & -8 \end{bmatrix} \begin{bmatrix} t_1 \\ t_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$

$$R_2 \rightarrow R_2 + 2R_1, R_1 / -2 \Rightarrow \begin{bmatrix} 1 & -2 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} t_1 \\ t_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\Rightarrow t_1 - 2t_2 = 0$$

When  $t_2 = 1$ , we get  $t_1 = 2$  and thus, the solution set is  $x_2 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$  is the eigen vector corresponding to  $\lambda_2 = 5$

## Step-5

The characteristic equation of  $A = \begin{bmatrix} a & b \\ b & a \end{bmatrix}$  is  $\det(A - \lambda I) = \begin{vmatrix} a - \lambda & b \\ b & a - \lambda \end{vmatrix} = 0$

$$\Rightarrow (\lambda - a)^2 - b^2 = 0$$

$$\Rightarrow \lambda^2 - 2a\lambda + (a^2 - b^2) = 0$$

$$\Rightarrow \lambda = \frac{2a + \sqrt{4a^2 - 4(a^2 - b^2)}}{2}, \frac{2a - \sqrt{4a^2 - 4(a^2 - b^2)}}{2}$$

$$\Rightarrow \lambda_1 = a + b, \lambda_2 = a - b$$

## Step-6

Let the eigen vector corresponding to  $\lambda_1$  is  $x_1$

Then  $x_1$  satisfies the equation  $(A - \lambda_1 I)x_1 = 0$

$$\Rightarrow \begin{bmatrix} -b & b \\ b & -b \end{bmatrix} \begin{bmatrix} t_1 \\ t_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Applying row operation  $R_2 \rightarrow R_2 + R_1, R_2 / -b$ , we get  $\begin{bmatrix} 1 & -1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} t_1 \\ t_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$

$$\Rightarrow t_1 - t_2 = 0$$

Putting  $t_1 = 1$ , we get  $t_2 = 1$  and thus, the respective eigen vector is  $x_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$

## Step-7

Similarly,

Let the eigen vector corresponding to  $\lambda_2$  is  $x_2$

Then  $x_2$  satisfies the equation  $(A - \lambda_2 I)x_2 = 0$

$$\Rightarrow \begin{bmatrix} b & b \\ b & b \end{bmatrix} \begin{bmatrix} t_1 \\ t_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Applying row operation  $R_2 \rightarrow R_2 - R_1, R_2 / b$ , we get  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} t_1 \\ t_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$

$$\Rightarrow t_1 + t_2 = 0$$

Putting  $t_1 = 1$ , we get  $t_2 = -1$  and thus, the respective eigen vector is  $x_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$