1. A town has an initial population of 70000. The town's population for the next 10 years is provided below. Which type of function would be most appropriate to model the town's population?

Year	1	2	3	4	5	6	7	8	9
Pop.	69944	69898	69852	69806	69744	69698	69652	69606	69544

- A. Non-Linear Power
- B. Logarithmic
- C. Exponential
- D. Linear
- E. None of the above
- 2. For the information below, construct a linear model that describes the total time T spent on the path in terms of the distance of a particular part of the path if we know that the time spent on each path was equal.

A bicyclist is training for a race on a hilly path. Their bike keeps track of their speed at any time, but not the distance traveled. Their speed traveling up a hill is 4 mph, 8 mph when traveling down a hill, and 6 mph when traveling along a flat portion.

- A. 18.000*D*
- B. 192.000*D*
- C. 0.542D
- D. The model can be found with the information provided, but isn't options 1-3
- E. The model cannot be found with the information provided.
- 3. What is the **best** way to describe the domain of the scenario below?

Bridges on highways often have expansion joints, which are small gaps in the roadway between one bridge section and the next. The gaps are put there so the bridge will have room to expand when the weather gets hot. Assume the gap width varies constantly with the temperature. Suppose a bridge has a gap of 1.3 cm when the temperature is 22

degrees C and that the gap narrows to 0.9 cm when the temperature warms to 30 degrees C.

- A. Subset of the Rational numbers
- B. Subset of the Integers
- C. There is no restricted domain in this scenario
- D. Proper subset of the Real numbers
- E. Subset of the Natural numbers
- 4. For the information provided below, construct a linear model that describes her total budget, B, as a function of the number of months, x she is at UF.

Aubrey is a college student going into her first year at UF. She will receive Bright Futures, which covers her tuition plus a \$1000 educational expense each year. Before college, Aubrey saved up \$8000. She knows she will need to pay \$1200 in rent a month, \$50 for food a week, and \$64 in other weekly expenses.

A.
$$B(x) = 7686x$$

B.
$$B(x) = 7344x$$

C.
$$B(x) = 9000 - 1314x$$

D.
$$B(x) = 9000 - 1656x$$

- E. None of the above.
- 5. Using the situation below, construct a linear model that describes the cost of the coffee beans C(h) in terms of the weight of the low-quality coffee beans h.

Veronica needs to prepare 200 of blended coffee beans selling for \$5.09 per pound. She has a high-quality bean that sells for \$5.71 a pound and a low-quality bean that sells for \$2.86 a pound.

A.
$$C(h) = 2.85h + 572.00$$

- B. C(h) = 2.86h
- C. C(h) = 4.29h
- D. C(h) = -2.85h + 1142.00
- E. None of the above.
- 6. For the information provided below, construct a linear model that describes her total budget, B, as a function of the number of months, x she is at UF.

Aubrey is a college student going into her first year at UF. She will receive Bright Futures, which covers her tuition plus a \$1000 educational expense each year. Before college, Aubrey saved up \$5000. She knows she will need to pay \$800 in rent a month, \$80 for food a week, and \$56 in other weekly expenses.

- A. B(x) = 5000x + 1000
- B. B(x) = 6000 1344x
- C. B(x) = 6000 936x
- D. B(x) = 1000x + 5000
- E. None of the above.
- 7. A town has an initial population of 40000. The town's population for the next 10 years is provided below. Which type of function would be most appropriate to model the town's population?

Year	1	2	3	4	5	6	7	8	9
Pop.	39947	39897	39855	39805	39747	39697	39655	39605	39547

- A. Logarithmic
- B. Linear
- C. Exponential
- D. Non-Linear Power
- E. None of the above

8. For the information provided below, construct a linear model that describes the total distance of the path, D, in terms of the time spent on a particular path if we know that the time spent on each path was equal.

A bicyclist is training for a race on a hilly path. Their bike keeps track of their speed at any time, but not the distance traveled. Their speed traveling up a hill is 5 mph, 9 mph when traveling down a hill, and 6 mph when traveling along a flat portion.

- A. 20t
- B. 0.478t
- C. 270t
- D. The model can be found with the information provided, but isn't options 1-3
- E. The model cannot be found with the information provided.
- 9. Using the situation below, construct a linear model that describes the cost of the coffee beans C(h) in terms of the weight of the low-quality coffee beans h.

Veronica needs to prepare 150 of blended coffee beans selling for \$5.09 per pound. She has a high-quality bean that sells for \$5.83 a pound and a low-quality bean that sells for \$3.66 a pound.

A.
$$C(h) = 3.66h$$

B.
$$C(h) = 4.75h$$

C.
$$C(h) = -2.17h + 874.50$$

D.
$$C(h) = 2.17h + 549.00$$

- E. None of the above.
- 10. What is the **best** way to describe the domain of the scenario below?

Veronica needs to prepare 170 lbs of blended coffee beans to sell for \$4.71 per pound. She has a high-quality bean that sells for \$6.00 a

Version C

pound and a low-quality been that sells for \$3.25 a pound.

- A. Subset of the Rational numbers
- B. Subset of the Natural numbers
- C. There is no restricted domain in this scenario
- D. Proper subset of the Real numbers
- E. Subset of the Integers