This key should allow you to understand why you choose the option you did (beyond just getting a question right or wrong). More instructions on how to use this key can be found here.

If you have a suggestion to make the keys better, please fill out the short survey here.

Note: This key is auto-generated and may contain issues and/or errors. The keys are reviewed after each exam to ensure grading is done accurately. If there are issues (like duplicate options), they are noted in the offline gradebook. The keys are a work-in-progress to give students as many resources to improve as possible.

1. Construct the lowest-degree polynomial given the zeros below. Then, choose the intervals that contain the coefficients of the polynomial in the form $x^3 + bx^2 + cx + d$.

$$-5 + 4i$$
 and 1

The solution is $x^3 + 9x^2 + 31x - 41$, which is option D.

- A. $b \in [0, 8], c \in [-9, -1]$, and $d \in [-1, 6]$ $x^3 + x^2 - 5x + 4$, which corresponds to multiplying out (x - 4)(x - 1).
- B. $b \in [0,8], c \in [0,10]$, and $d \in [-6,2]$ $x^3 + x^2 + 4x 5$, which corresponds to multiplying out (x+5)(x-1).
- C. $b \in [-14, -8], c \in [30, 39]$, and $d \in [40, 48]$ $x^3 - 9x^2 + 31x + 41$, which corresponds to multiplying out (x - (-5 + 4i))(x - (-5 - 4i))(x + 1).
- D. $b \in [7, 15], c \in [30, 39]$, and $d \in [-46, -29]$ * $x^3 + 9x^2 + 31x - 41$, which is the correct option.
- E. None of the above.

This corresponds to making an unanticipated error or not understanding how to use nonreal complex numbers to create the lowest-degree polynomial. If you chose this and are not sure what you did wrong, please contact the coordinator for help.

General Comment: Remember that the conjugate of a + bi is a - bi. Since these zeros always come in pairs, we need to multiply out (x - (-5 + 4i))(x - (-5 - 4i))(x - (1)).

2. Construct the lowest-degree polynomial given the zeros below. Then, choose the intervals that contain the coefficients of the polynomial in the form $ax^3 + bx^2 + cx + d$.

$$\frac{1}{4}, \frac{2}{3}, \text{ and } -7$$

The solution is $12x^3 + 73x^2 - 75x + 14$, which is option D.

- A. $a \in [3, 16], b \in [70, 76], c \in [-84, -73],$ and $d \in [-18, -11]$ $12x^3 + 73x^2 - 75x - 14$, which corresponds to multiplying everything correctly except the constant
- B. $a \in [3, 16], b \in [92, 96], c \in [72, 87], \text{ and } d \in [8, 15]$ $12x^3 + 95x^2 + 79x + 14, \text{ which corresponds to multiplying out } (4x + 1)(3x + 2)(x + 7).$
- C. $a \in [3, 16], b \in [78, 85], c \in [-44, -33], \text{ and } d \in [-18, -11]$ $12x^3 + 79x^2 - 37x - 14$, which corresponds to multiplying out (4x + 1)(3x - 2)(x + 7).

2107-1615 test

D. $a \in [3, 16], b \in [70, 76], c \in [-84, -73], \text{ and } d \in [8, 15]$

* $12x^3 + 73x^2 - 75x + 14$, which is the correct option.

E. $a \in [3, 16], b \in [-76, -67], c \in [-84, -73], \text{ and } d \in [-18, -11]$


 $12x^3 - 73x^2 - 75x - 14$, which corresponds to multiplying out (4x + 1)(3x + 2)(x - 7).


General Comment: To construct the lowest-degree polynomial, you want to multiply out (4x - 1)(3x - 2)(x + 7)


3. Describe the zero behavior of the zero x=3 of the polynomial below.

$$f(x) = -9(x-6)^5(x+6)^4(x+3)^{11}(x-3)^8$$

The solution is the graph below, which is option C.









В.

2107-1615 test






E. None of the above.


General Comment: You will need to sketch the entire graph, then zoom in on the zero the question asks about.

4. Describe the zero behavior of the zero x = -8 of the polynomial below.


$$f(x) = -8(x-6)^{7}(x+6)^{3}(x+8)^{8}(x-8)^{7}$$


The solution is the graph below, which is option C.





A.






D.

E. None of the above.

General Comment: You will need to sketch the entire graph, then zoom in on the zero the question asks about.

5. Which of the following equations could be of the graph presented below?



The solution is $-8x^9(x+4)^9(x-3)^7$, which is option C.

A.
$$-17x^9(x+4)^6(x-3)^{11}$$

The factor -4 should have been an odd power.

B.
$$-19x^{10}(x+4)^6(x-3)^7$$

The factors -4 and 0 have have been odd power.

C.
$$-8x^9(x+4)^9(x-3)^7$$

* This is the correct option.

D.
$$13x^{11}(x+4)^7(x-3)^7$$

This corresponds to the leading coefficient being the opposite value than it should be.

E.
$$16x^7(x+4)^{10}(x-3)^9$$


The factor (x + 4) should have an odd power and the leading coefficient should be the opposite sign.

General Comment: General Comments: Draw the x-axis to determine which zeros are touching (and so have even multiplicity) or cross (and have odd multiplicity).


6. Describe the end behavior of the polynomial below.

$$f(x) = -3(x+5)^5(x-5)^8(x+9)^2(x-9)^3$$

The solution is the graph below, which is option B.



C.



E. None of the above.

General Comment: Remember that end behavior is determined by the leading coefficient AND whether the **sum** of the multiplicities is positive or negative.

7. Construct the lowest-degree polynomial given the zeros below. Then, choose the intervals that contain the coefficients of the polynomial in the form $x^3 + bx^2 + cx + d$.

$$4 - 3i \text{ and } -1$$

The solution is $x^3 - 7x^2 + 17x + 25$, which is option B.

A. $b \in [6, 11], c \in [16, 18]$, and $d \in [-26, -24]$ $x^3 + 7x^2 + 17x - 25$, which corresponds to multiplying out (x - (4 - 3i))(x - (4 + 3i))(x - 1).

B. $b \in [-9, -4], c \in [16, 18]$, and $d \in [20, 28]$ $* x^3 - 7x^2 + 17x + 25$, which is the correct option.

C. $b \in [-2, 6], c \in [-4, -2], \text{ and } d \in [-4, -1]$ $x^3 + x^2 - 3x - 4, \text{ which corresponds to multiplying out } (x - 4)(x + 1).$

D. $b \in [-2, 6], c \in [3, 5], \text{ and } d \in [-1, 8]$ $x^3 + x^2 + 4x + 3, \text{ which corresponds to multiplying out } (x + 3)(x + 1).$

E. None of the above.

This corresponds to making an unanticipated error or not understanding how to use nonreal complex numbers to create the lowest-degree polynomial. If you chose this and are not sure what you did wrong, please contact the coordinator for help.

General Comment: Remember that the conjugate of a + bi is a - bi. Since these zeros always come in pairs, we need to multiply out (x - (4 - 3i))(x - (4 + 3i))(x - (-1)).

8. Construct the lowest-degree polynomial given the zeros below. Then, choose the intervals that contain the coefficients of the polynomial in the form $ax^3 + bx^2 + cx + d$.

$$\frac{-2}{3}, \frac{-1}{4}, \text{ and } \frac{7}{4}$$

The solution is $48x^3 - 40x^2 - 69x - 14$, which is option B.


- A. $a \in [44, 54], b \in [-110, -98], c \in [21, 29], \text{ and } d \in [9, 22]$ $48x^3 - 104x^2 + 27x + 14$, which corresponds to multiplying out (3x - 2)(4x + 1)(4x - 7).
- B. $a \in [44, 54], b \in [-41, -35], c \in [-74, -64], \text{ and } d \in [-15, -11]$ * $48x^3 - 40x^2 - 69x - 14$, which is the correct option.
- C. $a \in [44, 54], b \in [-41, -35], c \in [-74, -64]$, and $d \in [9, 22]$ $48x^3 - 40x^2 - 69x + 14$, which corresponds to multiplying everything correctly except the constant term.
- D. $a \in [44, 54], b \in [-134, -124], c \in [85, 86], \text{ and } d \in [-15, -11]$ $48x^3 - 128x^2 + 85x - 14, \text{ which corresponds to multiplying out } (3x - 2)(4x - 1)(4x - 7).$
- E. $a \in [44, 54], b \in [40, 42], c \in [-74, -64], \text{ and } d \in [9, 22]$ $48x^3 + 40x^2 - 69x + 14, \text{ which corresponds to multiplying out } (3x - 2)(4x - 1)(4x + 7).$

General Comment: To construct the lowest-degree polynomial, you want to multiply out (3x + 2)(4x + 1)(4x - 7)


9. Describe the end behavior of the polynomial below.

$$f(x) = 3(x-3)^{2}(x+3)^{7}(x+9)^{5}(x-9)^{7}$$

The solution is the graph below, which is option D.




--



E. None of the above.

General Comment: Remember that end behavior is determined by the leading coefficient AND whether the **sum** of the multiplicities is positive or negative.

10. Which of the following equations *could* be of the graph presented below?



The solution is $4(x-1)^{11}(x-3)^{11}(x+4)^9$, which is option E.

A.
$$17(x-1)^{10}(x-3)^6(x+4)^{11}$$

The factors 1 and 3 have have been odd power.

B.
$$-20(x-1)^{10}(x-3)^{11}(x+4)^{11}$$

The factor (x-1) should have an odd power and the leading coefficient should be the opposite sign.

C.
$$4(x-1)^8(x-3)^{11}(x+4)^5$$

The factor 1 should have been an odd power.

D.
$$-20(x-1)^{11}(x-3)^5(x+4)^5$$

This corresponds to the leading coefficient being the opposite value than it should be.

E.
$$4(x-1)^{11}(x-3)^{11}(x+4)^9$$

General Comment: General Comments: Draw the x-axis to determine which zeros are touching (and so have even multiplicity) or cross (and have odd multiplicity).

2107-1615 test

^{*} This is the correct option.