

This key should allow you to understand why you choose the option you did (beyond just getting a question right or wrong). More instructions on how to use this key can be found [here](#).

If you have a suggestion to make the keys better, please fill out the short survey [here](#).

Note: This key is auto-generated and may contain issues and/or errors. The keys are reviewed after each exam to ensure grading is done accurately. If there are issues (like duplicate options), they are noted in the offline gradebook. The keys are a work-in-progress to give students as many resources to improve as possible.

1. A town has an initial population of 30000. The town's population for the next 10 years is provided below. Which type of function would be most appropriate to model the town's population?

Year	1	2	3	4	5	6	7	8	9
Pop.	29980	29960	29940	29920	29900	29880	29860	29840	29820

The solution is Linear, which is option A.

A. Linear

This suggests a constant growth. You would be able to add or subtract the same amount year-to-year if this is the correct answer.

B. Exponential

This suggests the fastest of growths that we know.

C. Non-Linear Power

This suggests a growth faster than constant but slower than exponential.

D. Logarithmic

This suggests the slowest of growths that we know.

E. None of the above

Please contact the coordinator to discuss why you believe none of the options model the population.

General Comment: We are trying to compare the growth rate of the population. Growth rates can be characterized from slowest to fastest as: logarithmic, indirect, linear, direct, exponential. The best way to approach this is to first compare it to linear (is it linear, faster than linear, or slower than linear)? If faster, is it as fast as exponential? If slower, is it as slow as logarithmic?

2. For the information below, construct a linear model that describes the total time T spent on the path in terms of the distance of a particular part of the path *if we know that all parts of the path are equal length*.

A bicyclist is training for a race on a hilly path. Their bike keeps track of their speed at any time, but not the distance traveled. Their speed traveling up a hill is 5 mph, 8 mph when traveling down a hill, and 6 mph when traveling along a flat portion.

The solution is $0.492D$, which is option B.

A. $240.000D$

The coefficient here is calculated by multiplying the distances together rather than adding.

B. $0.492D$

* This is the correct option.

C. $19.000D$

The coefficient here is calculated as if you were trying to model the distance on the total path.

D. The model can be found with the information provided, but isn't options 1-3.

Since we know all parts of the path are equal length, we can treat all distance variables as the same variable, D .

E. The model cannot be found with the information provided.

If you chose this option, please contact the coordinator to discuss why you think we cannot model the situation.

General Comment: Be sure you pay attention to the variable we are writing the model in terms of. To create the model with a single variable, we have to know that variable is the same throughout each path!

3. For the information provided below, construct a linear model that describes her total costs, C , as a function of the number of months, x she is at UF.

Aubrey is a college student going into her first year at UF. She will receive Bright Futures, which covers her tuition plus a \$800 educational expense each year. Before college, Aubrey saved up \$5000. She knows she will need to pay \$900 in rent a month, \$60 for food a week, and \$56 in other weekly expenses.

The solution is None of the above., which is option E.

A. $C(x) = 1016x$

This treats weekly expenses as monthly expenses rather than multiplying each weekly expense by 4.

B. $C(x) = 5800$

This describes the student's income, not costs.

C. $C(x) = 5800x$

This describes the student's income as if they received the savings and educational expense each month.

D. $C(x) = 1016$

This treats weekly expenses as month expenses rather than multiplying each weekly expense by 4 AND does not account for these expenses per month.

E. None of the above.

* This is the correct option as the model should be $C(x) = 1364x$.

General Comment: This is a Costs, Profit, Revenue question! The most common issues here are: (1) not converting the weekly costs to monthly costs, (2) treating the one-time values like savings and educational expense as happening per month, and (3) not checking that your model is for cost, profit [income], or revenue [budget].

4. A town has an initial population of 20000. The town's population for the next 10 years is provided below. Which type of function would be most appropriate to model the town's population?

Year	1	2	3	4	5	6	7	8	9
Pop.	19970	19940	19910	19880	19850	19820	19790	19760	19730

The solution is Linear, which is option A.

A. Linear

This suggests a constant growth. You would be able to add or subtract the same amount year-to-year if this is the correct answer.

B. Logarithmic

This suggests the slowest of growths that we know.

C. Non-Linear Power

This suggests a growth faster than constant but slower than exponential.

D. Exponential

This suggests the fastest of growths that we know.

E. None of the above

Please contact the coordinator to discuss why you believe none of the options model the population.

General Comment: We are trying to compare the growth rate of the population. Growth rates can be characterized from slowest to fastest as: logarithmic, indirect, linear, direct, exponential. The best way to approach this is to first compare it to linear (is it linear, faster than linear, or slower than linear)? If faster, is it as fast as exponential? If slower, is it as slow as logarithmic?

5. What is the **best** way to describe the domain of the scenario below?

Bridges on highways often have expansion joints, which are small gaps in the roadway between one bridge section and the next. The gaps are put there so the bridge will have room to expand when the weather gets hot. Assume the gap width varies constantly with the temperature. Suppose a bridge has a gap of 1.3 cm when the temperature is 22 degrees C and that the gap narrows to 0.9 cm when the temperature warms to 30 degrees C.

The solution is There is no restricted domain in this scenario, which is option D.

A. Subset of the Integers

Recall that the Integers are the positive and negative counting numbers: ..., -3, -2, -1, 0, 1, 2, 3, ...

B. Subset of the Natural numbers

Recall that the Naturals are the counting numbers: 1, 2, 3, ...

C. Proper subset of the Real numbers

This means we have a domain of the Real numbers but need to throw out values based on the context.

D. There is no restricted domain in this scenario

This means we have a domain of the Real numbers and we don't need to remove any values even in the real-world context.

E. Subset of the Rational numbers

Recall that the Rationals are fractions with Integers in the numerator and denominator.

General Comment: We often have to remove values in the domain when working with real-world models.

6. For the information provided below, construct a linear model that describes her total income, I , as a function of the number of months, x she is at UF.

Aubrey is a college student going into her first year at UF. She will receive Bright Futures, which covers her tuition plus a \$1000 educational expense each year. Before college, Aubrey saved up \$10000. She knows she will need to pay \$1100 in rent a month, \$80 for food a week, and \$64 in other weekly expenses.

The solution is $I(x) = 11000$, which is option C.

A. $I(x) = 11000x$

This treats the educational expense and savings as something you get every month rather than a 1-time payment.

B. $I(x) = 1000x + 10000$

This treats the educational expense as something you get every month rather than a 1-time payment.

C. $I(x) = 11000$

* This is the correct option.

D. $I(x) = 10000x + 1000$

This treats the savings as something you get every month rather than a 1-time payment.

E. None of the above.

You may have chosen this as you thought you were modeling total costs or total budget.

General Comment: This is a Costs, Profit, Revenue question! The most common issues here are: (1) not converting the weekly costs to monthly costs, (2) treating the one-time values like savings and educational expense as happening per month, and (3) not checking that your model is for cost, profit [income], or revenue [budget].

-
7. What is the **best** way to describe the domain of the scenario below?

Chemists commonly create a solution by mixing two products of differing concentrations together. A 10% and 30% solution can make an acid solution of some value between these, such as a 24% acid solution. The chemist wants to make differing solution percentages of 7 liters each.

The solution is Proper subset of the Real numbers, which is option C.

A. Subset of the Natural numbers

Recall that the Naturals are the counting numbers: 1, 2, 3, ...

B. Subset of the Rational numbers

Recall that the Rationals are fractions with Integers in the numerator and denominator.

C. Proper subset of the Real numbers

This means we have a domain of the Real numbers but need to throw out values based on the context.

D. Subset of the Integers

Recall that the Integers are the positive and negative counting numbers: ..., -3, -2, -1, 0, 1, 2, 3, ...

- E. There is no restricted domain in this scenario

This means we have a domain of the Real numbers and we don't need to remove any values even in the real-world context.

General Comment: We often have to remove values in the domain when working with real-world models.

8. Using the situation below, construct a linear model that describes the cost of the coffee beans $C(h)$ in terms of the weight of the high-quality coffee beans h .

Veronica needs to prepare 70 of blended coffee beans selling for \$3.84 per pound. She has a high-quality bean that sells for \$4.79 a pound and a low-quality bean that sells for \$3.19 a pound.

The solution is $C(h) = 1.60h + 223.30$, which is option D.

- A. $C(h) = -1.60h + 335.30$

This would be correct if the question asked you to construct the cost model in terms of the weight of the low-quality bean.

- B. $C(h) = 3.99h$

This assumes that exactly half of the high- and low- quality beans are mixed to create the blended coffee beans.

- C. $C(h) = 4.79h$

This models the cost of the high-quality bean only, not the blended beans.

- D. $C(h) = 1.60h + 223.30$

* This is the correct option since the questions asked you to construct the cost model in terms of the weight of the high-quality bean.

- E. None of the above.

If you chose this option, please talk to the coordinator to discuss why.

General Comment: This is exactly like the chemistry mixture question from the homework! If you are having trouble with this problem, be sure to review the video for building linear models.

9. For the information below, construct a linear model that describes the total time T spent on the path in terms of the distance of a particular part of the path *if we know that all parts of the path are equal length*.

A bicyclist is training for a race on a hilly path. Their bike keeps track of their speed at any time, but not the distance traveled. Their speed traveling up a hill is 5 mph, 11 mph when traveling down a hill, and 8 mph when traveling along a flat portion.

The solution is $0.416D$, which is option A.

- A. $0.416D$

* This is the correct option.

- B. $440.000D$

The coefficient here is calculated by multiplying the distances together rather than adding.

- C. $24.000D$

The coefficient here is calculated as if you were trying to model the distance on the total path.

- D. The model can be found with the information provided, but isn't options 1-3.

Since we know all parts of the path are equal length, we can treat all distance variables as the same variable, D .

- E. The model cannot be found with the information provided.

If you chose this option, please contact the coordinator to discuss why you think we cannot model the situation.

General Comment: Be sure you pay attention to the variable we are writing the model in terms of. To create the model with a single variable, we have to know that variable is the same throughout each path!

10. Using the situation below, construct a linear model that describes the cost of the coffee beans $C(h)$ in terms of the weight of the high-quality coffee beans h .

Veronica needs to prepare 240 of blended coffee beans selling for \$3.23 per pound. She has a high-quality bean that sells for \$4.19 a pound and a low-quality bean that sells for \$2.56 a pound.

The solution is $C(h) = 1.63h + 614.40$, which is option C.

- A. $C(h) = 4.19h$

This models the cost of the high-quality bean only, not the blended beans.

- B. $C(h) = 3.38h$

This assumes that exactly half of the high- and low- quality beans are mixed to create the blended coffee beans.

- C. $C(h) = 1.63h + 614.40$

* This is the correct option since the questions asked you to construct the cost model in terms of the weight of the high-quality bean.

- D. $C(h) = -1.63h + 1005.60$

This would be correct if the question asked you to construct the cost model in terms of the weight of the low-quality bean.

- E. None of the above.

If you chose this option, please talk to the coordinator to discuss why.

General Comment: This is exactly like the chemistry mixture question from the homework! If you are having trouble with this problem, be sure to review the video for building linear models.
