This key should allow you to understand why you choose the option you did (beyond just getting a question right or wrong). More instructions on how to use this key can be found here.

If you have a suggestion to make the keys better, please fill out the short survey here.

Note: This key is auto-generated and may contain issues and/or errors. The keys are reviewed after each exam to ensure grading is done accurately. If there are issues (like duplicate options), they are noted in the offline gradebook. The keys are a work-in-progress to give students as many resources to improve as possible.

36. Solve the equation for x and choose the interval that contains the solution (if it exists).

$$2^{3x-2} = 27^{2x+4}$$

The solution is x = -3.229

A. $x \in [-2.3, -1.1]$

x = -1.330, which corresponds to distributing the $\ln(base)$ to the first term of the exponent only.

B. $x \in [5.6, 6.6]$

x = 6.000, which corresponds to solving the numerators as equal while ignoring the bases are different.

C. $x \in [12.4, 16]$

x = 14.570, which corresponds to distributing the $\ln(base)$ to the second term of the exponent only.

D. $x \in [-4.8, -2.1]$

* x = -3.229, which is the correct option.

E. There is no Real solution to the equation.

This corresponds to believing there is no solution since the bases are not powers of each other.

General Comments: This question was written so that the bases could not be written the same. You will need to take the log of both sides.

37. Solve the equation for x and choose the interval that contains x (if it exists).

$$5 = \sqrt[7]{\frac{20}{e^{8x}}}$$

The solution is x = -1.034

A. $x \in [-0.4, 0.3]$

x = -0.028, which corresponds to treating any root as a square root.

B. $x \in [-2.2, -0.8]$

* x = -1.034, which is the correct option.

C. $x \in [-5.6, -4.6]$

x = -4.749, which corresponds to thinking you don't need to take the natural log of both sides before reducing, as if the equation already had a natural log on the right side.

D. There is no Real solution to the equation.

This corresponds to believing you cannot solve the equation.

E. None of the above.

This corresponds to making an unexpected error.

General Comments: After using the properties of logarithmic functions to break up the right-hand side, use ln(e) = 1 to reduce the question to a linear function to solve. You can put ln(20) into a calculator if you are having trouble.

38. Which of the following intervals describes the Domain of the function below?

$$f(x) = \log_2\left(x - 5\right) - 3$$

The solution is $(5, \infty)$

A. $(-\infty, a), a \in [-7.3, -4]$

 $(-\infty, -5)$, which corresponds to flipping the Domain. Remember: the general for is $a * \log(x - h) + k$, where a does not affect the domain.

B. $[a, \infty), a \in [-3.1, -0.7]$

 $[-3,\infty)$, which corresponds to using the vertical shift when shifting the Domain AND including the endpoint.

C. $(-\infty, a], a \in [-1.6, 3.6]$

 $(-\infty,3]$, which corresponds to using the negative vertical shift AND including the endpoint AND flipping the domain.

D. $(a, \infty), a \in [4.9, 7.1]$

* $(5, \infty)$, which is the correct option.

E. $(-\infty, \infty)$

This corresponds to thinking of the range of the log function (or the domain of the exponential function).

General Comments: The domain of a basic logarithmic function is $(0, \infty)$ and the Range is $(-\infty, \infty)$. We can use shifts when finding the Domain, but the Range will always be all Real numbers.

39. Solve the equation for x and choose the interval that contains the solution (if it exists).

$$\log_2(3x+6) + 6 = 3$$

The solution is x = -1.958

A. $x \in [-2.38, -1.44]$

* x = -1.958, which is the correct option.

B. $x \in [0.43, 0.84]$

x = 0.667, which corresponds to ignoring the vertical shift when converting to exponential form.

C. $x \in [0.97, 1.39]$

x = 1.000, which corresponds to reversing the base and exponent when converting.

D. $x \in [4.93, 5.77]$

x = 5.000, which corresponds to reversing the base and exponent when converting and reversing the value with x.

E. There is no Real solution to the equation.

Corresponds to believing a negative coefficient within the log equation means there is no Real solution.

General Comments: First, get the equation in the form $\log_b{(cx+d)} = a$. Then, convert to $b^a = cx + d$ and solve.

40. Which of the following intervals describes the Range of the function below?

$$f(x) = e^{x+1} + 2$$

The solution is $(2, \infty)$

A. $[a, \infty), a \in [1.8, 2.8]$

 $[2,\infty)$, which corresponds to including the endpoint.

B. $(a, \infty), a \in [1.8, 2.8]$

* $(2, \infty)$, which is the correct option.

C. $(-\infty, a), a \in [-4.1, 1.4]$

 $(-\infty, -2)$, which corresponds to using the negative vertical shift AND flipping the Range interval.

D. $(-\infty, a], a \in [-4.1, 1.4]$

 $(-\infty, -2]$, which corresponds to using the negative vertical shift AND flipping the Range interval AND including the endpoint.

E. $(-\infty, \infty)$

This corresponds to confusing range of an exponential function with the domain of an exponential function.

General Comments: Domain of a basic exponential function is $(-\infty, \infty)$ while the Range is $(0, \infty)$. We can shift these intervals [and even flip when a < 0!] to find the new Domain/Range.