

1. A town has an initial population of 20000. The town's population for the next 10 years is provided below. Which type of function would be most appropriate to model the town's population?

Year	1	2	3	4	5	6	7	8	9
Pop.	20080	20160	20320	20640	21280	22560	25120	30240	40480

- A. Linear
- B. Non-Linear Power
- C. Logarithmic
- D. Exponential
- E. None of the above

2. For the information below, construct a linear model that describes the total time  $T$  spent on the path in terms of the distance of a particular part of the path *if we know that all parts of the path are equal length*.

*A bicyclist is training for a race on a hilly path. Their bike keeps track of their speed at any time, but not the distance traveled. Their speed traveling up a hill is 5 mph, 11 mph when traveling down a hill, and 7 mph when traveling along a flat portion.*

- A.  $23.000D$
- B.  $385.000D$
- C.  $0.434D$
- D. The model can be found with the information provided, but isn't options 1-3
- E. The model cannot be found with the information provided.

3. Using the situation below, construct a linear model that describes the cost of the coffee beans  $C(h)$  in terms of the weight of the low-quality coffee beans  $h$ .

*Veronica needs to prepare 130 of blended coffee beans selling for \$3.91 per pound. She has a high-quality bean that sells for \$5.46 a pound and a low-quality bean that sells for \$2.63 a pound.*

- A.  $C(h) = 4.04h$
  - B.  $C(h) = 2.83h + 341.90$
  - C.  $C(h) = 2.63h$
  - D.  $C(h) = -2.83h + 709.80$
  - E. None of the above.
- 

4. What is the **best** way to describe the domain of the scenario below?

*Fred is a store manager at Publix. The store normally orders two pallets of water bottles a week and sells 1000 bottles per day. However, a hurricane is coming and Fred expects water bottle sales to increase tenfold for three days, then decrease by half of normal sales for four days. How many more pallets of water bottles should Fred order the week before the hurricane?*

- A. Subset of the Integers
  - B. There is no restricted domain in this scenario
  - C. Subset of the Rational numbers
  - D. Proper subset of the Real numbers
  - E. Subset of the Natural numbers
- 

5. What is the **best** way to describe the domain of the scenario below?

*Fred is a store manager at Publix. The store normally orders two pallets of water bottles a week and sells 1000 bottles per day. However, a hurricane is coming and Fred expects water bottle sales to increase tenfold for three days, then decrease by half of normal sales for four days. How many more pallets of water bottles should Fred order the week before the hurricane?*

- A. Subset of the Natural numbers
- B. There is no restricted domain in this scenario
- C. Proper subset of the Real numbers

- D. Subset of the Integers
  - E. Subset of the Rational numbers
- 

6. For the information provided below, construct a linear model that describes the total distance of the path,  $D$ , in terms of the time spent on a particular path *if we know that the time spent on each path was equal*.

*A bicyclist is training for a race on a hilly path. Their bike keeps track of their speed at any time, but not the distance traveled. Their speed traveling up a hill is 4 mph, 10 mph when traveling down a hill, and 7 mph when traveling along a flat portion.*

- A.  $21t$
  - B.  $0.493t$
  - C.  $280t$
  - D. The model can be found with the information provided, but isn't options 1-3
  - E. The model cannot be found with the information provided.
- 

7. Using the situation below, construct a linear model that describes the cost of the coffee beans  $C(h)$  in terms of the weight of the high-quality coffee beans  $h$ .

*Veronica needs to prepare 90 of blended coffee beans selling for \$3.63 per pound. She has a high-quality bean that sells for \$4.14 a pound and a low-quality bean that sells for \$2.50 a pound.*

- A.  $C(h) = -1.64h + 372.60$
  - B.  $C(h) = 1.64h + 225.00$
  - C.  $C(h) = 3.32h$
  - D.  $C(h) = 4.14h$
  - E. None of the above.
-

8. For the information provided below, construct a linear model that describes her total budget,  $B$ , as a function of the number of months,  $x$  she is at UF.

*Aubrey is a college student going into her first year at UF. She will receive Bright Futures, which covers her tuition plus a \$800 educational expense each year. Before college, Aubrey saved up \$9000. She knows she will need to pay \$800 in rent a month, \$50 for food a week, and \$48 in other weekly expenses.*

- A.  $B(x) = 9000x + 800$
  - B.  $B(x) = 9800 - 898x$
  - C.  $B(x) = 9800 - 1192x$
  - D.  $B(x) = 800x + 9000$
  - E. None of the above.
- 

9. For the information provided below, construct a linear model that describes her total costs,  $C$ , as a function of the number of months,  $x$  she is at UF.

*Aubrey is a college student going into her first year at UF. She will receive Bright Futures, which covers her tuition plus a \$600 educational expense each year. Before college, Aubrey saved up \$10000. She knows she will need to pay \$700 in rent a month, \$40 for food a week, and \$32 in other weekly expenses.*

- A.  $C(x) = 10600$
  - B.  $C(x) = 772x$
  - C.  $C(x) = 772$
  - D.  $C(x) = 10600x$
  - E. None of the above.
- 

10. A town has an initial population of 100000. The town's population for the next 10 years is provided below. Which type of function would be

most appropriate to model the town's population?

Year	1	2	3	4	5	6	7	8	9
Pop.	100020	100040	100060	100080	100100	100120	100140	100160	100180

- A. Non-Linear Power
  - B. Logarithmic
  - C. Exponential
  - D. Linear
  - E. None of the above
-