

1. For the scenario below, use the model for the volume of a cylinder as $V = \pi r^2 h$ to find the coefficient for the model of the new volume $V_{extnew} = kr^2 h$.

Pepsi wants to increase the volume of soda in their cans. They've decided to increase the radius by 17 percent and increase the height by 10 percent. They want to model the new volume based on the radius and height of the original cans.

2. A town has an initial population of 100000. The town's population for the next 9 years is provided below. Which type of function would be most appropriate to model the town's population?

Year	1	2	3	4	5	6	7	8	9
Pop	100000	99986	99978	99972	99967	99964	99961	99958	99956

3. What model type would best describe the scenario below?

Social distancing is a common tactic to counter potential epidemics. This is due to the exponential increase in number of people infected as the density of people living in an area increases.

4. For the scenario below, model the rate of vibration (cm/s) of the string in terms of the length of the string. Then determine the variation constant k of the model (if possible). The constant should be in terms of cm and s.

The rate of vibration of a string under constant tension varies based on the type of string and the length of the string. The rate of vibration of string ω decreases as the square length of the string increases. For example, when string ω is 5 mm long, the rate of vibration is 24 cm/s.

5. For the scenario below, use the model for the volume of a cylinder as $V = \pi r^2 h$ to find the coefficient for the model of the new volume $V_{extnew} = kr^2 h$.

Pepsi wants to increase the volume of soda in their cans. They've decided to decrease the radius by 13 percent and increase the height by 16 percent. They want to model the new volume based on the radius and height of the original cans.

6. A town has an initial population of 100000. The town’s population for the next 9 years is provided below. Which type of function would be most appropriate to model the town’s population?

Year	1	2	3	4	5	6	7	8	9
Pop	100018	100046	100058	100086	100098	100126	100138	100166	100

7. For the scenario below, model the rate of vibration (cm/s) of the string in terms of the length of the string. Then determine the variation constant k of the model (if possible). The constant should be in terms of cm and s.

The rate of vibration of a string under constant tension varies based on the type of string and the length of the string. The rate of vibration of string ω decreases as the cube length of the string increases. For example, when string ω is 3 mm long, the rate of vibration is 20 cm/s.

8. What model type would best describe the scenario below?

Social distancing is a common tactic to counter potential epidemics. This is due to the exponential increase in number of people infected as the density of people living in an area increases.

9. For the scenario below, find the variation constant k of the model (if possible).

In an alternative galaxy, the cube of the time, T (Earth years), required for a planet to orbit Sun χ increases as the cube of the distance, d (AUs), that the planet is from Sun χ increases. For example, when Ea's average distance from Sun χ is 2, it takes 82 Earth days to complete an orbit.

10. For the scenario below, find the variation constant k of the model (if possible).

In an alternative galaxy, the cube of the time, T (Earth years), required for a planet to orbit Sun χ decreases as the cube of the distance, d (AUs), that the planet is from Sun χ decreases. For example, when Ea's average distance from Sun χ is 4, it takes 67 Earth days to complete an orbit.

11. For the scenario below, use the model for the volume of a cylinder as $V = \pi r^2 h$ to find the coefficient for the model of the new volume $V_{extnew} = kr^2 h$.

Pepsi wants to increase the volume of soda in their cans. They've decided to increase the radius by 19 percent and increase the height by 12 percent. They want to model the new volume based on the radius and height of the original cans.

12. A town has an initial population of 80000. The town's population for the next 9 years is provided below. Which type of function would be most appropriate to model the town's population?

Year	1	2	3	4	5	6	7	8	9
Pop	80017	80045	80057	80085	80097	80125	80137	80165	80177

13. What model type would best describe the scenario below?

Big O notation is common in computer science to describe how fast a program can solve a particular problem. Big O notation categorizes functions according to their growth rates, the same way we have categorized modeling real-world problems by certain types of functions. When analyzing a particular program, a student found the computer to need x^x time to complete, where x was the number of inputs into the program.

14. For the scenario below, model the rate of vibration (cm/s) of the string in terms of the length of the string. Then determine the variation constant k of the model (if possible). The constant should be in terms of cm and s.

The rate of vibration of a string under constant tension varies based on the type of string and the length of the string. The rate of vibration of string ω decreases as the quartic length of the string increases. For example, when string ω is 5 mm long, the rate of vibration is 21 cm/s.

15. For the scenario below, use the model for the volume of a cylinder as $V = \pi r^2 h$ to find the coefficient for the model of the new volume $V_{extnew} = kr^2 h$.

Pepsi wants to increase the volume of soda in their cans. They've decided to decrease the radius by 11 percent and increase the height by 18 percent. They want to model the new volume based on the radius and height of the original cans.

16. A town has an initial population of 80000. The town's population for the next 9 years is provided below. Which type of function would be most appropriate to model the town's population?

Year	1	2	3	4	5	6	7	8	9
Pop	80000	80013	80021	80027	80032	80035	80038	80041	80043

17. For the scenario below, model the rate of vibration (cm/s) of the string in terms of the length of the string. Then determine the variation constant k of the model (if possible). The constant should be in terms of cm and s.

The rate of vibration of a string under constant tension varies based on the type of string and the length of the string. The rate of vibration of string ω decreases as the quartic length of the string increases. For example, when string ω is 4 mm long, the rate of vibration is 30 cm/s.

18. What model type would best describe the scenario below?

In economics, there are two common equations to model interest earned. The compound interest formula is $A = P(1 + \frac{r}{n})^{nt}$, where A is the amount of money you end up with, P is your starting money, r is the interest rate, n is the number of times compounded in a year, and t is the total number of years. For example, if you were a parent and wanted to save \$10,000 in 3 years-time at 3.5% interest compounded monthly, you would need to invest about \$9,000.

19. For the scenario below, find the variation constant k of the model (if possible).

In an alternative galaxy, the cube of the time, T (Earth years), required for a planet to orbit Sun χ increases as the square of the distance, d (AUs), that the planet is from Sun χ increases. For example, when Ea's average distance from Sun χ is 7, it takes 56 Earth days to complete an orbit.

20. For the scenario below, find the variation constant k of the model (if possible).

In an alternative galaxy, the quartic of the time, T (Earth years), required for a planet to orbit Sun χ increases as the quartic of the distance, d (AUs), that the planet is from Sun χ increases. For example, when Ea's average distance from Sun χ is 3, it takes 95 Earth days to complete an orbit.

21. For the scenario below, use the model for the volume of a cylinder as $V = \pi r^2 h$ to find the coefficient for the model of the new volume $V_{extnew} = kr^2 h$.

Pepsi wants to increase the volume of soda in their cans. They've decided to decrease the radius by 20 percent and decrease the height by 12 percent. They want to model the new volume based on the radius and height of the original cans.

22. A town has an initial population of 90000. The town's population for the next 9 years is provided below. Which type of function would be most appropriate to model the town's population?

Year	1	2	3	4	5	6	7	8	9
Pop	90000	90034	90054	90069	90080	90089	90097	90103	90109

23. What model type would best describe the scenario below?

In economics, there are two common equations to model interest earned. The compound interest formula is $A = P(1 + \frac{r}{n})^{nt}$, where A is the amount of money you end up with, P is your starting money, r is the interest rate, n is the number of times compounded in a year, and t is the total number of years. For example, if you were a parent and wanted to save \$10,000 in 3 years-time at 3.5% interest compounded monthly, you would need to invest about \$9,000.

24. For the scenario below, model the rate of vibration (cm/s) of the string in terms of the length of the string. Then determine the variation constant k of the model (if possible). The constant should be in terms of cm and s.

The rate of vibration of a string under constant tension varies based on the type of string and the length of the string. The rate of vibration of string ω increases as the cube length of the string decreases. For example, when string ω is 3 mm long, the rate of vibration is 31 cm/s.

25. For the scenario below, use the model for the volume of a cylinder as $V = \pi r^2 h$ to find the coefficient for the model of the new volume $V_{extnew} = kr^2 h$.

Pepsi wants to increase the volume of soda in their cans. They've decided to increase the radius by 16 percent and increase the height by 11 percent. They want to model the new volume based on the radius and height of the original cans.

26. A town has an initial population of 20000. The town's population for the next 9 years is provided below. Which type of function would be most appropriate to model the town's population?

Year	1	2	3	4	5	6	7	8	9
Pop	20000	19972	19956	19944	19935	19928	19922	19916	19912

27. For the scenario below, model the rate of vibration (cm/s) of the string in terms of the length of the string. Then determine the variation constant k of the model (if possible). The constant should be in terms of cm and s.

The rate of vibration of a string under constant tension varies based on the type of string and the length of the string. The rate of vibration of string ω increases as the square length of the string decreases. For example, when string ω is 5 mm long, the rate of vibration is 22 cm/s.

28. What model type would best describe the scenario below?

Big O notation is common in computer science to describe how fast a program can solve a particular problem. Big O notation categorizes functions according to their growth rates, the same way we have categorized modeling real-world problems by certain types of functions. When analyzing a particular program, a student found the computer to need x^x time to complete, where x was the number of inputs into the program.

29. For the scenario below, find the variation constant k of the model (if possible).

In an alternative galaxy, the quartic of the time, T (Earth years), required for a planet to orbit Sun χ increases as the square of the distance, d (AUs), that the planet is from Sun χ increases. For example, when Ea's average distance from Sun χ is 2, it takes 91 Earth days to complete an orbit.

30. For the scenario below, find the variation constant k of the model (if possible).

In an alternative galaxy, the square of the time, T (Earth years), required for a planet to orbit Sun χ increases as the quartic of the distance, d (AUs), that the planet is from Sun χ increases. For example, when Ea's average distance from Sun χ is 2, it takes 97 Earth days to complete an orbit.