1. Using the scenario below, model the situation using an exponential function and a base of  $\frac{1}{2}$ . Then, solve for the half-life of the element, rounding to the nearest day.

The half-life of an element is the amount of time it takes for the element to decay to half of its initial starting amount. There is initially 967 grams of element X and after 5 years there is 107 grams remaining.

- A. About 365 days
- B. About 2190 days
- C. About 0 days
- D. About 730 days
- E. None of the above
- 2. Using the scenario below, model the population of bacteria  $\alpha$  in terms of the number of minutes, t that pass. Then, choose the correct approximate (rounded to the nearest minute) replication rate of bacteria- $\alpha$ .

A newly discovered bacteria,  $\alpha$ , is being examined in a lab. The lab started with a petri dish of 3 bacteria- $\alpha$ . After 1 hours, the petri dish has 10 bacteria- $\alpha$ . Based on similar bacteria, the lab believes bacteria- $\alpha$  doubles after some undetermined number of minutes.

- A. About 329 minutes
- B. About 54 minutes
- C. About 50 minutes
- D. About 304 minutes
- E. None of the above
- 3. The temperature of an object, T, in a different surrounding temperature  $T_s$  will behave according to the formula  $T(t) = Ae^{kt} + T_s$ , where t is minutes, A is a constant, and k is a constant. Use this formula and

Summer C 2020 Version B

the situation below to construct a model that describes the uranium's temperature, T, based on the amount of time t (in minutes) that have passed. Choose the correct constant k from the options below.

Uranium is taken out of the reactor with a temperature of 120° C and is placed into a 17° C bath to cool. After 34 minutes, the uranium has cooled to 77° C.


A. 
$$k = -0.02039$$

B. 
$$k = -0.01589$$

C. 
$$k = -0.02137$$

D. 
$$k = -0.02081$$

- E. None of the above
- 4. Determine the appropriate model for the graph of points below.



- A. Non-linear Power model
- B. Exponential model
- C. Logarithmic model
- D. Linear model
- E. None of the above

Summer C 2020 Version B

5. A town has an initial population of 20000. The town's population for the next 10 years is provided below. Which type of function would be most appropriate to model the town's population?

Year	1	2	3	4	5	6	7	8	9
Pop.	19970	19520	17570	12320	1250	0	0	0	0

- A. Exponential
- B. Logarithmic
- C. Non-Linear Power
- D. Linear
- E. None of the above

Summer C 2020 Version B