

This key should allow you to understand why you choose the option you did (beyond just getting a question right or wrong). More instructions on how to use this key can be found [here](#).

If you have a suggestion to make the keys better, please fill out the short survey [here](#).

Note: This key is auto-generated and may contain issues and/or errors. The keys are reviewed after each exam to ensure grading is done accurately. If there are issues (like duplicate options), they are noted in the offline gradebook. The keys are a work-in-progress to give students as many resources to improve as possible.

1. Factor the quadratic below. Then, choose the intervals that contain the constants in the form $(ax + b)(cx + d); b \leq d$.

$$24x^2 + 38x + 15$$

The solution is $(4x + 3)(6x + 5)$

A. $a \in [3.99, 4.27]$, $b \in [2, 8]$, $c \in [5.98, 6.1]$, and $d \in [2, 7]$

* $(4x + 3)(6x + 5)$, which is the correct option.

B. $a \in [1.23, 1.49]$, $b \in [2, 8]$, $c \in [17.17, 18.57]$, and $d \in [2, 7]$

$(1.33333333333333x + 3)(18x + 5)$, which corresponds to associating some factor of c to a .

C. $a \in [7.87, 8.11]$, $b \in [2, 8]$, $c \in [2.99, 3.71]$, and $d \in [2, 7]$

$(8x + 3)(3.0x + 5)$, which corresponds to associating some factor of a to c .

D. $a \in [0.58, 1.25]$, $b \in [13, 20]$, $c \in [0.61, 1.5]$, and $d \in [18, 21]$


$(x + 18)(x + 20)$, which corresponds to factoring $x^2 + 38x + 360$.

E. None of the above.

Corresponds to a different factoring than any of the predicted options. If you get this, please let the coordinator know so they can work with you to figure out what went wrong with your factoring.

General Comment: General Comments: ac had many factors in this problem. It is best to list out the possible pairs in order to make sure you don't miss any.

2. Write the equation of the graph presented below in the form $f(x) = ax^2 + bx + c$, assuming $a = 1$ or $a = -1$. Then, choose the intervals that a, b , and c belong to.



The solution is $f(x) = -x^2 + 8x - 20$

A. $a \in [0, 5]$, $b \in [4, 10]$, and $c \in [7, 13]$

$f(x) = x^2 + 8x + 12$, which corresponds to incorrectly using vertex form as $f(x) = a(x + h)^2 + k$ AND making a the opposite sign than it should be.

B. $a \in [-3, 0]$, $b \in [4, 10]$, and $c \in [-22, -17]$

* $f(x) = -x^2 + 8x - 20$, which is the correct option.

C. $a \in [-3, 0]$, $b \in [-10, -5]$, and $c \in [-14, -8]$

$f(x) = -x^2 - 8x - 12$, which corresponds to incorrectly using vertex form as $f(x) = a(x + h)^2 - k$.

D. $a \in [0, 5]$, $b \in [-10, -5]$, and $c \in [7, 13]$

$f(x) = x^2 - 8x + 12$, which corresponds to making a the opposite sign than it should be.

E. $a \in [-3, 0]$, $b \in [-10, -5]$, and $c \in [-22, -17]$


$f(x) = -x^2 - 8x - 20$, which corresponds to incorrectly using vertex form as $f(x) = a(x + h)^2 + k$.

General Comment: General Comments: When the graph is pointing up, $a = 1$. When the graph is pointing down, $a = -1$. Be sure to use Vertex Form: $y = a(x - h)^2 + k$.

3. Graph the equation below.

$$f(x) = -(x - 4)^2 - 19$$

The solution is



<p>A.</p>	<p>B.</p>
<p>C.</p>	<p>D.</p>
<p>E. None of the figures above.</p>	

A. Used the incorrect general form $f(x) = a(x + h)^2 + k$

B. This is the correct option.

C. Used the incorrect general form $f(x) = -a(x + h)^2 + k$

D. Used the incorrect general form $f(x) = -a(x - h)^2 + k$

General Comment: General Comments: Remember that Vertex Form is $y = a(x - h)^2 + k$, where the vertex is (h, k) .

4. Solve the quadratic equation below. Then, choose the intervals that the solutions belong to, with $x_1 \leq x_2$ (if they exist).

$$-14x^2 + 8x + 9 = 0$$

The solution is $x_1 = -0.565$ and $x_2 = 1.137$

A. $x_1 \in [-0.59, -0.31]$ and $x_2 \in [0.6, 1.9]$

* $x_1 = -0.565$ and $x_2 = 1.137$, which is the correct option.

B. $x_1 \in [-1.62, -0.68]$ and $x_2 \in [-0.1, 0.7]$

$x_1 = -1.137$ and $x_2 = 0.565$, which corresponds to writing the Quadratic Formula as $\frac{b \pm \sqrt{b^2 - 4ac}}{2a}$

C. $x_1 \in [-16.48, -15.25]$ and $x_2 \in [6.9, 10.1]$

$x_1 = -15.916$ and $x_2 = 7.916$, which corresponds to using the Quadratic Formula with $a = 1$

D. $x_1 \in [-24.23, -22.57]$ and $x_2 \in [23.9, 25.2]$

$x_1 = -23.547$ and $x_2 = 24.118$, which corresponds to writing the Quadratic Formula as $-\frac{b}{2a} \pm \sqrt{b^2 - 4ac}$.

E. There are no Real solutions.

Corresponds to getting a negative under the radical or believing that since the quadratic cannot be factored, it has no Real solutions.

General Comment: General Comments: This requires Quadratic Formula. Just be sure to use the correct formula and watch your signs.

0. Solve the quadratic equation below. Then, choose the intervals that the solutions x_1 and x_2 belong to, with $x_1 \leq x_2$.

$$10x^2 - 37x - 36 = 0$$

The solution is $x_1 = -0.800$ and $x_2 = 4.500$

A. $x_1 \in [-1, -0.56]$ and $x_2 \in [3.81, 4.76]$

* $x_1 = -0.800$ and $x_2 = 4.500$, which is the correct option. Obtained by solving the factored version $(5x + 4)(2x - 9)$

B. $x_1 \in [-8.46, -7.58]$ and $x_2 \in [44.2, 46.52]$

$x_1 = -8.000$ and $x_2 = 45.000$, which corresponds to solving the factored version $(x + 8)(x - 45)$

C. $x_1 \in [-4.38, -3.54]$ and $x_2 \in [-0.16, 1.8]$

$x_1 = -4.000$ and $x_2 = 0.900$, which corresponds to solving the factored version $(x + 4)(10x - 9)$

D. $x_1 \in [-1.75, -1.5]$ and $x_2 \in [0.98, 3.03]$

$x_1 = -1.600$ and $x_2 = 2.250$, which corresponds to solving the factored version $(5x + 8)(2x - 4)$

E. $x_1 \in [-0.36, -0.01]$ and $x_2 \in [13.28, 13.84]$

$x_1 = -0.267$ and $x_2 = 13.500$, which corresponds to solving the factored version $(5x + 1)(2x - 27)$

General Comment: General Comments: This question can be factored, but it may be faster to find the solutions via the Quadratic Equation.
