

1. For the information below, construct a linear model that describes the total time T spent on the path in terms of the distance of a particular part of the path *if we know that all parts of the path are equal length*.

A bicyclist is training for a race on a hilly path. Their bike keeps track of their speed at any time, but not the distance traveled. Their speed traveling up a hill is 3 mph, 9 mph when traveling down a hill, and 5 mph when traveling along a flat portion.

- A. $0.644D$
 - B. $135.000D$
 - C. $17.000D$
 - D. The model can be found with the information provided, but isn't options 1-3
 - E. The model cannot be found with the information provided.
-

2. For the information below, construct a linear model that describes the total time T spent on the path in terms of the distance of a particular part of the path *if we know that all parts of the path are equal length*.

A bicyclist is training for a race on a hilly path. Their bike keeps track of their speed at any time, but not the distance traveled. Their speed traveling up a hill is 5 mph, 10 mph when traveling down a hill, and 7 mph when traveling along a flat portion.

- A. $350.000D$
 - B. $0.443D$
 - C. $22.000D$
 - D. The model can be found with the information provided, but isn't options 1-3
 - E. The model cannot be found with the information provided.
-

3. A town has an initial population of 90000. The town's population for the next 9 years is provided below. Which type of function would be most appropriate to model the town's population?

Year	1	2	3	4	5	6	7	8	9
Pop	89840	89360	87440	79760	49040	0	0	0	0

- A. Non-Linear Power
- B. Linear
- C. Exponential
- D. Logarithmic
- E. None of the above

4. What is the **best** way to describe the domain of the scenario below?

Chemists commonly create a solution by mixing two products of differing concentrations together. A 10% and 30% solution can make an acid solution of some value between these, such as a 24% acid solution. The chemist wants to make differing solution percentages of 7 liters each.

- A. Subset of the Natural numbers
- B. Proper subset of the Real numbers
- C. Subset of the Integers
- D. Subset of the Rational numbers
- E. There is no restricted domain in this scenario

5. Using the situation below, construct a linear model that describes the cost of the coffee beans $C(h)$ in terms of the weight of the low-quality coffee beans h .

Veronica needs to prepare 230 of blended coffee beans selling for \$5.55 per pound. She has a high-quality bean that sells for \$6.77 a pound and a low-quality bean that sells for \$3.97 a pound.

- A. $C(h) = 5.37h$
- B. $C(h) = 3.97h$
- C. $C(h) = -2.80h + 1557.10$

D. $C(h) = 2.80h + 913.10$

E. None of the above.

-
6. What is the **best** way to describe the domain of the scenario below?

Bridges on highways often have expansion joints, which are small gaps in the roadway between one bridge section and the next. The gaps are put there so the bridge will have room to expand when the weather gets hot. Assume the gap width varies constantly with the temperature.

Suppose a bridge has a gap of 1.3 cm when the temperature is 22 degrees C and that the gap narrows to 0.9 cm when the temperature warms to 30 degrees C.

- A. Subset of the Integers
- B. Subset of the Rational numbers
- C. There is no restricted domain in this scenario
- D. Subset of the Natural numbers
- E. Proper subset of the Real numbers

-
7. For the information provided below, construct a linear model that describes her total costs, C , as a function of the number of months, x she is at UF.

Aubrey is a college student going into her first year at UF. She will receive Bright Futures, which covers her tuition plus a \$800 educational expense each year. Before college, Aubrey saved up \$10000. She knows she will need to pay \$1200 in rent a month, \$70 for food a week, and \$64 in other weekly expenses.

- A. $C(x) = 1334x$
- B. $C(x) = 1736x$
- C. $C(x) = 1736$
- D. $C(x) = 1334$

E. None of the above.

-
8. Using the situation below, construct a linear model that describes the cost of the coffee beans $C(h)$ in terms of the weight of the high-quality coffee beans h .

Veronica needs to prepare 200 of blended coffee beans selling for \$5.72 per pound. She has a high-quality bean that sells for \$6.79 a pound and a low-quality bean that sells for \$4.77 a pound.

- A. $C(h) = 6.79h$
 B. $C(h) = -2.02h + 1358.00$
 C. $C(h) = 5.78h$
 D. $C(h) = 2.02h + 954.00$
 E. None of the above.

-
9. A town has an initial population of 50000. The town's population for the next 9 years is provided below. Which type of function would be most appropriate to model the town's population?

Year	1	2	3	4	5	6	7	8	9
Pop	50000	49986	49978	49972	49967	49964	49961	49958	49956

- A. Exponential
 B. Logarithmic
 C. Linear
 D. Non-Linear Power
 E. None of the above

-
10. For the information provided below, construct a linear model that describes her total income, I , as a function of the number of months, x she is at UF.

Aubrey is a college student going into her first year at UF. She will receive Bright Futures, which covers her tuition plus a \$600 educational expense each year. Before college, Aubrey saved up \$11000. She knows she will need to pay \$800 in rent a month, \$70 for food a week, and \$32 in other weekly expenses.

- A. $I(x) = 902x$
- B. $I(x) = 1208$
- C. $I(x) = 902$
- D. $I(x) = 1208x$
- E. None of the above.

-
11. For the information below, construct a linear model that describes the total time T spent on the path in terms of the distance of a particular part of the path *if we know that all parts of the path are equal length.*

A bicyclist is training for a race on a hilly path. Their bike keeps track of their speed at any time, but not the distance traveled. Their speed traveling up a hill is 6 mph, 11 mph when traveling down a hill, and 8 mph when traveling along a flat portion.

- A. $528.000D$
- B. $0.383D$
- C. $25.000D$
- D. The model can be found with the information provided, but isn't options 1-3
- E. The model cannot be found with the information provided.

-
12. For the information provided below, construct a linear model that describes the total distance of the path, D , in terms of the time spent on a particular path *if we know that all parts of the path are equal length.*

A bicyclist is training for a race on a hilly path. Their bike keeps track of their speed at any time, but not the distance traveled. Their

speed traveling up a hill is 2 mph, 7 mph when traveling down a hill, and 5 mph when traveling along a flat portion.

- A. $70t$
- B. $0.843t$
- C. $14t$
- D. The model can be found with the information provided, but isn't options 1-3
- E. The model cannot be found with the information provided.

13. A town has an initial population of 100000. The town's population for the next 9 years is provided below. Which type of function would be most appropriate to model the town's population?

Year	1	2	3	4	5	6	7	8	9
Pop	99960	99920	99880	99840	99800	99760	99720	99680	99640

- A. Exponential
- B. Logarithmic
- C. Non-Linear Power
- D. Linear
- E. None of the above

14. What is the **best** way to describe the domain of the scenario below?

Bridges on highways often have expansion joints, which are small gaps in the roadway between one bridge section and the next. The gaps are put there so the bridge will have room to expand when the weather gets hot. Assume the gap width varies constantly with the temperature.

Suppose a bridge has a gap of 1.3 cm when the temperature is 22 degrees C and that the gap narrows to 0.9 cm when the temperature warms to 30 degrees C.

- A. Subset of the Integers
- B. Subset of the Natural numbers

- C. Subset of the Rational numbers
 - D. There is no restricted domain in this scenario
 - E. Proper subset of the Real numbers
-

15. Using the situation below, construct a linear model that describes the cost of the coffee beans $C(h)$ in terms of the weight of the low-quality coffee beans h .

Veronica needs to prepare 150 of blended coffee beans selling for \$3.66 per pound. She has a high-quality bean that sells for \$4.20 a pound and a low-quality bean that sells for \$2.43 a pound.

- A. $C(h) = 1.77h + 364.50$
 - B. $C(h) = 3.32h$
 - C. $C(h) = 2.43h$
 - D. $C(h) = -1.77h + 630.00$
 - E. None of the above.
-

16. What is the **best** way to describe the domain of the scenario below?

Hannah plans to pay off a no-interest loan from her parents. Her loan balance is \$1,000. She plans to pay \$35 at the end of every week until her balance is \$0. How many weeks will it be until she has paid off her loan?

- A. Subset of the Natural numbers
 - B. There is no restricted domain in this scenario
 - C. Subset of the Rational numbers
 - D. Proper subset of the Real numbers
 - E. Subset of the Integers
-

17. For the information provided below, construct a linear model that describes her total income, I , as a function of the number of months, x she is at UF.

Aubrey is a college student going into her first year at UF. She will receive Bright Futures, which covers her tuition plus a \$800 educational expense each year. Before college, Aubrey saved up \$8000. She knows she will need to pay \$900 in rent a month, \$80 for food a week, and \$40 in other weekly expenses.

- A. $I(x) = 8800x$
- B. $I(x) = 800x + 8000$
- C. $I(x) = 8800$
- D. $I(x) = 8000x + 800$
- E. None of the above.

-
18. Using the situation below, construct a linear model that describes the cost of the coffee beans $C(h)$ in terms of the weight of the high-quality coffee beans h .

Veronica needs to prepare 100 of blended coffee beans selling for \$3.27 per pound. She has a high-quality bean that sells for \$3.77 a pound and a low-quality bean that sells for \$2.73 a pound.

- A. $C(h) = 3.77h$
- B. $C(h) = -1.04h + 377.00$
- C. $C(h) = 3.25h$
- D. $C(h) = 1.04h + 273.00$
- E. None of the above.

-
19. A town has an initial population of 70000. The town's population for the next 9 years is provided below. Which type of function would be most appropriate to model the town's population?

Year	1	2	3	4	5	6	7	8	9
Pop	69970	69940	69910	69880	69850	69820	69790	69760	69730

A. Non-Linear Power

B. Linear

C. Logarithmic

D. Exponential

E. None of the above

20. For the information provided below, construct a linear model that describes her total income, I , as a function of the number of months, x she is at UF.

Aubrey is a college student going into her first year at UF. She will receive Bright Futures, which covers her tuition plus a \$1000 educational expense each year. Before college, Aubrey saved up \$9000. She knows she will need to pay \$800 in rent a month, \$40 for food a week, and \$56 in other weekly expenses.

A. $I(x) = 9000x + 1000$

B. $I(x) = 10000$

C. $I(x) = 1000x + 9000$

D. $I(x) = 10000x$

E. None of the above.

21. For the information provided below, construct a linear model that describes the total distance of the path, D , in terms of the time spent on a particular path *if we know that the time spent on each path was equal.*

A bicyclist is training for a race on a hilly path. Their bike keeps track of their speed at any time, but not the distance traveled. Their speed traveling up a hill is 5 mph, 10 mph when traveling down a hill, and 6 mph when traveling along a flat portion.

A. $300t$

B. $0.467t$

C. $21t$

D. The model can be found with the information provided, but isn't options 1-3

E. The model cannot be found with the information provided.

22. For the information below, construct a linear model that describes the total time T spent on the path in terms of the distance of a particular part of the path *if we know that all parts of the path are equal length*.

A bicyclist is training for a race on a hilly path. Their bike keeps track of their speed at any time, but not the distance traveled. Their speed traveling up a hill is 2 mph, 8 mph when traveling down a hill, and 5 mph when traveling along a flat portion.

A. $15.000D$

B. $80.000D$

C. $0.825D$

D. The model can be found with the information provided, but isn't options 1-3

E. The model cannot be found with the information provided.

23. A town has an initial population of 30000. The town's population for the next 9 years is provided below. Which type of function would be most appropriate to model the town's population?

Year	1	2	3	4	5	6	7	8	9
Pop	30000	30034	30054	30069	30080	30089	30097	30103	30109

A. Non-Linear Power

B. Logarithmic

C. Linear

D. Exponential

E. None of the above

24. What is the **best** way to describe the domain of the scenario below?

Bridges on highways often have expansion joints, which are small gaps in the roadway between one bridge section and the next. The gaps are put there so the bridge will have room to expand when the weather gets hot. Assume the gap width varies constantly with the temperature.

Suppose a bridge has a gap of 1.3 cm when the temperature is 22 degrees C and that the gap narrows to 0.9 cm when the temperature warms to 30 degrees C.

- A. Subset of the Natural numbers
 - B. Proper subset of the Real numbers
 - C. There is no restricted domain in this scenario
 - D. Subset of the Integers
 - E. Subset of the Rational numbers
-

25. Using the situation below, construct a linear model that describes the cost of the coffee beans $C(h)$ in terms of the weight of the high-quality coffee beans h .

Veronica needs to prepare 100 of blended coffee beans selling for \$5.80 per pound. She has a high-quality bean that sells for \$6.93 a pound and a low-quality bean that sells for \$4.78 a pound.

- A. $C(h) = -2.15h + 693.00$
 - B. $C(h) = 2.15h + 478.00$
 - C. $C(h) = 6.93h$
 - D. $C(h) = 5.86h$
 - E. None of the above.
-

26. What is the **best** way to describe the domain of the scenario below?

Fred is a store manager at Publix. The store normally orders two pallets of water bottles a week and sells 1000 bottles per day. However, a hurricane is coming and Fred expects water bottle sales to increase tenfold for three days, then decrease by half of normal sales for four days. How many more pallets of water bottles should Fred order the week before the hurricane?

- A. Subset of the Rational numbers
- B. There is no restricted domain in this scenario
- C. Subset of the Natural numbers
- D. Subset of the Integers
- E. Proper subset of the Real numbers

-
27. For the information provided below, construct a linear model that describes her total costs, C , as a function of the number of months, x she is at UF.

Aubrey is a college student going into her first year at UF. She will receive Bright Futures, which covers her tuition plus a \$600 educational expense each year. Before college, Aubrey saved up \$11000. She knows she will need to pay \$1100 in rent a month, \$50 for food a week, and \$48 in other weekly expenses.

- A. $C(x) = 1492x$
- B. $C(x) = 1198$
- C. $C(x) = 1198x$
- D. $C(x) = 1492$
- E. None of the above.

-
28. Using the situation below, construct a linear model that describes the cost of the coffee beans $C(h)$ in terms of the weight of the high-quality coffee beans h .

Veronica needs to prepare 160 of blended coffee beans selling for \$3.55 per pound. She has a high-quality bean that sells for \$4.27 a pound and a low-quality bean that sells for \$2.73 a pound.

- A. $C(h) = 4.27h$
- B. $C(h) = 1.54h + 436.80$
- C. $C(h) = 3.50h$
- D. $C(h) = -1.54h + 683.20$
- E. None of the above.

29. A town has an initial population of 30000. The town's population for the next 9 years is provided below. Which type of function would be most appropriate to model the town's population?

Year	1	2	3	4	5	6	7	8	9
Pop	29960	29920	29880	29840	29800	29760	29720	29680	29640

- A. Logarithmic
- B. Non-Linear Power
- C. Linear
- D. Exponential
- E. None of the above

30. For the information provided below, construct a linear model that describes her total income, I , as a function of the number of months, x she is at UF.

Aubrey is a college student going into her first year at UF. She will receive Bright Futures, which covers her tuition plus a \$600 educational expense each year. Before college, Aubrey saved up \$11000. She knows she will need to pay \$1000 in rent a month, \$80 for food a week, and \$64 in other weekly expenses.

- A. $I(x) = 11600$

B. $I(x) = 600x + 11000$

C. $I(x) = 11000x + 600$

D. $I(x) = 11600x$

E. None of the above.