1. A town has an initial population of 60000. The town's population for the next 9 years is provided below. Which type of function would be most appropriate to model the town's population?

Year	1	2	3	4	5	6	7	8	9
Pop	60000	59979	59967	59958	59951	59946	59941	59937	59934

- A. Logarithmic
- B. Exponential
- C. Linear
- D. Non-Linear Power
- E. None of the above
- 2. What is the **best** way to describe the domain of the scenario below?

Bridges on highways often have expansion joints, which are small gaps in the roadway between one bridge section and the next. The gaps are put there so the bridge will have room to expand when the weather gets hot. Assume the gap width varies constantly with the temperature.

Suppose a bridge has a gap of 1.3 cm when the temperature is 22 degrees C and that the gap narrows to 0.9 cm when the temperature warms to 30 degrees C.

- A. Subset of the Natural numbers
- B. Subset of the Rational numbers
- C. Proper subset of the Real numbers
- D. There is no restricted domain in this scenario
- E. Subset of the Integers
- 3. For the information below, construct a linear model that describes the total time T spent on the path in terms of the distance of a particular part of the path if we know that all parts of the path are equal length.

A bicyclist is training for a race on a hilly path. Their bike keeps track of their speed at any time, but not the distance traveled. Their

5493-4176 Summer C 2021

speed traveling up a hill is 5 mph, 10 mph when traveling down a hill, and 7 mph when traveling along a flat portion.

- A. 22.000D
- B. 0.443*D*
- C. 350.000*D*
- D. The model can be found with the information provided, but isn't options 1-3
- E. The model cannot be found with the information provided.
- 4. What is the **best** way to describe the domain of the scenario below?

Veronica needs to prepare 170 lbs of blended coffee beans to sell for \$4.71 per pound. She has a high-quality bean that sells for \$6.00 a pound and a low-quality been that sells for \$3.25 a pound.

- A. Subset of the Rational numbers
- B. There is no restricted domain in this scenario
- C. Subset of the Integers
- D. Subset of the Natural numbers
- E. Proper subset of the Real numbers
- 5. For the information provided below, construct a linear model that describes her total budget, B, as a function of the number of months, x she is at UF.

Aubrey is a college student going into her first year at UF. She will receive Bright Futures, which covers her tuition plus a \$400 educational expense each year. Before college, Aubrey saved up \$9000. She knows she will need to pay \$800 in rent a month, \$70 for food a week, and \$64 in other weekly expenses.

- A. B(x) = 8466x
- B. B(x) = 8064x

5493-4176 Summer C 2021

- C. B(x) = 9400 1336x
- D. B(x) = 9400 934x
- E. None of the above.
- 6. For the information below, construct a linear model that describes the total time T spent on the path in terms of the distance of a particular part of the path if we know that all parts of the path are equal length.

A bicyclist is training for a race on a hilly path. Their bike keeps track of their speed at any time, but not the distance traveled. Their speed traveling up a hill is 4 mph, 9 mph when traveling down a hill, and 7 mph when traveling along a flat portion.

- A. 252.000*D*
- B. 0.504*D*
- C. 20.000*D*
- D. The model can be found with the information provided, but isn't options 1-3
- E. The model cannot be found with the information provided.
- 7. A town has an initial population of 60000. The town's population for the next 9 years is provided below. Which type of function would be most appropriate to model the town's population?

Year	1	2	3	4	5	6	7	8	9
Pop	59980	59960	59940	59920	59900	59880	59860	59840	59820

- A. Exponential
- B. Linear
- C. Logarithmic
- D. Non-Linear Power
- E. None of the above

8. Using the situation below, construct a linear model that describes the cost of the coffee beans C(h) in terms of the weight of the low-quality coffee beans h.

Veronica needs to prepare 110 of blended coffee beans selling for \$5.71 per pound. She has a high-quality bean that sells for \$6.89 a pound and a low-quality bean that sells for \$4.48 a pound.

A.
$$C(h) = 2.41h + 492.80$$

B.
$$C(h) = 4.48h$$

C.
$$C(h) = 5.69h$$

D.
$$C(h) = -2.41h + 757.90$$

- E. None of the above.
- 9. Using the situation below, construct a linear model that describes the cost of the coffee beans C(h) in terms of the weight of the high-quality coffee beans h.

Veronica needs to prepare 70 of blended coffee beans selling for \$5.22 per pound. She has a high-quality bean that sells for \$6.79 a pound and a low-quality bean that sells for \$4.69 a pound.

A.
$$C(h) = -2.10h + 475.30$$

B.
$$C(h) = 2.10h + 328.30$$

C.
$$C(h) = 6.79h$$

D.
$$C(h) = 5.74h$$

- E. None of the above.
- 10. For the information provided below, construct a linear model that describes her total income, I, as a function of the number of months, x she is at UF.

Aubrey is a college student going into her first year at UF. She will receive Bright Futures, which covers her tuition plus a \$1000

educational expense each year. Before college, Aubrey saved up \$6000. She knows she will need to pay \$700 in rent a month, \$40 for food a week, and \$56 in other weekly expenses.

A.
$$I(x) = 6000x + 1000$$

B.
$$I(x) = 7000x$$

C.
$$I(x) = 7000$$

D.
$$I(x) = 1000x + 6000$$

E. None of the above.

5493-4176 Summer C 2021