
1. A town has an initial population of 60000. The town's population for the next 10 years is provided below. Which type of function would be most appropriate to model the town's population?

Year	1	2	3	4	5	6	7	8	9
Pop	60000	60027	60043	60055	60064	60071	60077	60083	60087

- A. Non-Linear Power
- B. Logarithmic
- C. Exponential
- D. Linear
- E. None of the above
- 2. The temperature of an object, T, in a different surrounding temperature T_s will behave according to the formula $T(t) = Ae^{kt} + T_s$, where t is minutes, A is a constant, and k is a constant. Use this formula and the situation below to construct a model that describes the uranium's temperature, T, based on the amount of time t (in minutes) that have passed. Choose the correct constant k from the options below.

Uranium is taken out of the reactor with a temperature of 200° C and is placed into a 18° C bath to cool. After 15 minutes, the uranium has cooled to 149° C.

- A. k = -0.02821
- B. k = -0.05239
- C. k = -0.05160
- D. k = -0.08354
- E. None of the above
- 3. Determine the appropriate model for the graph of points below.



- A. Non-linear Power model
- B. Linear model
- C. Exponential model
- D. Logarithmic model
- E. None of the above
- 4. Using the scenario below, model the population of bacteria α in terms of the number of minutes, t that pass. Then, choose the correct approximate (rounded to the nearest minute) replication rate of bacteria- α .

A newly discovered bacteria, α, is being examined in a lab. The lab started with a petri dish of 2 bacteria-α. After 2 hours, the petri dish has 133 bacteria-α. Based on similar bacteria, the lab believes bacteria-α triples after some undetermined number of minutes.

- A. About 188 minutes
- B. About 263 minutes
- C. About 31 minutes
- D. About 43 minutes
- E. None of the above

5170-5105 Summer C 2021

5. A town has an initial population of 70000. The town's population for the next 10 years is provided below. Which type of function would be most appropriate to model the town's population?

Year	1	2	3	4	5	6	7	8	9
Pop	69964	69924	69884	69844	69804	69764	69724	69684	69644

- A. Linear
- B. Logarithmic
- C. Exponential
- D. Non-Linear Power
- E. None of the above
- 6. Using the scenario below, model the situation using an exponential function and a base of $\frac{1}{2}$. Then, solve for the half-life of the element, rounding to the nearest day.


The half-life of an element is the amount of time it takes for the element to decay to half of its initial starting amount. There is initially 979 grams of element X and after 5 years there is 97 grams remaining.

- A. About 2190 days
- B. About 730 days
- C. About 365 days
- D. About 1 day
- E. None of the above
- 7. The temperature of an object, T, in a different surrounding temperature T_s will behave according to the formula $T(t) = Ae^{kt} + T_s$, where t is minutes, A is a constant, and k is a constant. Use this formula and the situation below to construct a model that describes the uranium's

temperature, T, based on the amount of time t (in minutes) that have passed. Choose the correct constant k from the options below.

Uranium is taken out of the reactor with a temperature of 140° C and is placed into a 18° C bath to cool. After 22 minutes, the uranium has cooled to 75° C.

- A. k = -0.03166
- B. k = -0.03459
- C. k = -0.03093
- D. k = -0.04085
- E. None of the above
- 8. Determine the appropriate model for the graph of points below.



- A. Logarithmic model
- B. Exponential model
- C. Non-linear Power model
- D. Linear model
- E. None of the above
- 9. Using the scenario below, model the situation using an exponential

function and a base of $\frac{1}{2}$. Then, solve for the half-life of the element, rounding to the nearest day.

The half-life of an element is the amount of time it takes for the element to decay to half of its initial starting amount. There is initially 920 grams of element X and after 7 years there is 131 grams remaining.

- A. About 730 days
- B. About 1095 days
- C. About 1 day
- D. About 2920 days
- E. None of the above
- 10. Using the scenario below, model the population of bacteria α in terms of the number of minutes, t that pass. Then, choose the correct approximate (rounded to the nearest minute) replication rate of bacteria- α .

A newly discovered bacteria, α, is being examined in a lab. The lab started with a petri dish of 3 bacteria-α. After 3 hours, the petri dish has 889 bacteria-α. Based on similar bacteria, the lab believes bacteria-α triples after some undetermined number of minutes.

- A. About 47 minutes
- B. About 21 minutes
- C. About 284 minutes
- D. About 131 minutes
- E. None of the above

5170-5105 Summer C 2021