

1. Add the following functions, then choose the domain of the resulting function from the list below.

$$f(x) = 3x^3 + 8x + 5 \text{ and } g(x) = \sqrt{-5x - 18}$$

- A. The domain is all Real numbers less than or equal to $x = a$, where $a \in [-7.6, 1.4]$
 - B. The domain is all Real numbers except $x = a$, where $a \in [-8.6, -3.6]$
 - C. The domain is all Real numbers greater than or equal to $x = a$, where $a \in [4, 9]$
 - D. The domain is all Real numbers except $x = a$ and $x = b$, where $a \in [3.75, 10.75]$ and $b \in [1.2, 8.2]$
 - E. The domain is all Real numbers.
-

2. Find the inverse of the function below. Then, evaluate the inverse at $x = 7$ and choose the interval that $f^{-1}(7)$ belongs to.

$$f(x) = \ln(x - 5) - 3$$

- A. $f^{-1}(7) \in [22026.47, 22032.47]$
 - B. $f^{-1}(7) \in [162748.79, 162754.79]$
 - C. $f^{-1}(7) \in [58.6, 62.6]$
 - D. $f^{-1}(7) \in [22015.47, 22023.47]$
 - E. $f^{-1}(7) \in [-2.61, 10.39]$
-

3. Determine whether the function below is 1-1.

$$f(x) = (4x + 22)^3$$

- A. No, because the domain of the function is not $(-\infty, \infty)$.
- B. Yes, the function is 1-1.
- C. No, because the range of the function is not $(-\infty, \infty)$.

- D. No, because there is an x -value that goes to 2 different y -values.
E. No, because there is a y -value that goes to 2 different x -values.
-

4. Find the inverse of the function below. Then, evaluate the inverse at $x = 4$ and choose the interval that $f^{-1}(4)$ belongs to.

$$f(x) = e^{x-2} - 2$$

- A. $f^{-1}(4) \in [-0.26, 0.56]$
B. $f^{-1}(4) \in [-2.74, -0.56]$
C. $f^{-1}(4) \in [-0.26, 0.56]$
D. $f^{-1}(4) \in [-2.74, -0.56]$
E. $f^{-1}(4) \in [3.68, 4.18]$
-

5. Determine whether the function below is 1-1.

$$f(x) = 9x^2 + 126x + 441$$

- A. No, because there is a y -value that goes to 2 different x -values.
B. No, because the range of the function is not $(-\infty, \infty)$.
C. No, because there is an x -value that goes to 2 different y -values.
D. No, because the domain of the function is not $(-\infty, \infty)$.
E. Yes, the function is 1-1.
-

6. Choose the interval below that f composed with g at $x = 1$ is in.

$$f(x) = 3x^3 + 2x^2 - 4x \text{ and } g(x) = 3x^3 + 2x^2 - 4x + 1$$

- A. $(f \circ g)(1) \in [-13, -3]$
B. $(f \circ g)(1) \in [21, 25]$
C. $(f \circ g)(1) \in [16, 21]$

- D. $(f \circ g)(1) \in [1, 4]$
E. It is not possible to compose the two functions.
-

7. Choose the interval below that f composed with g at $x = -1$ is in.

$$f(x) = x^3 - 2x^2 - x + 2 \text{ and } g(x) = -x^3 + 4x^2 + x$$

- A. $(f \circ g)(-1) \in [35, 40]$
B. $(f \circ g)(-1) \in [29, 31]$
C. $(f \circ g)(-1) \in [-4, 1]$
D. $(f \circ g)(-1) \in [7, 18]$
E. It is not possible to compose the two functions.
-

8. Find the inverse of the function below (if it exists). Then, evaluate the inverse at $x = -14$ and choose the interval that $f^{-1}(-14)$ belongs to.

$$f(x) = \sqrt[3]{5x - 4}$$

- A. $f^{-1}(-14) \in [547, 548.6]$
B. $f^{-1}(-14) \in [-551.3, -548.6]$
C. $f^{-1}(-14) \in [-548.6, -546.7]$
D. $f^{-1}(-14) \in [549, 552.1]$
E. The function is not invertible for all Real numbers.
-

9. Find the inverse of the function below (if it exists). Then, evaluate the inverse at $x = -15$ and choose the interval that $f^{-1}(-15)$ belongs to.

$$f(x) = 3x^2 - 4$$

- A. $f^{-1}(-15) \in [2.27, 2.72]$
B. $f^{-1}(-15) \in [2.73, 3.19]$

- C. $f^{-1}(-15) \in [5.61, 6.24]$
 - D. $f^{-1}(-15) \in [1.77, 2.01]$
 - E. The function is not invertible for all Real numbers.
-

10. Subtract the following functions, then choose the domain of the resulting function from the list below.

$$f(x) = \frac{3}{3x - 14} \text{ and } g(x) = 2x^2 + 3x + 7$$

- A. The domain is all Real numbers except $x = a$, where $a \in [3.67, 13.67]$
 - B. The domain is all Real numbers greater than or equal to $x = a$, where $a \in [-8.67, -1.67]$
 - C. The domain is all Real numbers less than or equal to $x = a$, where $a \in [-3, 0]$
 - D. The domain is all Real numbers except $x = a$ and $x = b$, where $a \in [2.67, 6.67]$ and $b \in [-4.67, 1.33]$
 - E. The domain is all Real numbers.
-