

1. Find the inverse of the function below. Then, evaluate the inverse at $x = 6$ and choose the interval that $f^{-1}(6)$ belongs to.

$$f(x) = e^{x-4} - 4$$

- A. $f^{-1}(6) \in [-7.31, -2.31]$
 - B. $f^{-1}(6) \in [-1.7, 3.3]$
 - C. $f^{-1}(6) \in [-7.31, -2.31]$
 - D. $f^{-1}(6) \in [6.3, 9.3]$
 - E. $f^{-1}(6) \in [-1.7, 3.3]$
-

2. Multiply the following functions, then choose the domain of the resulting function from the list below.

$$f(x) = 8x + 2 \text{ and } g(x) = \sqrt{3x + 14}$$

- A. The domain is all Real numbers less than or equal to $x = a$, where $a \in [-2.17, 1.83]$
 - B. The domain is all Real numbers greater than or equal to $x = a$, where $a \in [-5.67, -2.67]$
 - C. The domain is all Real numbers except $x = a$, where $a \in [2.25, 7.25]$
 - D. The domain is all Real numbers except $x = a$ and $x = b$, where $a \in [4.67, 12.67]$ and $b \in [6.67, 10.67]$
 - E. The domain is all Real numbers.
-

3. Find the inverse of the function below (if it exists). Then, evaluate the inverse at $x = -14$ and choose the interval that $f^{-1}(-14)$ belongs to.

$$f(x) = 5x^2 - 4$$

- A. $f^{-1}(-14) \in [4.27, 4.55]$
- B. $f^{-1}(-14) \in [1.76, 2.08]$
- C. $f^{-1}(-14) \in [1.22, 1.83]$

- D. $f^{-1}(-14) \in [6.66, 7.48]$
- E. The function is not invertible for all Real numbers.

-
4. Subtract the following functions, then choose the domain of the resulting function from the list below.

$$f(x) = 7x + 8 \text{ and } g(x) = \sqrt{-4x + 11}$$

- A. The domain is all Real numbers greater than or equal to $x = a$, where $a \in [-5.6, -3.6]$
- B. The domain is all Real numbers less than or equal to $x = a$, where $a \in [-0.25, 3.75]$
- C. The domain is all Real numbers except $x = a$, where $a \in [-11.2, -6.2]$
- D. The domain is all Real numbers except $x = a$ and $x = b$, where $a \in [-9.8, -4.8]$ and $b \in [-4.2, 1.8]$
- E. The domain is all Real numbers.

-
5. Choose the interval below that f composed with g at $x = 1$ is in.

$$f(x) = -x^3 + 2x^2 + x - 3 \text{ and } g(x) = 2x^3 + 2x^2 - 2x$$

- A. $(f \circ g)(1) \in [1.69, 3.52]$
- B. $(f \circ g)(1) \in [-2.82, -0.76]$
- C. $(f \circ g)(1) \in [3.04, 5.12]$
- D. $(f \circ g)(1) \in [7.85, 8.59]$
- E. It is not possible to compose the two functions.

-
6. Choose the interval below that f composed with g at $x = 2$ is in.

$$f(x) = -2x^3 + 3x^2 + 2x \text{ and } g(x) = -2x^3 + 2x^2 + 4x$$

- A. $(f \circ g)(2) \in [-0.5, 0.1]$

- B. $(f \circ g)(2) \in [-0.5, 0.1]$
 - C. $(f \circ g)(2) \in [9.4, 11]$
 - D. $(f \circ g)(2) \in [5.2, 9.2]$
 - E. It is not possible to compose the two functions.
-

7. Determine whether the function below is 1-1.

$$f(x) = -20x^2 - 247x - 713$$

- A. No, because there is a y -value that goes to 2 different x -values.
 - B. Yes, the function is 1-1.
 - C. No, because the domain of the function is not $(-\infty, \infty)$.
 - D. No, because there is an x -value that goes to 2 different y -values.
 - E. No, because the range of the function is not $(-\infty, \infty)$.
-

8. Determine whether the function below is 1-1.

$$f(x) = 36x^2 + 300x + 625$$

- A. No, because the domain of the function is not $(-\infty, \infty)$.
 - B. No, because the range of the function is not $(-\infty, \infty)$.
 - C. Yes, the function is 1-1.
 - D. No, because there is a y -value that goes to 2 different x -values.
 - E. No, because there is an x -value that goes to 2 different y -values.
-

9. Find the inverse of the function below. Then, evaluate the inverse at $x = 10$ and choose the interval that $f^{-1}(10)$ belongs to.

$$f(x) = \ln(x - 3) + 5$$

- A. $f^{-1}(10) \in [442417.39, 442419.39]$

- B. $f^{-1}(10) \in [150.41, 156.41]$
 - C. $f^{-1}(10) \in [140.41, 151.41]$
 - D. $f^{-1}(10) \in [3269014.37, 3269023.37]$
 - E. $f^{-1}(10) \in [1097.63, 1104.63]$
-

10. Find the inverse of the function below (if it exists). Then, evaluate the inverse at $x = 10$ and choose the interval that $f^{-1}(10)$ belongs to.

$$f(x) = \sqrt[3]{5x + 4}$$

- A. $f^{-1}(10) \in [-199.5, -199.1]$
 - B. $f^{-1}(10) \in [198.4, 199.8]$
 - C. $f^{-1}(10) \in [199.5, 202.3]$
 - D. $f^{-1}(10) \in [-203.4, -200.2]$
 - E. The function is not invertible for all Real numbers.
-