

This key should allow you to understand why you choose the option you did (beyond just getting a question right or wrong). More instructions on how to use this key can be found [here](#).

If you have a suggestion to make the keys better, please fill out the short survey [here](#).

*Note: This key is auto-generated and may contain issues and/or errors. The keys are reviewed after each exam to ensure grading is done accurately. If there are issues (like duplicate options), they are noted in the offline gradebook. The keys are a work-in-progress to give students as many resources to improve as possible.*

1. A town has an initial population of 40000. The town's population for the next 10 years is provided below. Which type of function would be most appropriate to model the town's population?

Year	1	2	3	4	5	6	7	8	9
Pop	40000	40027	40043	40055	40064	40071	40077	40083	40087

The solution is Logarithmic, which is option B.

A. Exponential

This suggests the fastest of growths that we know.

B. Logarithmic

This suggests the slowest of growths that we know.

C. Linear

This suggests a constant growth. You would be able to add or subtract the same amount year-to-year if this is the correct answer.

D. Non-Linear Power

This suggests a growth faster than constant but slower than exponential.

E. None of the above

Please contact the coordinator to discuss why you believe none of the options model the population.

**General Comment:** We are trying to compare the growth rate of the population. Growth rates can be characterized from slowest to fastest as: logarithmic, indirect, linear, direct, exponential. The best way to approach this is to first compare it to linear (is it linear, faster than linear, or slower than linear)? If faster, is it as fast as exponential? If slower, is it as slow as logarithmic?

2. The temperature of an object,  $T$ , in a different surrounding temperature  $T_s$  will behave according to the formula  $T(t) = Ae^{kt} + T_s$ , where  $t$  is minutes,  $A$  is a constant, and  $k$  is a constant. Use this formula and the situation below to construct a model that describes the uranium's temperature,  $T$ , based on the amount of time  $t$  (in minutes) that have passed. Choose the correct constant  $k$  from the options below.

*Uranium is taken out of the reactor with a temperature of  $110^\circ\text{C}$  and is placed into a  $17^\circ\text{C}$  bath to cool. After 37 minutes, the uranium has cooled to  $68^\circ\text{C}$ .*

The solution is  $k = -0.01624$ , which is option D.

A.  $k = -0.01864$

This uses  $A$  as the initial temperature and solves for  $k$  incorrectly.

B.  $k = -0.02077$

This uses  $A$  as the initial temperature and solves for  $k$  correctly.

C.  $k = -0.01921$

This uses  $A$  correctly but solves for  $k$  incorrectly.

D.  $k = -0.01624$

\* This is the correct option.


E. None of the above

If you chose this, please contact the coordinator to discuss why you believe none of the other answers are correct.

**General Comment:** The initial temperature is when  $t = 0$ . Unlike power models, that means  $A$  is not the initial temperature!

---

3. Determine the appropriate model for the graph of points below.



The solution is Linear model, which is option A.

A. Linear model

For this to be the correct option, we need to see a mostly straight line of points.

B. Exponential model

For this to be the correct option, we want an extremely slow change early, then a rapid change later.

C. Logarithmic model

For this to be the correct option, we want a rapid change early, then an extremely slow change later.

D. Non-linear Power model

For this to be the correct option, we need to see a polynomial or rational shape.

E. None of the above

For this to be the correct option, we want to see no pattern in the points.

**General Comment:** This question is testing if you can associate the models with their graphical representation. If you are having trouble, go back to the corresponding Core module to learn about the specific function you are having trouble recognizing.

---

4. Using the scenario below, model the population of bacteria  $\alpha$  in terms of the number of minutes,  $t$  that pass. Then, choose the correct approximate (*rounded to the nearest minute*) replication rate of bacteria- $\alpha$ .

*A newly discovered bacteria,  $\alpha$ , is being examined in a lab. The lab started with a petri dish of 3 bacteria- $\alpha$ . After 2 hours, the petri dish has 105 bacteria- $\alpha$ . Based on similar bacteria, the lab believes bacteria- $\alpha$  triples after some undetermined number of minutes.*

The solution is None of the above, which is option E.

- A. About 46 minutes

This uses the wrong base and does not solve for the constant correctly.

- B. About 140 minutes

This uses the wrong base and solves for the constant correctly but converted incorrectly.

- C. About 23 minutes

This uses the wrong base.

- D. About 276 minutes

This uses the wrong base, does not solve for the constant correctly, AND converted incorrectly.

- E. None of the above

\* This is the correct option as all other options used the wrong base in their model.

**General Comment:** Your model should be  $P(t) = P_0(b)^{kt}$ , where  $P(t)$  is the population at some time  $t$ ,  $P_0$  is the initial population, and  $k$  is the replication rate. Be sure you convert the hours into minutes!

5. A town has an initial population of 50000. The town's population for the next 10 years is provided below. Which type of function would be most appropriate to model the town's population?

Year	1	2	3	4	5	6	7	8	9
Pop	50026	50056	50094	50124	50146	50176	50214	50244	50266

The solution is Non-Linear Power, which is option A.

- A. Non-Linear Power

This suggests a growth faster than constant but slower than exponential.

- B. Linear

This suggests a constant growth. You would be able to add or subtract the same amount year-to-year if this is the correct answer.

- C. Logarithmic

This suggests the slowest of growths that we know.

- D. Exponential

This suggests the fastest of growths that we know.

- E. None of the above

Please contact the coordinator to discuss why you believe none of the options model the population.

**General Comment:** We are trying to compare the growth rate of the population. Growth rates can be characterized from slowest to fastest as: logarithmic, indirect, linear, direct, exponential. The best way to approach this is to first compare it to linear (is it linear, faster than linear, or slower than linear)? If faster, is it as fast as exponential? If slower, is it as slow as logarithmic?

---

6. Using the scenario below, model the situation using an exponential function and a base of  $\frac{1}{2}$ . Then, solve for the half-life of the element, rounding to the nearest day.

*The half-life of an element is the amount of time it takes for the element to decay to half of its initial starting amount. There is initially 630 grams of element X and after 2 years there is 126 grams remaining.*

The solution is About 0 days, which is option D.

- A. About 730 days

This uses the correct model but solves for the exponential constant incorrectly.

- B. About 365 days

This uses the correct model but a base of  $e$  rather than  $\frac{1}{2}$ .

- C. About 1 day

This models half-life as a linear function.

- D. About 0 days

\* This is the correct option.

- E. None of the above

Please contact the coordinator if you believe all the options above are incorrect.

**General Comment:** The model should be  $A(t) = A_0(\frac{1}{2})^{kt}$ , where  $A(t)$  is the amount after  $t$  years,  $A_0$  is the initial amount, and  $k$  is decay constant. To find the half-life, you need to solve for  $k$  by using the amount after  $x$  years, then solve for the time  $t$  when  $A = \frac{A_0}{2}$ . Your answer would be in years, so convert to days.

---

7. The temperature of an object,  $T$ , in a different surrounding temperature  $T_s$  will behave according to the formula  $T(t) = Ae^{kt} + T_s$ , where  $t$  is minutes,  $A$  is a constant, and  $k$  is a constant. Use this formula and the situation below to construct a model that describes the uranium's temperature,  $T$ , based on the amount of time  $t$  (in minutes) that have passed. Choose the correct constant  $k$  from the options below.

*Uranium is taken out of the reactor with a temperature of  $100^\circ\text{C}$  and is placed into a  $12^\circ\text{C}$  bath to cool. After 21 minutes, the uranium has cooled to  $30^\circ\text{C}$ .*

The solution is None of the above, which is option E.

- A.  $k = -0.08166$

This uses  $A$  as the initial temperature and solves for  $k$  correctly.

- B.  $k = -0.02513$

This uses  $A$  correctly and solves for  $k$  incorrectly.

C.  $k = -0.03950$

This uses  $A$  as the bath temperature and solves for  $k$  incorrectly.

D.  $k = -0.02456$

This uses  $A$  as the initial temperature and solves for  $k$  incorrectly.


E. None of the above

\* This is the correct answer as  $k = -0.07557$ .

**General Comment:** The initial temperature is when  $t = 0$ . Unlike power models, that means  $A$  is not the initial temperature!

---

8. Determine the appropriate model for the graph of points below.



The solution is None of the above, which is option E.

A. Logarithmic model

For this to be the correct option, we want a rapid change early, then an extremely slow change later.

B. Linear model

For this to be the correct option, we need to see a mostly straight line of points.

C. Exponential model

For this to be the correct option, we want an extremely slow change early, then a rapid change later.

D. Non-linear Power model

For this to be the correct option, we need to see a polynomial or rational shape.

E. None of the above

For this to be the correct option, we want to see no pattern in the points.

**General Comment:** This question is testing if you can associate the models with their graphical representation. If you are having trouble, go back to the corresponding Core module to learn about the specific function you are having trouble recognizing.

---

9. Using the scenario below, model the situation using an exponential function and a base of  $\frac{1}{2}$ . Then, solve for the half-life of the element, rounding to the nearest day.

*The half-life of an element is the amount of time it takes for the element to decay to half of its initial starting amount. There is initially 779 grams of element X and after 2 years there is 194 grams remaining.*

The solution is About 0 days, which is option C.

- A. About 1 day

This models half-life as a linear function.

- B. About 730 days

This uses the correct model but solves for the exponential constant incorrectly.

- C. About 0 days

\* This is the correct option.

- D. About 365 days

This uses the correct model but a base of  $e$  rather than  $\frac{1}{2}$ .

- E. None of the above

Please contact the coordinator if you believe all the options above are incorrect.

**General Comment:** The model should be  $A(t) = A_0(\frac{1}{2})^{kt}$ , where  $A(t)$  is the amount after  $t$  years,  $A_0$  is the initial amount, and  $k$  is decay constant. To find the half-life, you need to solve for  $k$  by using the amount after  $x$  years, then solve for the time  $t$  when  $A = \frac{A_0}{2}$ . Your answer would be in years, so convert to days.

---

10. Using the scenario below, model the population of bacteria  $\alpha$  in terms of the number of minutes,  $t$  that pass. Then, choose the correct approximate (*rounded to the nearest minute*) replication rate of bacteria- $\alpha$ .

*A newly discovered bacteria,  $\alpha$ , is being examined in a lab. The lab started with a petri dish of 2 bacteria- $\alpha$ . After 2 hours, the petri dish has 149 bacteria- $\alpha$ . Based on similar bacteria, the lab believes bacteria- $\alpha$  triples after some undetermined number of minutes.*

The solution is None of the above, which is option E.

- A. About 115 minutes

This uses the wrong base and solves for the constant correctly but converted incorrectly.

- B. About 19 minutes

This uses the wrong base.

- C. About 199 minutes

This uses the wrong base, does not solve for the constant correctly, AND converted incorrectly.

- D. About 33 minutes

This uses the wrong base and does not solve for the constant correctly.

- E. None of the above

\* This is the correct option as all other options used the wrong base in their model.

**General Comment:** Your model should be  $P(t) = P_0(b)^{kt}$ , where  $P(t)$  is the population at some time  $t$ ,  $P_0$  is the initial population, and  $k$  is the replication rate. Be sure you convert the hours into minutes!

---