

1. Choose the model type that would best describe the scenario below.

Social distancing is a common tactic to counter potential epidemics. This is due to the exponential increase in number of people infected as the density of people living in an area increases.

- A. Indirect variation
- B. Direct variation
- C. Joint variation
- D. None of the above

-
2. For the scenario below, find the variation constant k of the model (if possible).

In an alternative galaxy, the square of the time, T (Earth years), required for a planet to orbit Sun χ decreases as the square of the distance, d (AUs), that the planet is from Sun χ decreases. For example, when Ea's average distance from Sun χ is 8, it takes 99 Earth days to complete an orbit.

- A. $k = 4.028$
- B. $k = 3.518$
- C. $k = 153.141$
- D. $k = 627264.000$
- E. Unable to compute the constant based on the information given.

-
3. For the scenario below, model the rate of vibration (cm/s) of the string in terms of the length of the string. Then determine the variation constant k of the model (if possible). The constant should be in terms of cm and s.

The rate of vibration of a string under constant tension varies based on the type of string and the length of the string. The rate of vibration of string ω increases as the cube length of the string decreases. For example, when string ω is 2 mm long, the rate of vibration is 22 cm/s.

- A. $k = 2.75$
- B. $k = 0.18$
- C. $k = 176.00$
- D. $k = 2750.00$
- E. None of the above.

4. For the scenario below, model the rate of vibration (cm/s) of the string in terms of the length of the string. Then determine the variation constant k of the model (if possible). The constant should be in terms of cm and s.

The rate of vibration of a string under constant tension varies based on the type of string and the length of the string. The rate of vibration of string ω increases as the quartic length of the string decreases. For example, when string ω is 4 mm long, the rate of vibration is 20 cm/s.

- A. $k = 781.25$
- B. $k = 0.51$
- C. $k = 0.08$
- D. $k = 5120.00$
- E. None of the above.

5. For the scenario below, find the variation constant k of the model (if possible).

In an alternative galaxy, the cube of the time, T (Earth years), required for a planet to orbit Sun χ increases as the cube of the distance, d (AUs), that the planet is from Sun χ increases. For example, when Ea's average distance from Sun χ is 8, it takes 68 Earth days to complete an orbit.

- A. $k = 614.125$
- B. $k = 160989184.000$

- C. $k = 2.041$
- D. $k = 4.028$
- E. Unable to compute the constant based on the information given.

6. For the scenario below, use the model for the volume of a cylinder as $V = \pi r^2 h$ to find the coefficient for the model of the new volume $V_{\text{new}} = k r^2 h$.

Pepsi wants to increase the volume of soda in their cans. They've decided to decrease the radius by 10 percent and increase the height by 14 percent. They want to model the new volume based on the radius and height of the original cans.

- A. $k = 0.00440$
- B. $k = 0.00140$
- C. $k = 2.90095$
- D. $k = 0.92340$
- E. None of the above.

7. Choose the model type that would best describe the scenario below.

In economics, there are two common equations to model interest earned. The compound interest formula is $A = P(1 + \frac{r}{n})^{nt}$, where A is the amount of money you end up with, P is your starting money, r is the interest rate, n is the number of times compounded in a year, and t is the total number of years. For example, if you were a parent and wanted to save \$10,000 in 3 years-time at 3.5% interest compounded monthly, you would need to invest about \$9,000.

- A. Joint variation
- B. Direct variation
- C. Indirect variation
- D. None of the above

8. A town has an initial population of 60000. The town's population for the next 9 years is provided below. Which type of function would be most appropriate to model the town's population?

| Year | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Pop | 60000 | 59979 | 59967 | 59958 | 59951 | 59946 | 59941 | 59937 | 59934 |

- A. Logarithmic
- B. Exponential
- C. Non-Linear Power
- D. Linear
- E. None of the above

9. For the scenario below, use the model for the volume of a cylinder as $V = \pi r^2 h$ to find the coefficient for the model of the new volume $V_{\text{new}} = k r^2 h$.

Pepsi wants to increase the volume of soda in their cans. They've decided to decrease the radius by 13 percent and decrease the height by 17 percent. They want to model the new volume based on the radius and height of the original cans.

- A. $k = 1.97363$
- B. $k = 0.62823$
- C. $k = 0.00287$
- D. $k = 0.00903$
- E. None of the above.

10. A town has an initial population of 60000. The town's population for the next 9 years is provided below. Which type of function would be most appropriate to model the town's population?

| Year | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
|-------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Pop | 60000 | 59986 | 59978 | 59972 | 59967 | 59964 | 59961 | 59958 | 59956 |

- A. Exponential
- B. Linear
- C. Non-Linear Power
- D. Logarithmic
- E. None of the above