

1. Add the following functions, then choose the domain of the resulting function from the list below.

$$f(x) = \sqrt{-3x - 16} \text{ and } g(x) = 3x^3 + 9x$$

- A. The domain is all Real numbers except  $x = a$ , where  $a \in [4.2, 12.2]$
  - B. The domain is all Real numbers less than or equal to  $x = a$ , where  $a \in [-7.33, 0.67]$
  - C. The domain is all Real numbers greater than or equal to  $x = a$ , where  $a \in [5.33, 11.33]$
  - D. The domain is all Real numbers except  $x = a$  and  $x = b$ , where  $a \in [5.67, 8.67]$  and  $b \in [4.6, 12.6]$
  - E. The domain is all Real numbers.
- 

2. Find the inverse of the function below. Then, evaluate the inverse at  $x = 7$  and choose the interval that  $f^{-1}(7)$  belongs to.

$$f(x) = \ln(x - 5) - 2$$

- A.  $f^{-1}(7) \in [153.41, 157.41]$
  - B.  $f^{-1}(7) \in [8107.08, 8112.08]$
  - C.  $f^{-1}(7) \in [2.39, 7.39]$
  - D.  $f^{-1}(7) \in [8096.08, 8099.08]$
  - E.  $f^{-1}(7) \in [162748.79, 162755.79]$
- 

3. Determine whether the function below is 1-1.

$$f(x) = 25x^2 + 220x + 484$$

- A. Yes, the function is 1-1.
- B. No, because the range of the function is not  $(-\infty, \infty)$ .
- C. No, because there is an  $x$ -value that goes to 2 different  $y$ -values.

- D. No, because there is a  $y$ -value that goes to 2 different  $x$ -values.
- E. No, because the domain of the function is not  $(-\infty, \infty)$ .

- 
4. Find the inverse of the function below. Then, evaluate the inverse at  $x = 10$  and choose the interval that  $f^{-1}(10)$  belongs to.

$$f(x) = e^{x+5} - 3$$

- A.  $f^{-1}(10) \in [-1.1, -0.84]$
- B.  $f^{-1}(10) \in [7.32, 8.32]$
- C.  $f^{-1}(10) \in [-0.32, 0.71]$
- D.  $f^{-1}(10) \in [-2.55, -1.81]$
- E.  $f^{-1}(10) \in [-2.19, -1.3]$

- 
5. Determine whether the function below is 1-1.

$$f(x) = (5x + 17)^3$$

- A. No, because the range of the function is not  $(-\infty, \infty)$ .
- B. Yes, the function is 1-1.
- C. No, because there is a  $y$ -value that goes to 2 different  $x$ -values.
- D. No, because there is an  $x$ -value that goes to 2 different  $y$ -values.
- E. No, because the domain of the function is not  $(-\infty, \infty)$ .

- 
6. Choose the interval below that  $f$  composed with  $g$  at  $x = -1$  is in.

$$f(x) = -4x^3 - 2x^2 + 4x \text{ and } g(x) = -3x^3 - 4x^2 - x - 3$$

- A.  $(f \circ g)(-1) \in [10, 23]$
- B.  $(f \circ g)(-1) \in [83, 91]$
- C.  $(f \circ g)(-1) \in [75, 84]$

- D.  $(f \circ g)(-1) \in [3, 15]$   
E. It is not possible to compose the two functions.
- 

7. Choose the interval below that  $f$  composed with  $g$  at  $x = 1$  is in.

$$f(x) = 2x^3 + 2x^2 - 3x \text{ and } g(x) = -4x^3 + x^2 + 2x - 3$$

- A.  $(f \circ g)(1) \in [-9, -3]$   
B.  $(f \circ g)(1) \in [1, 4]$   
C.  $(f \circ g)(1) \in [-84, -83]$   
D.  $(f \circ g)(1) \in [-76, -73]$   
E. It is not possible to compose the two functions.
- 

8. Find the inverse of the function below (if it exists). Then, evaluate the inverse at  $x = -10$  and choose the interval that  $f^{-1}(-10)$  belongs to.

$$f(x) = \sqrt[3]{4x - 3}$$

- A.  $f^{-1}(-10) \in [-250, -247]$   
B.  $f^{-1}(-10) \in [246.4, 250.6]$   
C.  $f^{-1}(-10) \in [249.9, 253]$   
D.  $f^{-1}(-10) \in [-250.9, -249.9]$   
E. The function is not invertible for all Real numbers.
- 

9. Find the inverse of the function below (if it exists). Then, evaluate the inverse at  $x = -15$  and choose the interval that  $f^{-1}(-15)$  belongs to.

$$f(x) = 5x^2 - 2$$

- A.  $f^{-1}(-15) \in [5.33, 5.75]$   
B.  $f^{-1}(-15) \in [1.81, 2.05]$

- C.  $f^{-1}(-15) \in [1.48, 1.65]$
  - D.  $f^{-1}(-15) \in [4.33, 4.91]$
  - E. The function is not invertible for all Real numbers.
- 

10. Subtract the following functions, then choose the domain of the resulting function from the list below.

$$f(x) = 9x^2 + 8x + 6 \text{ and } g(x) = 5x^3 + 8x^2 + 5x + 6$$

- A. The domain is all Real numbers less than or equal to  $x = a$ , where  $a \in [0.5, 5.5]$
  - B. The domain is all Real numbers except  $x = a$ , where  $a \in [6.2, 7.2]$
  - C. The domain is all Real numbers greater than or equal to  $x = a$ , where  $a \in [2.67, 9.67]$
  - D. The domain is all Real numbers except  $x = a$  and  $x = b$ , where  $a \in [-7.33, 1.67]$  and  $b \in [-5.25, -2.25]$
  - E. The domain is all Real numbers.
-