

Monte Carlo Simulation of Linear Accelerator for Dosimetry Analysis

Under the guidance of Dr.Jayaraj P B and Dr.Pournami P N

Darshana Suresh B160373CS

Shalini Nath B160613CS

Saai Lakshmi B160584CS

Department of Computer Science and Engineering
NIT Calicut

05 March 2020

Introduction - Problem Statement

Monte Carlo simulation of a linear accelerator for treatment planning of cancer:

- Simulation of radiation beam production in LINAC.
- Simulation of beam transport from LINAC head to phantom.
- Dosimetry analysis of radiation.

- Installed GATE (Geant4 Application for Tomographic Emission) and compared the ease of usage with Geant4 for simulation purposes.
- Obtained sample specifications of an Elekta Synergy LINAC from "Grid Monte Carlo Simulation of a Medical Linear Accelerator" paper and updated the simulation of LINAC using GATE v8.2 [1]
- Researched on the various types of interactions of photons with matter and studied its relevance in the radiotherapy treatment process.


Photon Interactions with Matter

- Radiotherapy treatment uses high energy photons
- If the source of photon is from the nucleus of a radioactive atom, they are called gamma rays
- If it is created electronically, for eg. using LINAC, they form X-rays
- The maximum energy of the X-ray spectrum produced can be controlled, unlike the case of gamma ray
- X-ray, on getting produced from the linac, goes through several interactions with matter.

Photon Interactions with Matter

- The energy of photons is imparted to matter in a two stage process.
- In the first stage, the photon radiation transfers energy to the secondary charged particles (electrons) through various photon interactions
- In the second stage, the charged particle transfers energy to the medium through atomic excitations and ionizations.
- The measure of energy transferred from radiation to matter is called **KERMA (Kinetic Energy Released per unit MAss)**.

Photoelectric Effect




The incoming photon transfers its energy to an atomic electron, ejecting it out from the atom and ionizing its neighbouring molecules

Photoelectric Effect

- Energy conservation is given by $E_e = h\nu - E_b$
- This can only take place with an atom as a whole and not with free electrons
- The most probable origin of the photoelectron is the most tightly bound shell (K-shell) of the atom
- The PE interaction is most probable for **low energy and high atomic number and high density materials**
- This effect is predominant in tissues in the energy range **10-25 keV**. [4]

Compton Effect



A high energy photon collides with a free electron and gets deflected with respect to its original direction through an angle θ .

Compton Effect


- The transfer of energy is given by the formula

$$h\nu_0 = \frac{h\nu}{(1 + (h\nu/m_0c^2)(1 - \cos\theta))}$$

- The probability of this effect is independent of the matter's **atomic number** but is inversely proportional to the **energy of the incoming photon**
- The energy range for the effect is **25keV - 25MeV** [4]
- It is the most dominant interaction mechanism in tissue.

Coherent Scattering


There are two types of elastic scattering : Rayleigh scattering and Thomson scattering. [3]



- Energy of the X-ray or gamma photon is low (10 keV).
- The only significant change is a change of direction of the photon.
- It varies with the atomic number of the absorber (Z) and incident photon energy (E) by Z/E^2 .


Pair Production

- It which occurs when a photon is in close proximity to the nucleus of an atom with an energy of **at least 1.022 MeV**.^[4]




- Results in formation of a pair of particles, an electron and a positively charged positron.
- Related to the atomic number of a material through Z^2 .

Triplet Production



- Triplet Production is a special case of pair production which occurs in the vicinity of an orbital electron instead of a nucleus.
- Here, a positron (anti-electron) and an electron are produced spontaneously as a photon interacts with a strong electric field with an electron.[3]

Photodisintegration



- Photodisintegration occurs when a photon is absorbed by the nucleus of an atom.
- The nucleus becomes excited and becomes radioactive. To become stable, the nucleus emits any one among negatrons, protons, neutrons, alpha particles, clusters of fragments or gamma rays.
- The threshold for this effect is **over 10 MeV** for most nuclei.[3]

GATE- Geant4 Application for Tomographic Emission

- GATE is an open-source software developed by the international OpenGATE collaboration for numerical simulations in **medical imaging and radiotherapy**. [2]
- Several of the previous simulations of LINAC have been carried out using GATE.
- This software makes use of macros entirely to define geometry, beam transportation, and further simulations, therefore increasing the readability of the code.
- We have installed GATE and compared the ease of usage with Geant4 for simulation purposes.

GATE vs Geant4

- GATE uses only macro files for geometry, scoring, physics and beam, unlike in Geant4 which has separate header files, source files etc. written in C++.
- GATE is flexible and can design different kinds of detectors.
- Geometry designing is easier in GATE.
- GATE contains its own material database. New materials/alloys can be added to the db.

GATE - Implementing Photon Interactions

- Can implement Photoelectric effect, Compton effect, Pair production, Rayleigh scattering.
- Three models are available for electromagnetic processes: Standard, Livermore, and Penelope.
- The Penelope models have been specifically developed for Monte Carlo simulation. They are effective between 250 eV and 1 GeV.

Visuals of the simulation




Figure: GATE Linac Simulation

Visuals of the simulation




Figure: GATE Linac Simulation with added MLCs

Work Plan

Work to be done

- Study several research papers on Monte Carlo simulation of linacs and work on a review paper.
- Study the role of Monte Carlo methods in the simulations to be done.
- Simulation of radiation beam transportation from the LINAC treatment head to the phantom.

Conclusion

- We have installed GATE and compared the ease of usage with Geant4 for simulation purposes.
- Thorough understanding of photon interactions with matter.

References

-  [1] Didi et. al. (2018). Grid Monte Carlo Simulation of a Medical Linear Accelerator. 3. 40-43. 10.24018/ejers.2018.3.12.1001.
-  [2] GATE homepage, <http://www.opengatecollaboration.org/>. Last accessed 03 March 2020
-  [3]<http://ozradonc.wikidot.com/photon-interactions>
-  [4]<https://pdfs.semanticscholar.org/f6c6/e44bd495775ba1433c6c42ff68f71b7f823f.pdf>

Thank You