Supplementary Methods: Constrained Bivariate Gaussian Mixture Modeling

Darwin Brunel Ngoune Domo

1. Objective and Input Data

To determine whether the mRNA–protein response regimes identified by Differential Pattern Analysis (DiPa) reflect distinct biological subpopulations rather than arbitrary slices of a continuous cloud, we fitted constrained bivariate Gaussian Mixture Models (GMM) to the empirical $\log_2$ fold changes independently of the ± 0.5 cutoffs.

The input is a matrix $X \in \mathbb{R}^{N\times 2}$ ($N = 943$ core matched pairs), where for each gene i:

$$X_i = \begin{pmatrix} x_i \\ y_i \end{pmatrix} = \begin{pmatrix} \log_2 \mathrm{FC}_{\mathrm{RNA},i} \\ \log_2 \mathrm{FC}_{\mathrm{Prot},i} \end{pmatrix}. \tag{1}$$

2. Model Specification and Biophysical Constraints

The probability density of observation X_i under a mixture of $K \in \{1, 2, 3\}$ bivariate Gaussians is:

$$p(X_i \mid \theta) = \sum_{k=1}^{K} \pi_k \mathcal{N}_2 \left(X_i \mid \mu_k, \Sigma_k\right), \quad \text{with } \pi_k \geq 0, \quad \sum_{k=1}^{K} \pi_k = 1, \tag{2}$$

where $\mu_k = (\mu_{xk}, \mu_{yk})^\top$ are component centroids and $\Sigma_k = \begin{pmatrix} \sigma_{xk}^2 & \rho_k \sigma_{xk}\sigma_{yk} \\ \rho_k \sigma_{xk}\sigma_{yk} & \sigma_{yk}^2 \end{pmatrix}$ are covariance matrices.

To prevent unconstrained components from fitting arbitrary noise, three biophysical constraints were enforced:

1. **Component 1 (Diagonal / Co-regulation):** Represents canonical stoichiometric coupling. The correlation is bounded within a biologically realistic coupling window:
$$\rho_1 \in [0.70,\, 0.90]. \tag{3}$$
2. **Component 2 (Vertical / Protein-dominant decoupling):** Represents protein changes occurring without significant transcriptomic shifts. Coordinate correlation is set to zero, and variance is vertically elongated by factor c:
$$\rho_2 = 0, \qquad \sigma_{y2}^2 \geq c \cdot \sigma_{x2}^2. \tag{4}$$
3. **Component 3 (Horizontal / Post-transcriptional buffering, for $K = 3$):** Represents large mRNA changes buffered at the protein level. Correlation is set to zero, and variance is horizontally elongated:
$$\rho_3 = 0, \qquad \sigma_{x3}^2 \geq c \cdot \sigma_{y3}^2. \tag{5}$$

The elongation factor was systematically tested across $c \in \{1.3, 1.5, 2.0, 2.5, 4.0\}$ (aspect ratios $\sqrt{c} \in [1.14, 2.00]$).

3. Expectation-Maximization (EM) Algorithm

Parameters were estimated iteratively via the Expectation-Maximization (EM) algorithm [1, 2] until the log-likelihood converged ($|\ell^{(t+1)} - \ell^{(t)}| < 10^{-6}$):

- **Initialization:** Posterior weights $\gamma_{ik}^{(0)} \in [0, 1]$ were initialized deterministically based on coarse regions (vertical candidates: $|x_i| < 0.55, |y_i| > 0.40$; horizontal candidates: $|x_i| > 0.50, |y_i| < 0.50$; diagonal otherwise), normalized so $\sum_k \gamma_{ik}^{(0)} = 1$.

- **M-Step (Maximization):** Given weights γ_{ik}, component sample sizes $N_k = \sum_i \gamma_{ik}$ and mixing weights $\pi_k = N_k/N$ were computed. Centroids and sample dispersions were updated:

$$\mu_k = \frac{1}{N_k}\sum_{i=1}^{N}\gamma_{ik}X_i, \quad v_{xk} = \frac{1}{N_k}\sum_{i=1}^{N}\gamma_{ik}(x_i-\mu_{xk})^2, \quad v_{yk} = \frac{1}{N_k}\sum_{i=1}^{N}\gamma_{ik}(y_i-\mu_{yk})^2, \tag{6}$$

  and $c_{xy,1} = \frac{1}{N_1}\sum_{i=1}^{N}\gamma_{i1}(x_i-\mu_{x1})(y_i-\mu_{y1})$. Constraints were then directly applied to define the final dispersion parameters $\hat{\sigma}^2$ and matrices Σ_k:
  - For $k=1$: $\hat{\sigma}^2_{x1} = v_{x1}$, $\hat{\sigma}^2_{y1} = v_{y1}$, $\hat{\rho}_1 = \min(0.90, \max(0.70, c_{xy,1}/\sqrt{v_{x1}v_{y1}}))$, and $\Sigma_1 = \begin{pmatrix} \hat{\sigma}^2_{x1} & \hat{\rho}_1\hat{\sigma}_{x1}\hat{\sigma}_{y1} \\ \hat{\rho}_1\hat{\sigma}_{x1}\hat{\sigma}_{y1} & \hat{\sigma}^2_{y1} \end{pmatrix}$.
  - For $k=2$: $\hat{\sigma}^2_{x2} = v_{x2}$, $\hat{\sigma}^2_{y2} = \max(v_{y2},\, c\cdot v_{x2})$, and $\Sigma_2 = \mathrm{diag}(\hat{\sigma}^2_{x2},\, \hat{\sigma}^2_{y2})$.
  - For $k=3$: $\hat{\sigma}^2_{x3} = \max(v_{x3},\, c\cdot v_{y3})$, $\hat{\sigma}^2_{y3} = v_{y3}$, and $\Sigma_3 = \mathrm{diag}(\hat{\sigma}^2_{x3},\, \hat{\sigma}^2_{y3})$.
- **E-Step (Expectation):** Using the updated parameters, posterior responsibilities were recalculated:

$$\gamma_{ik}^{(t+1)} = \frac{\pi_k\,\mathcal{N}_2\left(X_i \mid \mu_k,\, \Sigma_k\right)}{\sum_{j=1}^{K}\pi_j\,\mathcal{N}_2\left(X_i \mid \mu_j,\, \Sigma_j\right)}, \quad \ell^{(t+1)} = \sum_{i=1}^{N}\ln\left(\sum_{k=1}^{K}\pi_k\,\mathcal{N}_2\left(X_i \mid \mu_k,\, \Sigma_k\right)\right). \tag{7}$$

4. Model Selection (BIC)

Model selection across $K \in \{1,2,3\}$ and $c \in \{1.3, 1.5, 2.0, 2.5, 4.0\}$ was evaluated using the Bayesian Information Criterion (BIC) [3]:

$$\mathrm{BIC} = -2\hat{\ell} + P\ln(N), \tag{8}$$

where $\hat{\ell}$ is the maximized log-likelihood, $N = 943$ ($\ln(943) \approx 6.849$), and P is the number of free parameters:

- $K=1$: 2 means + 2 variances + 1 covariance $\implies P = 5$.
- $K=2$ (constrained): 1 weight (π_1) + 4 means + 3 dispersion (Comp 1) + 2 variances (Comp 2) $\implies P = 10$.
- $K=3$ (constrained): 2 weights + 6 means + 3 (Comp 1) + 2 (Comp 2) + 2 (Comp 3) $\implies P = 15$.

A difference $\Delta\mathrm{BIC} < -10$ provides decisive evidence in favor of the lower-BIC model on the Kass–Raftery scale [4]. BIC was chosen over classical likelihood-ratio tests because testing component complexity (K vs. $K+1$) places the null hypothesis ($\pi_{K+1} = 0$) on the boundary of the probability simplex, violating Wilks' regularity conditions [2].

5. Confidence Ellipses

Concentration ellipses for component k at confidence levels $\alpha \in \{0.50, 0.90, 0.95\}$ were plotted using the spectral decomposition $\hat{\Sigma}_k = V\Lambda V^\top$:

$$E_k(\alpha) = \hat{\mu}_k + \sqrt{F^{-1}_{\chi^2_2}(\alpha)}\cdot V\Lambda^{1/2}\begin{pmatrix}\cos\phi \\ \sin\phi\end{pmatrix}, \quad \phi \in [0, 2\pi), \tag{9}$$

with radii $\sqrt{F^{-1}_{\chi^2_2}(0.50)} \approx 1.177$, $\sqrt{F^{-1}_{\chi^2_2}(0.90)} \approx 2.146$, and $\sqrt{F^{-1}_{\chi^2_2}(0.95)} \approx 2.448$.

The complete analysis pipeline is implemented in base R and `data.table` (see script `Generate_Supplementary_Figure_6_Master_Plate.R`).

References

[1] A. P. Dempster, N. M. Laird, and D. B. Rubin. Maximum likelihood from incomplete data via the EM algorithm. *J. R. Stat. Soc. Series B*, 39(1):1–22, 1977.

[2] G. J. McLachlan and D. Peel. *Finite Mixture Models*. John Wiley & Sons, New York, 2000.

[3] G. Schwarz. Estimating the dimension of a model. *Ann. Stat.*, 6(2):461–464, 1978.

[4] R. E. Kass and A. E. Raftery. Bayes factors. *J. Am. Stat. Assoc.*, 90(430):773–795, 1995.