

Human alteration of natural light cycles: causes and ecological consequences

Author(s): Kevin J. Gaston, James P. Duffy, Sian Gaston, Jonathan Bennie and Thomas W. Davies

Source: *Oecologia*, Vol. 176, No. 4 (December 2014), pp. 917-931

Published by: Springer in cooperation with International Association for Ecology

Stable URL: <https://www.jstor.org/stable/43671445>

Accessed: 05-09-2019 18:06 UTC

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <https://about.jstor.org/terms>



JSTOR

Springer, International Association for Ecology are collaborating with JSTOR to digitize, preserve and extend access to *Oecologia*

Human alteration of natural light cycles: causes and ecological consequences

Kevin J. Gaston · James P. Duffy · Sian Gaston ·
Jonathan Bennie · Thomas W. Davies

Received: 9 June 2014 / Accepted: 4 September 2014 / Published online: 20 September 2014
© The Author(s) 2014. This article is published with open access at Springerlink.com

Abstract Artificial light at night is profoundly altering natural light cycles, particularly as perceived by many organisms, over extensive areas of the globe. This alteration comprises the introduction of light at night at places and times at which it has not previously occurred, and with different spectral signatures. Given the long geological periods for which light cycles have previously been consistent, this constitutes a novel environmental pressure, and one for which there is evidence for biological effects that span from molecular to community level. Here we provide a synthesis of understanding of the form and extent of this alteration, some of the key consequences for terrestrial and aquatic ecosystems, interactions and synergies with other anthropogenic pressures on the environment, major uncertainties, and future prospects and management options. This constitutes a compelling example of the need for a thoroughly interdisciplinary approach to understanding and managing the impact of one particular anthropogenic pressure. The former requires insights that span molecular biology to ecosystem ecology, and the latter contributions of biologists, policy makers and engineers.

Keywords Day · Diurnal · Night · Nocturnal · Skyglow

Introduction

Ecological systems are organized foremost by light, and particularly by daily and seasonal cycles of light and dark

(Bradshaw and Holzapfel 2010; Kronfeld-Schor et al. 2013). Humans are profoundly altering these cycles as detected and/or perceived by many organisms. This is occurring by the introduction of artificial light at night (ALAN) in the environment, predominantly from electric lighting sources associated with human settlement, transport networks and industry, the impact of which extends across much of the globe (Cinzano et al. 2001). In turn this is influencing biological systems from the molecule to the ecosystem, including impacts on gene expression, physiology and behaviour of organisms, abundance and distribution of species, ecological interactions, and the composition of communities (e.g. recent examples include Bird et al. 2004; Davies et al. 2012; Dwyer et al. 2012; Dominioni et al. 2013a; Le Tallec et al. 2013; Mazarin et al. 2013; Picchi et al. 2013). This then almost inevitably affects the function and process of ecosystems, and thus other fundamental ecological cycles. This paper reviews the form and extent of the human alteration of natural light cycles, key consequences for terrestrial and aquatic ecosystems, interactions and synergies with other anthropogenic environmental pressures, major uncertainties, and future prospects and management options. Several of these topics have not previously been well developed. As a synthesis, this is an illustrative rather than an exhaustive compilation of relevant studies, which are numerous but highly scattered within the literature (see Rich and Longcore 2006; Hölker et al. 2010; Perkin et al. 2011; Gaston et al. 2012, 2013; Bogard 2013; Gaston and Bennie 2014).

Communicated by Nina Farwig.

K. J. Gaston (✉) · J. P. Duffy · S. Gaston · J. Bennie ·
T. W. Davies
Environment and Sustainability Institute, University of Exeter,
Penryn, Cornwall TR10 9FE, UK
e-mail: k.j.gaston@exeter.ac.uk

Human alteration of natural light cycles

Natural light cycles are, of course, driven entirely from an extraplanetary source, the sun. The primary cycles take

three distinct forms (others with longer periods are not relevant here). First, rotation of the Earth partitions time into a regular cycle of day and night, such that the intensity of incident light at a site typically varies by ca. eight orders of magnitude (Fig. 1). Second, the Earth's orbital motion and tilt of its axis cause marked seasonal variation in the distribution of the approximately 4,400 h of each year spent under conditions of relative darkness (including twilight, moonlight and starlight). Third, the nighttime light environment is subject to systematic variation as a consequence of the orbit of the moon around the Earth, and the level and pattern of reflected sunlight (moonlight) during the night. The effect of all three of these cycles on the light received at ground level (Fig. 2a–c) is modified to some degree by local topography, habitat and weather (especially cloud cover; Fig. 2d), and by other intermittent natural sources of light (e.g. lightning strikes and fires; Fig. 2e, f). However, the amplitude and frequency of these cycles prevail.

These daily, annual and lunar geophysical cycles have also remained rather invariant over long periods of time. For example, the Earth's period of revolution around the

Sun has been effectively constant. Its period of rotation around its axis, presently 24 h, has experienced deceleration, such that day length has increased through geological time, from ~21 h at the beginning of the Cambrian (Wells 1963). However, this amounts to a rate of increase only of ~0.002 s per century (Wahr 1988). This background means that ALAN is rather unusual amongst anthropogenic environmental pressures. Most others (e.g. changes in CO₂, precipitation, temperature) have historical analogues, having previously altered naturally over geological or evolutionary time in similar ways to those presently experienced, albeit often at different rates. The most fundamental human-caused change to light cycles has two key characteristics, changes in the spatial and temporal occurrence of light and changes in its spectrum.

Changing occurrence of light

ALAN has introduced light in places, times and at intensities at which it does not naturally occur (Figs. 1, 2). The extent of these changes remains to be fully evaluated. Data





Fig. 1 Change in illumination at the Earth's surface with solar (positive) and lunar altitude (negative) above the horizon; typical illumination levels of artificial light at night (ALAN); and levels at which nighttime lighting has been observed to have biological effects [arrows; Sharma et al. (1997) (a), Zubidat et al. (2007) (b), Johnson (1979) (c), Stone et al. (2009) (d), Kuijper et al. (2008) (e), Riley et al. (2012) (f), Bedrosian et al. (2011) (g), Miller (2006) and Kramer and Birney (2001) (h), Falkenberg and Clarke (1998) and

Clarke et al. (1996) (h), Santos et al. (2010) (j), Dauchy et al. (1997) and Cos et al. (2006) (k), Bachleitner et al. (2007) (l), Evans et al. (2007a) (m), Larsen and Pedersen (1982) and Dice (1945) (n), Dice (1945) (o–q); studies of levels at which nighttime lighting has biological effects are from Gaston et al. (2013, Table 3)]. Main figure modified from Beier (2006), with additional data from Kurtze (1974); Rich and Longcore (2006) and Gaston et al. (2013). SS Sunset, CT civil twilight, NT nautical twilight, AT astronomical twilight

Fig. 2a–g Global distributions of natural and artificial light. **a–c** Modelled yearly levels of day-light, moonlight and twilight in total hours, respectively, following equations in Meeus (2008), **d** cloud cover [composite of 12 monthly cloud fraction images for 2012 (Stockli 2013)], **e** mean annual lightning flash rate [flashes $\text{km}^{-2} \text{ year}^{-1}$ for 2012; Lightning Imaging Sensor/Optical Transient Detector gridded lightning climatology data set (NASA 2012)], **f** fire [sum of 12 cloud-corrected fire pixel images for 2012; NASA Land Processes Distributed Active Archive Center (2013)], and **g** artificial nighttime lighting {represented as digital number in 2010 [US Defense Meteorological Satellite Program (DMSP)/Operational Linescan System (OLS) 2012]}



principally arise from satellite imagery and aerial surveys of zenith-directed light emissions on cloud-free nights (e.g. Elvidge et al. 1999; Levin and Duke 2012; Miller et al. 2012; Mazar et al. 2013; Bennie et al. 2014a). Data from a recent satellite image, at a resolution of 810×810 m, indicate occurrence of ALAN in 3.4 % of cells globally (defined as a digital number >5), and 0.2 % of marine cells, but 11.4 % of terrestrial ones (Fig. 3). Where it does occur, this light varies markedly in intensity, even given saturation in the sensors at higher levels of emissions (Fig. 3). Spatial variation in the level of nighttime lighting tends to be well correlated with the level of development, built density, population density and economic activity of an area (Sutton 2003; Amaral et al. 2006; Doll et al. 2006; Chen and Nordhaus 2011; Pun and So 2011; Li et al. 2012; Hale et al. 2013). Recent decades have seen widespread increases in the number of spatially distinct lights and in the lit area, with changes particularly marked in Asia (Small and Elvidge 2011, 2013). These estimates focus on temporally persistent sources of light, and thus underestimate the overall scale of change.

Light emissions detected from satellite imagery and aerial surveys can bear complex relations to those experienced at ground level. Key features of ALAN on the

ground include a marked degree of spatial heterogeneity, with maximum values of intensity in areas of direct illumination, complex patterns of shading on the ground surface due to the number and location of light sources, and large areas affected by lower intensity illumination from reflected and scattered light in the atmosphere. It varies markedly in intensity, with areas such as sports fields and parking lots often lit to illuminance values of several hundred lux or above, ground-level illuminance in the vicinity of street lights around 10–40 lux and reduced to <1 lux several metres away.

ALAN that is emitted or reflected upwards can be scattered by water, dust and gas molecules in the atmosphere, resulting in skyglow. Studies of sky irradiance have been made for individual sites, sets of sites, and cities (Kyba et al. 2011a, b; Biggs et al. 2012; Davies et al. 2013). Skyglow can be detected over a much wider area than direct artificial lighting—extending tens and perhaps hundreds of kilometres from the source—particularly because of the contribution of light that is emitted or reflected upward at relatively shallow angles to the horizontal (Crawford 2000). Local levels tend to be closely associated with prevailing land use, being greater in more highly developed areas and declining away from these (Garstang 1986;




Fig. 3 A raw nighttime stable lights image (2012) from the US DMSP/OLS (in Behrmann equal-area projection at a resolution of 810×810 m). Digital number indicates light intensity with 0 representing darkness and 63 indicating the brightest pixels. Histograms show data from this image **a** without and **b** with 0 digital number values. **c** Grey graphs show the mean digital number for particular columns/rows (latitudes/longitudes)

Crawford 2000; Biggs et al. 2012). It can attain levels of up to 0.2–0.5 lux (Kurtze 1974; Eisenbeis 2006), and under cloudy conditions in urban areas skyglow has been shown to be of an equivalent or greater magnitude than high-elevation summer moonlight (Kyba et al. 2011b). Indeed, cloud cover (which varies markedly; Fig. 2d) increases ALAN, the reverse of what happens during daytime (Kyba et al. 2011b). On clear nights skyglow reduces the visibility of stars and other celestial objects (Kyba et al. 2013).

Modelling techniques enable global sky brightness estimates to be obtained using satellite imagery of nighttime lights (e.g. Cinzano et al. 2001; Cinzano and Elvidge 2004). One such exercise estimated that at the turn of the twenty-first century about two-thirds of the global human population already lived in areas where sky brightness is above the threshold set for polluted status, about one fifth had lost naked-eye visibility of the Milky Way, and for about a tenth sky brightness was such they no longer viewed nighttime skies with the eye adapted to night vision (Cinzano et al. 2001).

Changing spectra of light

Not only does ALAN change the spatial and temporal structure and intensity of natural light cycles, it also occurs with spectra different from those of sunlight, moonlight or starlight (Fig. 4). Some types of artificial lighting are restricted to narrow bandwidths (e.g. low-pressure sodium lighting emits a single narrow peak in the visible spectrum at 589.3 nm). Others emit over a wide range of wavelengths [high-pressure sodium lighting emits a yellow light allowing some colour discrimination in humans; high-intensity discharge lamps emit a whiter light, with significant peaks in blue and ultra-violet wavelengths, and light-emitting diode (LED)-based white street lighting typically emits at all wavelengths between around 400 and 700 nm, with peaks in the blue and green (Elvidge et al. 2010)]. The prevailing sources tend to vary from one region to another, and hence the nature of the resultant ALAN. There is, however, a trend towards the adoption of lighting technologies with a broader spectrum of ‘white’ light. This increases the amount of skyglow visible to people and many other organisms (Van Tichelen et al. 2007, p. 91).

Sources of change

There is a diversity of principal public and private sources of ALAN.

Street lights

Street lighting appears from satellite and aerial imagery to be the dominant terrestrial source of ALAN, albeit not that with the most intense emissions (Kuechly et al. 2012). To some extent this is because street lights are more prone to upward unshielded or reflected light emissions, but it also results from the huge numbers of such lights and the lighting being unconstrained by other factors (e.g. lighting from within buildings is filtered through windows). Good estimates of the actual numbers of street lights appear to be lacking, although one recent figure suggests there are ca. 60 million in EU countries alone (Van Tichelen et al. 2007). However, the global paved road network, much of which is typically lit at night, is estimated at 18,015,713 km (CIA 2011), giving some indication of the potential extent of this source of ALAN.

Buildings

The internal and, particularly, external lighting of buildings contributes substantially to nighttime views of major cities and conurbations, rendering some iconic in this regard (e.g. Paris, Las Vegas, Hong Kong, Shanghai).




Fig. 4 The spectral radiance of **a** daylight, **b** incandescent, **c** low-pressure sodium, **d** light-emitting diode, **e** mercury vapour, and **f** fluorescent lighting. Daylight spectra from pveducation.org and artificial light spectra from http://ngdc.noaa.gov/eog/night_sat/spectra.html

Urban areas are typically defined in terms of the level of coverage by buildings and associated infrastructure. Estimates of urban land cover are highly variable (Gaston 2010), but typical global figures are of the order of 2–3 % of land [excluding permanent ice cover (e.g. Millennium Ecosystem Assessment 2005)]. However, regional coverage may be substantially larger; figures for 165 countries vary from close to zero to 32 % (World Resources Institute 2007).

Road vehicles

Terrestrially, the headlights of road vehicles produce substantial quantities of ALAN. On all but the busiest roads these emissions are temporally highly variable, occur predominantly in the horizontal plane, and are thus underestimated from satellite and aerial imagery. The orientation of these emissions means they may propagate over long distances. They have also progressively increased with major developments in headlight technology (Mainster and Timberlake 2003). Globally, in 2012 there were an estimated 833,342,000 passenger cars and 309,888,000 commercial vehicles (Organisation Internationale des Constructeurs d'Automobiles 2014), although it is unclear what proportion of these are used at night and with what frequency. The

ecological impacts of ALAN from traffic has been little explored (Lyytimäki et al. 2012).

Vessels

In the marine environment, significant ALAN is produced by shipping and offshore infrastructure such as oil and gas platforms. Particular attention has been paid to that generated by fishing fleets [especially those employing banks of lights to attract squid (e.g. Kiyofuji and Saitoh 2004; Elvidge et al. 2001)]. Although these lights are transient, much shipping is aggregated along common routes around coastlines and across oceans (Kareiva et al. 2007), and fishing fleets whilst operating over much larger extents tend disproportionately to focus activities on quite constrained areas (Jennings and Lee 2012).

Effects on terrestrial ecosystems

There have been a number of reviews of the ecological effects of ALAN, focusing principally on evidence of their breadth, in terms of different processes or levels of biological organization (Fig. 5; Longcore and Rich 2004, 2006; Gaston et al. 2013; Gaston and Bennie 2014), taxonomic




Fig. 5a–d The effects of ALAN on animals. **a** Loggerhead turtle hatchlings crawl towards artificial light when it is turned on, and the ocean when it is turned off (Salmon et al. 1995). **b** The effect of high-pressure sodium street lighting on the abundance of invertebrates within trophic groups. Bars represent the total number of individuals in each group collected from pitfall traps under lights (open bars) and between lights (grey bars) (Davies et al. 2012). **c** The influence of light intensity on the suppression of pineal melatonin content after

30 min of exposure. Bars indicate mean pineal melatonin content (for each group $n = 7$). $*p < 0.001$ (Brainard et al. 1984). **d** The effect of artificial night light on paternity gain for adult and yearling blue tits *Cyanistes caeruleus* occupying edge territories; data are point estimates and 95 % confidence intervals from a generalised linear mixed model in which age and territory category are fixed factors and male identity and season are random intercepts; numbers show sample sizes (Kempnaers et al. 2010)

levels (Rich and Longcore 2006) and research domain (Perkin et al. 2011). Here we highlight selected effects that in our opinion are emerging as likely to be of key significance in terrestrial and in aquatic ecosystems, starting with the former.

Individual health

Natural light cycles influence the timings of numerous physiological processes. In many animals melatonin plays a key role in this (Vivien-Roels and Pévet 1993; Arendt 1998). Exposure either to even brief periods of high-intensity ALAN, or to prolonged periods of low intensity, has been shown in the laboratory to be capable of substantially altering patterns of circadian clock gene expression and melatonin production (e.g. Dauchy et al. 1997; Bedrosian et al. 2013; Schwimmer et al. 2014). In turn this can result in changes in expression of heat shock proteins, cortisol production and immune function, and increased risk of cancer (e.g. Dauchy et al. 1997; Bedrosian et al. 2011, 2013; Ashkenazi and Haim 2012; Schwimmer et al. 2014). This suggests that wild populations may also experience

significant health impacts from ALAN. As yet, empirical evidence remains largely lacking, but so do studies whose goal is to obtain this evidence.

Time partitioning

The timing of life history events is fundamental to fitness in perhaps most organisms (Bradshaw and Holzapfel 2010). Many use the timings of dawn and dusk, and/or day length as a cue for daily (e.g. foraging) and phenological events (e.g. growth, reproduction, migration), as in much of the world (foremost excepting environments with limited seasonal variation and where information is unreliable or inaccessible) these are more reliable than key alternatives (Fig. 6; such as temperature), although the latter may be used to modulate responses (e.g. Basler and Körner 2012; Saikkonen et al. 2012; Helm et al. 2013). ALAN has been found to cause changes in many such timings, including of singing (Nordt and Klenke 2013), activity (Boldogh et al. 2007; Dominoni et al. 2013b, 2014), foraging (Bakken and Bakken 1977; Bird et al. 2004; Lebbin et al. 2007), and births (Boldogh et al. 2007).




Fig. 6a–d Actual temperature, and day length calculated in two different ways for 2007 at Penryn Campus, Cornwall. **a** Temperature (°C) recorded on a minute-by-minute basis, **b** mean daily temperature, **c** estimated day length based on latitude and the angle of the sun, and **d** estimated day length based on minute-by-minute energy

measurements (kW m^{-2} ; beginning of day was identified as the time when values increased from 0 to >0 and end when they decreased from >0 to 0. Day length was then assumed to be the time between these two points). Data courtesy of K. Anderson

Interspecific interactions

Diurnal and nocturnal species assemblages can differ markedly in their taxonomic and structural composition (e.g. Guevara and Avilés 2013; Bennie et al. 2014b). They typically comprise mixes of species that exhibit more obligate or more facultative time-partitioning behaviours (although most species probably have some flexibility), and there is some tendency for more detailed studies to reveal previously undocumented variation in time partitioning. The time-partitioning behaviour of animal species has been shown to be influenced by, and to influence, both competitive and predator–prey interspecific interactions (e.g. Fenn and Macdonald 1995; Schwartz et al. 2010; Pita et al. 2011). ALAN has thus been found to be able to change these interactions (e.g. Arlettaz et al. 2000). This can occur in two ways (Fig. 7). First, ALAN can directly influence the time partitioning of individuals of a focal species, with consequences for its interactions with others. Alternatively,

ALAN can influence the time partitioning of competitors, prey or predators, causing in turn that of the focal species to change.

Effects on aquatic ecosystems

The effects of ALAN on aquatic ecosystems, whether freshwater or marine, have been much less frequently studied than for terrestrial ecosystems (for reviews see Montevercchi 2006; Moore et al. 2006; Nightingale et al. 2006; Perkin et al. 2011; Davies et al. 2014). This is logical in as much as the majority of sources of ALAN are themselves terrestrial, and a greater proportion of the land mass is subject to ALAN than of the oceans. However, the high proportion of the global human population that is distributed close to major watercourses, lakes and along coasts (Small and Cohen 2004) suggests that some kinds of aquatic ecosystems may be disproportionately subject to ALAN (e.g.




Fig. 7 Routes by which influences of ALAN on interspecific interactions have consequences for community structure and ecosystem function, process and services

Aubrecht et al. 2008; Davies et al. 2014). Although many of the same influences are associated both with terrestrial and aquatic ecosystems, again, we highlight selected issues that are emerging, in our opinion, as likely to be of key significance for the latter.

Reproduction

Natural light regimes, and notably lunar cycles, are used widely by marine organisms, either in isolation or in combination with other environmental cues, to time key reproductive activities. These organisms include polychaetes, cnidarians, echinoderms, and arthropods (e.g. Rudloe 1980; Lessios 1991; Tanner 1996; Bentley et al. 1999; Naylor 1999; Mercier et al. 2007; Harrison 2011). ALAN has significant potential to provide misleading information about when to time these reproductive activities, particularly for species reproducing in coastal waters. In turn, this could reduce synchrony of these activities amongst individuals (with consequences for fertilization success, predator satiation, etc.), and interactions with other important environmental phenomena, such as oceanographic processes and resource availability.

Movements

Much attention has been paid to the influence of ALAN on the movements of organisms in terrestrial ecosystems (e.g. Frank 1988; Beier 1995; Gauthreaux and Belser 2006; Rydell 2006; Stone et al. 2009; Polak et al. 2011). However, patterns of light are arguably more important cues for movement in aquatic systems, where alternatives (e.g. use of landmarks) may often be severely lacking (Davies et al. 2014). Indeed, ALAN has already been shown to influence the movements (local, dispersive, migratory) of aquatic groups as diverse as zooplankton (Moore et al. 2000), fish (Ryer et al. 2009; Riley et al. 2012, 2013),

turtles (Philibosian 1976; Lorne and Salmon 2007; Bourgeois et al. 2009) and birds (Telfer et al. 1987; Rodríguez and Rodríguez 2009; Rodrigues et al. 2012; Rodríguez et al. 2012a, b). Of particular concern is the extent to which ALAN impacts on the vertical diel movements of zooplankton, which are argued to constitute the largest synchronized movement of biomass globally, with huge impacts on carbon cycling and ecosystem functioning. These diel movements have been found to occur even during the polar night, regulated by variation in light intensity at levels below the threshold of human perception (Berge et al. 2009). This suggests that such movements may be highly susceptible to ALAN.

Community structure

Effects of ALAN on births and deaths of species and/or their movements will result in shifts in community structure. Because the influences on demographic rates are likely to be site, time and species specific (Gaston and Bennie 2014), and to lead to shifts in competitive and predator–prey interactions, it is virtually impossible to predict *a priori* the form that these changes in community structure will take, and they are likely to appear quite idiosyncratic. Nonetheless, these changes have indeed been documented. For example, Meyer and Sullivan (2013) detail changes in the taxonomic and functional composition of aquatic and terrestrial invertebrate communities when natural streams were experimentally subjected to ALAN, reflecting changes in the fluxes between the two faunas. Likewise, Becker et al. (2013) document changes in the trophic and size structure of estuarine fish assemblages when artificial lighting conditions were manipulated. Given the links between community structure and composition and ecosystem functions and processes, ALAN will inevitably impact the latter, although to our knowledge these effects remain to be documented.

Interactions and synergies

ALAN is, of course, only one of many anthropogenic pressures to which natural environments are subject, including habitat loss and fragmentation, climate change, excessive nutrient load and other forms of pollution, overexploitation and unsustainable use, and invasive alien species. One could potentially ask how ALAN compares in terms of the relative impact that it has. However, given that the different pressures seldom act in isolation it seems more pertinent to consider their interactions and synergies with ALAN. Here we highlight several such possibilities.

Habitat loss and fragmentation

Most consideration of levels of habitat loss and fragmentation and their effects on ecosystems and biodiversity pertains to structural changes, such as in different kinds of land cover, in the physical sizes of patches, and in their degree of connectivity or isolation (Hanski 2005). ALAN can exacerbate these effects in ways that are not apparent from the daylight images (from aerial photographs and satellite sensors) that are typically employed to make such assessments. It renders areas of structurally unaltered habitat unusable by some organisms, available to others, and creates barriers to or corridors for movement that fragment and connect landscapes in different ways (Beier 1995, 2006; Eisenbeis 2006; Frank 2006; Stone et al. 2012; Threlfall et al. 2013). Indeed, full understanding of habitat loss and fragmentation needs to account both for diurnal and nocturnal effects, which may be rather different. Given the high proportion of species that are nocturnal (in addition to those that are crepuscular and cathemeral) in some groups of major conservation concern [e.g. 69 % of mammals (Bennie et al. 2014b)], it seems likely that the full impact of habitat loss and fragmentation has often been markedly underestimated.

Climate change

It has previously been observed that biotic responses to anthropogenic climate change are critically dependent on the fact that whilst temperatures are changing, geographic and annual patterns in natural light cycles are not (Bradshaw and Holzapfel 2010). Given that organisms use day length as a cue for anticipating seasonal changes, this creates strong selection pressures for altering the timing of seasonal events, some of which they are able to respond to and some of which they are not (Bradshaw and Holzapfel 2010). ALAN serves to complicate this picture. Typically it serves locally to extend apparent day lengths, and to obscure their seasonal patterns. In combination, higher temperatures and increased light levels at night may

allow species that are able to utilize the night light niche to extend their hours of activity (Garber 1978; Heiling 1999) and may alter predation patterns and/or competitive interactions.

Other forms of pollution

ALAN can be seen as a stressor on the physiologies of many organisms, particularly as mediated through melatonin production. It seems likely that this will be more challenging to deal with in the presence of other forms of pollution, which are imposing other demands. ALAN can also exacerbate other forms of pollution in a more direct fashion. Stark et al. (2011) showed that artificial lights can change nighttime atmospheric nitrogen chemistry. Dim nocturnal light has also been found to inhibit recovery from leaf damage caused by atmospheric ozone in some species of clover *Trifolium* (Futsaether et al. 2009; Vollsnes et al. 2009).

Overexploitation and unsustainable use

The harvesting of many marine species (e.g. shrimp, squid, fish) employs the use of artificial nighttime lights as attractants, sometimes on an industrial scale (Kiyofuji and Saitoh 2004). The effects of this source of ALAN on unexploited organisms is largely unknown; however, it seems likely to be potentially marked, particularly given the responsiveness of most marine organisms to light. Some forms of terrestrial harvesting, such as spotlighting, also employ ALAN, but this is on a more localized and transient scale.

Invasive alien species

Several examples exist of invasive alien species that have rapidly adapted photoperiodic responses to their new environment [over a few decades (Gomi and Takeda 1991; Urbanski et al. 2012)]. The ability to adapt phenology to changing light regimes may be a key determinant of success in colonizing latitudes outside of a species' historical range (Bradshaw and Holzapfel 2010), which is critical when species are introduced to new regions or spread due to climate change. Such phenotypic flexibility in photoperiodism may also be important in species response to extended hours of light due to ALAN.

Major uncertainties

Although the potential for ALAN to have significant biological impacts has long been recognized and a large body of studies has ensued (Rich and Longcore 2006; Gaston et al. 2013), substantial uncertainties remain. We would

highlight the following as being, in our view, particularly significant gaps in knowledge:

1. What are the biological effects of skyglow resulting from ALAN? Whilst the effects of more direct lighting are increasingly well understood, those of skyglow remain poorly explored. Studies to do so are challenging, although suggestions as to how these might be constructed have been made (Kyba and Hölker 2013).
2. What are the effects of ALAN on photosynthesis? The widespread use of artificial lighting in growing plants under controlled conditions suggests the potential for ALAN to influence photosynthesis. However, studies of these impacts, and of those on the photophysiology of plants and phytoplankton more broadly, remain scarce and it is difficult to extract any broad conclusions (Gaston et al. 2013; Poulin et al. 2013).
3. Do the influences of ALAN on stress and disease demonstrated for animals in the laboratory extend to the wild? Particularly because of concerns about effects of ALAN on human health, more laboratory studies of potential ecological relevance have been conducted than for most ecological issues. However, there are undoubtedly large differences between the ALAN treatments used in laboratory settings and what the majority of organisms experience in the field, especially when those organisms are mobile.
4. What shape are dose–response curves for ALAN? The literature on the ecological effects of ALAN is dominated by studies in which comparison is made, observationally or experimentally, between the state of a given ecological variable with and without ALAN, or perhaps with two different forms of ALAN (usually differing in intensity, but sometimes light spectrum). Almost nothing is known about the form of dose–response curves for ALAN, and thus critically how responses are likely to change when ALAN attains different levels.
5. What are the impacts of ALAN on ecosystem functions and processes? Broadly speaking, most is known about the impacts of ALAN on the physiology and behaviour of organisms, less about those on population dynamics, little about those on communities, and almost nothing about the impacts on ecosystem functions and processes. Given that ALAN can influence the abundances of species and trophic interactions there seems little doubt that such effects on ecosystem functions and processes do occur.

Future prospects and management options

ALAN will undoubtedly continue to spread globally, particularly given the rapid rate of population growth and

industrialization in many countries (Small and Elvidge 2013), although the pattern may be more complex in industrialised regions (Bennie et al. 2014a). However, it will also alter in form, as the predominant technologies employed change. From an ecological perspective key changes include increasing use of:

1. Central management systems in developed countries by which the timing and intensity of grid-based lighting can be controlled, already resulting in some broad-scale decreases in lighting during periods when it is not needed (Bennie et al. 2014a);
2. White light technologies, especially LEDs. LEDs can be modified to control the spectral composition of lighting, can require lower wattage for a given level of illumination than more traditional light sources, provide high light output for low radiant heat, can distribute light more uniformly and thus allow lower levels of lighting to be employed, are dimmable and more tolerant of switching on and off, and have long life times before failure (US Department of Energy 2012). Typical white LEDs emit considerably more light in the blue portion of the spectrum than conventional ‘white’ lighting (Fig. 4); while LED technology may allow more control over the spectra emitted, a movement towards white LED-based lighting systems is likely to lead to greater emissions within the blue portion of the spectra. LEDs also raise concerns around hazardous waste and resource depletion (Lim et al. 2011).
3. Off-grid lighting in developing countries, likely principally using combinations of LEDs and solar power (Mills 2005).

Various management options have been highlighted by which the ecological impacts of ALAN can be limited and/or reduced (Falchi et al. 2011; Gaston et al. 2012). In brief, these are:

1. Maintaining and creating dark areas. Faced with progressive loss of dark areas, particularly in more heavily urbanized regions it is important to protect those that remain and where possible recover others. There are a number of initiatives to identify presently dark areas, to highlight this status, and to encourage steps by which it is maintained [UNESCO 2009; International Dark Sky Association (IDSA) 2013; IUCN 2013]. There are also initiatives to encourage communities to reduce their overall levels of ALAN (IDSA 2001). Even quite localized changes (e.g. switching off a few key lights) can serve to reduce particular impacts (e.g. Yurk and Trites 2000).
2. Reducing light trespass. Lighting devices generally remain quite poorly designed and/or managed for the

- purposes of only directing light where it is actually required. Resolving this problem provides a ready means of dramatically decreasing the impacts of ALAN at a local scale. Indeed, reduction of light trespass has been shown to reduce the impacts of ALAN on organisms (e.g. Reed et al. 1985).
3. Dimming. Many areas are overlit compared with what is practically required. This provides opportunities for dimming of lighting without major negative consequences for human populations. Indeed, substantial progressive dimming may be possible without these populations being able to perceive that this is the case. The introduction of LED lighting provides further opportunities, given that colour rendering may be improved at lower intensities of lighting.
 4. Part-night lighting. Many areas are presently lit at times of day when this carries limited or negligible human benefit. Particularly following the global financial crisis, and pressure on public expenditure, numerous towns and cities have sought to reduce energy costs (and CO₂ emissions) by switching off street lights in low-risk areas from late at night until the early hours of the morning (Gaston et al. 2012). The ecological benefits of such part-night lighting remain poorly understood, and may only influence a relative minority of species that use the heart of the night rather than hours around dusk and dawn.
 5. Targeting spectra. There are doubtless substantial opportunities to reduce the ecological impacts of ALAN by employing spatially more nuanced approaches to the use of lighting with different spectral properties. Developing alternatives to presently installed systems, which have often evolved as different technologies have become available and affordable, will require balancing of multiple pressures. These include cost, practicality, human perceived and actual need, and environmental concerns. In general, there would seem to be a number of advantages to the use of reddened spectra in environmentally more sensitive areas because, relative to white or blue sources, these reduce skyglow (Kyba et al. 2012), penetrate the water column to a lesser extent, have less influence on melatonin levels and circadian rhythms of species (Bayarri et al. 2002; Lockley et al. 2003), are less attractive to some organisms (e.g. Evans et al. 2007b; Cowan and Gries 2009; Somers-Yeates et al. 2013) and less repellent to others (e.g. Downs et al. 2003, Widder et al. 2005). However, this is not always the case—reddened light may disrupt the magnetic orientation of migratory birds (Wiltschko et al. 1993) and light of lower wavelengths may be less disruptive to these species (Poot et al. 2008). Furthermore, reddened light sources have a stronger influence on plant development through the impact on phytochromes, which respond to the ratio of red to far red light (Stutte 2009).
- Arguably, there is a trade-off between the economic costs and the perceived social costs associated with implementing these different strategies to managing ALAN. This constitutes the major challenge to limiting its ecological impacts.
- Acknowledgments** We are grateful to two anonymous reviewers for their comments on a previous version of the manuscript, and to S. Rouillard for assistance with figures. The research leading to this paper has received funding from the European Research Council under the European Union's Seventh Framework Programme (FP7/2007–2013)/ERC grant agreement no. 268504 to K. J. G.
- Open Access** This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

- Amaral S, Monteiro AMV, Camara G, Quintanilha JA (2006) DMSP/OLS night-time light imagery for urban population estimates in the Brazilian Amazon. *Int J Remote Sens* 27:855–870
- Arendt J (1998) Melatonin and the pineal gland: influence on mammalian seasonal and circadian physiology. *Rev Reprod* 3:13–22
- Arlettaz R, Godat S, Meyer H (2000) Competition for food by expanding pipistrelle bat populations (*Pipistrellus pipistrellus*) might contribute to the decline of lesser horseshoe bats (*Rhinolophus hipposideros*). *Biol Conserv* 93:55–60
- Ashkenazi L, Haim A (2012) Light interference as a possible stressor altering HSP70 and its gene expression levels in brain and hepatic tissues of golden spiny mice. *J Exp Biol* 215:4034–4040
- Aubrecht C, Elvidge CD, Longcore T, Rich C, Safran J, Strong AE, Eakin CM, Baugh K, Tuttle BT, Howard AT, Erwin EH (2008) A global inventory of coral reef stressors based on satellite observed nighttime lights. *Geocarto Int* 23:467–479
- Bachleitner W, Kempinger L, Wülbeck C, Rieger D, Helfrich-Förster C (2007) Moonlight shifts the endogenous clock of *Drosophila melanogaster*. *Proc Nat Acad Sci USA* 104:3538–3543
- Bakken LE, Bakken GS (1977) American redstart feeding by artificial light. *Auk* 94:373–374
- Ball JR, Lukianchuk K, Bayne EM (2011) Nocturnal provisioning by Swainson's thrush. *Wilson J Ornithol* 123:508–514
- Basler D, Körner C (2012) Photoperiod sensitivity of bud burst in 14 temperate forest tree species. *Agric For Meteorol* 165:73–81
- Bayarri MJ, Madrid JA, Sánchez-Vázquez FJ (2002) Influence of light intensity, spectrum and orientation on sea bass plasma and ocular melatonin. *J Pineal Res* 32:34–40
- Becker A, Whitfield AK, Cowley PD, Järnegen J, Næsje TF (2013) Potential effects of artificial light associated with anthropogenic infrastructure on the abundance and foraging behaviour of estuary-associated fishes. *J Appl Ecol* 50:43–50
- Bedrosian TA, Fonken LK, Walton JC, Nelson RJ (2011) Chronic exposure to dim light at night suppresses immune responses in Siberian hamsters. *Biol Lett* 7:468–471
- Bedrosian TA, Galan A, Vaughn CA, Weil ZM, Nelson RJ (2013) Light at night alters daily patterns of cortisol and clock proteins in female Siberian hamsters. *J Neuroendocrinol* 25:590–596

- Beier P (1995) Dispersal of juvenile cougars in fragmented habitat. *J Wildl Manage* 59:228–237
- Beier P (2006) Effects of artificial night lighting on terrestrial mammals. In: Rich C, Longcore T (eds) *Ecological consequences of artificial night lighting*. Island Press, Washington, pp 19–42
- Bennie J, Davies T, Duffy J, Inger R, Gaston KJ (2014a) Contrasting trends in light pollution across Europe. *Sci Rep* 4:3789
- Bennie J, Duffy JP, Inger R, Gaston KJ (2014b) The biogeography of time partitioning in mammals. *Proc Nat Acad Sci USA* (in press)
- Bentley MG, Olive PJW, Last K (1999) Sexual satellites, moonlight and the nuptial dances of worms: the influence of the moon on the reproduction of marine animals. *Earth Moon Planets* 85–86:67–84
- Berge J, Cottier F, Last KS, Varpe Ø, Leu E, Soreide J, Eiane K, Falk-Petersen S, Willis K, Nygård H, Vogedes D, Griffiths C, Johnsen G, Lorentzen D, Brierley AS (2009) Diel vertical migration of Arctic zooplankton during the polar night. *Biol Lett* 5:69–72
- Biggs JD, Fouche T, Bilki F, Zadnik MG (2012) Measuring and mapping the night sky brightness of Perth, Western Australia. *Mon Not R Astron Soc* 421:1450–1464
- Bird BL, Branch LC, Miller DL (2004) Effects of coastal lighting on foraging behavior of beach mice. *Conserv Biol* 18:1435–1439
- Bishop JE (1969) Light control of aquatic insect activity and drift. *Ecology* 50:371–380
- Bogard P (2013) *The end of night: searching for natural darkness in an age of artificial light*. Fourth Estate, London
- Boldogh S, Dobrosi D, Samu P (2007) The effects of the illumination of buildings on house-dwelling bats and its conservation consequences. *Acta Chiropterol* 9:527–534
- Bourgeois S, Gilot-Fromont E, Viallefant A, Boussamba F, Deem SL (2009) Influence of artificial lights, logs and erosion on leatherback sea turtle orientation at Pongara National Park, Gabon. *Biol Conserv* 142:85–93
- Bradshaw WE, Holzapfel CM (2010) Light, time, and the physiology of biotic response to rapid climate change in animals. *Annu Rev Physiol* 72:147–166
- Brainard GC, Richardson BA, Hurlbut EC, Steinlechner S, Matthews SA, Reiter RJ (1984) The influence of various irradiances of artificial light, twilight, and moonlight on the suppression of pineal melatonin content in the Syrian hamster. *J Pineal Res* 1:105–119
- Chen X, Nordhaus WD (2011) Using luminosity as a proxy for economic statistics. *Proc Nat Acad Sci USA* 108:8589–8594
- CIA (2011) *The world factbook*. <https://www.cia.gov/library/publications/the-world-factbook/index.html>
- Cinzano P, Elvidge CD (2004) Night sky brightness at sites from DMSP-OLS satellite measurements. *Mon Not R Astron Soc* 353:1107–1116
- Cinzano P, Falchi F, Elvidge CD (2001) The first world atlas of the artificial night sky brightness. *Mon Not R Astron Soc* 328:689–707
- Clarke JA, Chopko JT, Mackessy SP (1996) The effect of moonlight on activity patterns of adult and juvenile prairie rattlesnakes (*Crotalus viridis viridis*). *J Herpetol* 2:192–197
- Cos S, Mediavilla D, Martínez-Campa C, González A, Alonso-González C, Sánchez-Barceló EJ (2006) Exposure to light-at-night increases the growth of DMBA-induced mammary adenocarcinomas in rats. *Cancer Lett* 235:266–271
- Cowan T, Gries G (2009) Ultraviolet and violet light: attractive orientation cues for the Indian meal moth, *Plodia interpunctella*. *Entomol Exp Appl* 131:148–158
- Crawford DL (2000) Light pollution, an environmental problem for astronomy and for mankind. *Mem Soc Astron Ital* 71:11–40
- Dauchy RT, Sauer LA, Blask DE, Vaughan GM (1997) Light contamination during the dark phase in “photoperiodically controlled” animal rooms: effect on tumor growth and metabolism in rats. *Lab Anim Sci* 47:511–518
- Davies TW, Bennie J, Gaston KJ (2012) Street lighting changes the composition of invertebrate communities. *Biol Lett* 8:764–767
- Davies TW, Bennie J, Inger R, Gaston KJ (2013) Artificial light alters natural regimes of night-time sky brightness. *Sci Rep* 3:1722
- Davies TW, Duffy J, Bennie J, Gaston KJ (2014) Marine light pollution: nature, extent and ecological implications. *Frontiers Ecol Environ* 12:347–355
- Dice JD (1945) Minimum intensities of illumination under which owls can find dead prey by sight. *Am Nat* 79:385–416
- DMSP/OLS (2012) Night time lights data set (Version 4). NOAA Earth Observation Group, Boulder, CO. <http://ngdc.noaa.gov/eo/dmsp/downloadV4composites.html>. Accessed Oct 2012
- Doll CNH, Muller J-P, Morley JG (2006) Mapping regional economic activity from night-time light satellite imagery. *Ecol Econ* 57:75–92
- Dominoni DM, Helm B, Lehmann M, Dowse HB, Partecke J (2013a) Clocks for the city: circadian differences between forest and city songbirds. *Proc R Soc B* 280:20130593
- Dominoni D, Quetting M, Partecke J (2013b) Artificial light at night advances avian reproductive physiology. *Proc R Soc B* 280:20123017
- Dominoni DM, Carmona-Wagner EO, Hofmann M, Kranstauber B, Partecke J (2014) Individual-based measurements of light intensity provide new insights into the effects of artificial light at night on daily rhythms of urban-dwelling songbirds. *J Anim Ecol* 83:681–692
- Downs NC, Beaton V, Guest J, Polanski J, Robinson SL, Racey PA (2003) The effects of illuminating the roost entrance on the emergence behavior of *Pipistrellus pygmaeus*. *Biol Conserv* 111:247–252
- Dwyer RG, Bearhop S, Campbell HA, Bryant DM (2012) Shedding light on light: benefits of anthropogenic illumination to a nocturnally foraging shorebird. *J Anim Ecol* 82:478–485
- Eisenbeis G (2006) Artificial night lighting and insects: attraction of insects to streetlamps in a rural setting in Germany. In: Rich C, Longcore T (eds) *Ecological consequences of artificial night lighting*. Island Press, Washington, pp 281–304
- Elvidge CD, Baugh KE, Dietz JB, Bland T, Sutton PC, Kroehl HW (1999) Radiance calibration of DMSP-OLS low-light imaging data of human settlements. *Remote Sens Environ* 68:77–88
- Elvidge CD, Imhoff ML, Baugh KE, Hobson VR, Nelson I, Safran J, Dietz JB, Tuttle BT (2001) Night-time lights of the world: 1994–1995. *ISPRS J Photogramm Remote Sens* 56:L81–L99
- Elvidge CD, Keith DM, Tuttle BT, Baugh KE (2010) Spectral identification of lighting type and character. *Sensors* 10:3961–3988
- Evans JA, Elliott JA, Gorman MR (2007a) Circadian effects of light no brighter than moonlight. *J Biol Rhythms* 22:356–367
- Evans WR, Akashi Y, Altman NS, Manville AM II (2007b) Response of night-migrating songbirds in cloud to colored and flashing light. *North Am Birds* 60:476–488
- Falchi F, Cinzano P, Elvidge CD, Keith DM, Haim A (2011) Limiting the impact of light pollution on human health, environment and stellar visibility. *J Environ Manage* 92:2714–2722
- Falkenberg JC, Clarke JA (1998) Microhabitat use of deer mice: effects of interspecific interaction risks. *J Mamm* 79:558–565
- Fenn MGP, Macdonald DW (1995) Use of middens by red foxes: risk reverses rhythms of rats. *J Mamm* 76:130–136
- Frank KD (1988) Impact of outdoor lighting on moths: an assessment. *J Lepid Soc* 42:63–93
- Frank KD (2006) Effect of artificial night lighting on moths. In: Rich C, Longcore T (eds) *Ecological consequences of artificial night lighting*. Island Press, Washington, pp 305–344
- Futsaether C, Vollsnes A, Kruse OMO, Otterholt E, Kvaal K, Eriksen AB (2009) Effects of the Nordic photoperiod on ozone

- sensitivity and repair in different clover species studied using infrared imaging. *Ambio* 38:437–443
- Garber SD (1978) Opportunistic feeding behaviour of *Anolis cristatellus* (Iguanidae: Reptilia) in Puerto Rico. *Trans Kansas Acad Sci* 81:19–80
- Garstang RH (1986) Model for artificial night-sky illumination. *Publ Astron Soc Pacific* 98:364–375
- Gaston KJ (2010) Urbanisation. In: Gaston KJ (ed) *Urban ecology*. Cambridge University Press, Cambridge, pp 10–34
- Gaston KJ, Bennie J (2014) Demographic effects of artificial nighttime lighting on animal populations. *Environ Rev* (in press)
- Gaston KJ, Davies TW, Bennie J, Hopkins J (2012) Reducing the ecological consequences of night-time light pollution: options and developments. *J Appl Ecol* 49:1256–1266
- Gaston KJ, Bennie J, Davies TW, Hopkins J (2013) The ecological impacts of nighttime light pollution: a mechanistic appraisal. *Biol Rev* 88:912–927
- Gauthreaux SA Jr, Belser CG (2006) Effects of artificial night lighting on migrating birds. In: Rich C, Longcore T (eds) *Ecological consequences of artificial night lighting*. Island Press, Washington, pp 67–93
- Gomi T, Takeda M (1991) Geographic variation in photoperiodic responses in an introduced insect, *Hyphantria cunea* Drury (Lepidoptera: Arctiidae) in Japan. *Appl Entomol Zool* 26:357–363
- Grant R, Halliday T, Chadwick E (2012) Amphibians' response to the lunar synodic cycle—a review of current knowledge, recommendations, and implications for conservation. *Behav Ecol* 24:53–62
- Guevara J, Avilés L (2013) Community-wide body size differences between nocturnal and diurnal insects. *Ecology* 94:537–543
- Hale JD, Davies G, Fairbrass AJ, Matthews TJ, Rogers CDF, Sadler JP (2013) Mapping lightscapes: spatial patterning of artificial lighting in an urban landscape. *PLoS One* 8:e61460
- Hanski I (2005) *The shrinking world: ecological consequences of habitat loss*. International Ecology Institute, Oldendorf
- Harrison PL (2011) Sexual reproduction of scleractinian corals. In: Dubinsky Z, Stambler N, Harrison PL (eds) *Coral reefs: an ecosystem in transition*. Springer, Dordrecht, pp 59–85
- Heiling AM (1999) Why do nocturnal orb-web spiders (Araneidae) search for light? *Behav Ecol Sociobiol* 46:43–49
- Helm B, Ben-Shlomo R, Sheriff MJ, Hut RA, Foster R, Barnes BM, Dominoni D (2013) Annual rhythms that underlie phenology: biological time-keeping meets environmental change. *Proc R Soc B* 280:20130016
- Hölker F, Moss T, Griefahn B, Kloas W, Voigt CC, Henckel D, Hänel A, Kappeler PM, Völker S, Schwöpe A, Franke S, Uhrlandt D, Fischer J, Klenke R, Wolter C, Tockner K (2010) The dark side of light: a transdisciplinary research agenda for light pollution policy. *Ecol Soc* 15:13
- IDSA (2001) International dark sky communities—guidelines. www.darksky.org/international-dark-sky-places
- IDSA (2013) Dark sky park program criteria. www.darksky.org
- IUCN (2013) World list of dark sky protected areas. IUCN Dark Skies Advisory Group. www.darkskeyparks.org
- Jennings S, Lee J (2012) Defining fishing grounds with vessel monitoring system data. *ICES J Mar Sci* 69:51–63
- Johnson K (1979) Control of lampenflora at Waitomo Caves, New Zealand. Cave management in Australia III: proceedings of the 3rd Australasian Cave Tourism and Management Conference, Mount Gambier. South Australian National Parks and Australian Speleological Federation, Adelaide, pp 105–122
- Kareiva P, Watts S, McDonald R, Boucher T (2007) Landscapes and ecosystems for human welfare. *Science* 316:1866–1869
- Kempnaers B, Borgström P, Löes P, Schlicht E, Valcu M (2010) Artificial night lighting affects dawn song, extra-pair siring success, and lay date in songbirds. *Curr Biol* 20:1735–1739
- Kiyofuji H, Saitoh S-I (2004) Use of nighttime visible images to detect Japanese common squid *Todarodes pacificus* fishing areas and potential migration routes in the Sea of Japan. *Mar Ecol Prog Ser* 276:173–186
- Kramer KM, Birney EC (2001) Effect of light intensity on activity patterns of Patagonian leaf-eared mice, *Phyllotis xanthopygus*. *J Mamm* 82:535–544
- Kronfeld-Schor N, Dominoni D, de la Iglesia H, Levy O, Herzog ED, Dayan T, Helfrich-Forster C (2013) Chronobiology by moonlight. *Proc R Soc B* 280:20123088
- Kuechly HU, Kyba CCM, Ruhtz T, Lindemann C, Wolter C, Fischer J, Höcker F (2012) Aerial survey and spatial analysis of sources of light pollution in Berlin, Germany. *Remote Sens Environ* 126:39–50
- Kuijper DPJ, Schut J, van Dulleman D, Toorman H, Goossens N, Ouwehand J, Limpens HJGA (2008) Experimental evidence of light disturbance along the commuting routes of pond bats (*Myotis dasycneme*). *Lutra* 51:37–49
- Kurtze W (1974) Synökologische und experimentelle Untersuchungen zur Nachtaktivität von Insekten. *Zool Jahrb Abt Syst Ökol Geogr Tier* 101:297–344
- Kyba CCM, Höcker F (2013) Do artificially illuminated skies affect biodiversity in nocturnal landscapes? *Landsc Ecol* 28:1637–1640
- Kyba CCM, Ruhtz T, Fischer J, Höcker F (2011a) Lunar skylight polarization signal polluted by urban lighting. *J Geophys Res* D 116:D24106
- Kyba CCM, Ruhtz T, Fischer J, Höcker F (2011b) Cloud coverage acts as an amplifier for ecological light pollution in urban ecosystems. *PLoS One* 6:e17307
- Kyba CCM, Ruhtz T, Fischer J, Höcker F (2012) Red is the new black: how the colour of urban skyglow varies with cloud cover. *Mon Not R Astron Soc* 425:701–708
- Kyba CCM, Wagner JM, Kuechly HU, Walker CE, Elvidge CD, Falchi F, Ruhtz T, Fischer J, Höcker F (2013) Citizen science provides valuable data for monitoring global night sky luminance. *Sci Rep* 3:1835
- Larsen LO, Pedersen JN (1982) The snapping response of the toad *Bufo bufo*, towards prey dummies at very low light intensities. *Amphib-Reptil* 2:321–327
- Le Tallec T, Perret M, Théry M (2013) Light pollution modifies the expression of daily rhythms and behavior patterns in a nocturnal primate. *PLoS One* 8:e79250
- Lebbin DJ, Harvey MG, Lenz TC, Andersen MJ, Ellis JM (2007) Nocturnal migrants foraging at night by artificial light. *Wilson J Ornith* 119:506–508
- Lessios HA (1991) Presence and absence of monthly reproductive rhythms among eight Caribbean echinoids off the coast of Panama. *J Exp Mar Biol Ecol* 153:27–47
- Levin N, Duke Y (2012) High spatial resolution night-time images for demographic and socio-economic studies. *Remote Sens Environ* 119:1–10
- Li X, Chen X, Zhao Y, Xu J, Chen F, Li H (2012) Automatic intercalibration of night-time light imagery using robust regression. *Remote Sens Lett* 4:46–55
- Lim S-R, Kang D, Ogunseitan OA, Schoenung JM (2011) Potential environmental impacts of light-emitting diodes (LEDs): metallic resources, toxicity, and hazardous waste classification. *Environ Sci Technol* 45:320–327
- Lockley SW, Brainard GC, Czeisler CA (2003) High sensitivity of the human circadian melatonin rhythm to resetting by short wavelength light. *J Clin Endocrinol Metabolism* 88:4502–4505
- Longcore T, Rich C (2004) Ecological light pollution. *Front Ecol Environ* 2:191–198
- Longcore T, Rich C (2006) Synthesis. In: Rich C, Longcore T (eds) *Ecological consequences of artificial night lighting*. Island Press, Washington, pp 413–430

- Lorne JK, Salmon M (2007) Effects of exposure to artificial lighting on orientation of hatchling sea turtles on the beach and in the ocean. *Endanger Sp Res* 3:23–30
- Lyytimäki J, Tapio P, Assmuth T (2012) Unawareness in environmental protection: the case of light pollution from traffic. *Land Use Policy* 29:598–604
- Mainster MA, Timberlake GT (2003) Why HID headlights bother older drivers. *Br J Ophthalmol* 87:113–117
- Mazor T, Levin N, Possingham HP, Levy Y, Rocchini D, Richardson AJ, Kark S (2013) Can satellite-based night lights be used for conservation? The case of nesting sea turtles in the Mediterranean. *Biol Conserv* 159:63–72
- Meeus J (2008) *Astronomical formulae for calculators*, 4th edn. Atlantic Books, London
- Mercier A, Ycaza RH, Hamel JF (2007) Long-term study of gamete release in a broadcast-spawning holothurian: predictable lunar and diel periodicities. *Mar Ecol Prog Ser* 329:179–189
- Meyer LA, Sullivan SMP (2013) Bright lights, big city: influences of ecological light pollution on reciprocal stream-riparian invertebrate fluxes. *Ecol Appl* 23:1322–1330
- Millennium Ecosystem Assessment (2005) *Ecosystems and human well-being: current state and trends*, vol I. Island Press, Washington
- Miller MW (2006) Apparent effects of light pollution on singing behavior of American robins. *Condor* 108:130–139
- Miller SD, Mills SP, Elvidge CD, Lindsey DT, Lee TF, Hawkins JD (2012) Suomi satellite brings to light a unique frontier of nighttime environmental sensing capabilities. *Proc Nat Acad Sci USA* 109:15706–15711
- Mills E (2005) The specter of fuel-based lighting. *Science* 308:1263–1264
- Montevicchi WA (2006) Influences of artificial light on marine birds. In: Rich C, Longcore T (eds) *Ecological consequences of artificial night lighting*. Island Press, Washington, pp 94–113
- Moore MV, Pierce SM, Walsh HM, Kvalvik SK, Lim JD (2000) Urban light pollution alters the diel vertical migration of *Daphnia*. *Verh Int Ver Limnol* 27:779–782
- Moore MV, Kohler SJ, Cheers MS (2006) Artificial light at night in freshwater habitats and its potential ecological effects. In: Rich C, Longcore T (eds) *Ecological consequences of artificial night lighting*. Island Press, Washington, pp 365–384
- NASA (2012) LIS/OTD gridded lightning climatology data set. NASA EoSDIS GHRC DAAC, Huntsville, AL. <http://lightning.nsstc.nasa.gov/data/>. Accessed October 2012
- NASA Land Processes Distributed Active Archive Center (LP DAAC) (2013) MOD14A1. USGS/Earth Resources Observation and Science (ERO) Center, Sioux Falls, SD
- Naylor E (1999) Marine animal behaviour in relation to lunar phase. *Earth Moon Planets* 85–86:291–302
- Nightingale B, Longcore T, Simenstad CA (2006) Artificial night lighting and fishes. In: Rich C, Longcore T (eds) *Ecological consequences of artificial night lighting*. Island Press, Washington, pp 257–276
- Nordt A, Klenke R (2013) Sleepless in town—drivers of the temporal shift in dawn song in urban blackbirds. *PLoS One* 8:e71476
- Organisation Internationale des Constructeurs d'Automobiles (OICA) (2014) <http://www.oica.net/category/vehicles-in-use/>
- Perkin EK, Hölker F, Richardson JS, Sadler JP, Wolter C, Tockner K (2011) The influence of artificial light on stream and riparian ecosystems: questions, challenges, and perspectives. *Ecosphere* 2:122
- Philibosian R (1976) Disorientation of hawksbill turtle hatchlings, *Eretmochelys imbricata*, by stadium lights. *Copeia* 1976:824
- Picchi MS, Avolio L, Azzani L, Brombin O, Camerini G (2013) Fireflies and land use in an urban landscape: the case of *Luciola italica* L. (Coleoptera: Lampyridae) in the city of Turin. *J Insect Conserv* 17:797–805
- Pita R, Mira A, Beja P (2011) Circadian activity rhythms in relation to season, sex and interspecific interactions in two Mediterranean voles. *Anim Behav* 81:1023–1030
- Polak T, Korine C, Yair S, Holderied MW (2011) Differential effects of artificial lighting on flight and foraging behaviour of two sympatric bat species in a desert. *J Zool* 285:21–27
- Poot H, Ens BJ, de Vries H, Donners MAH, Wernand MR, Marquenie JM (2008) Green light for nocturnally migrating birds. *Ecol Soc* 13(2):47
- Poulin C, Bruyant F, Laprise M-H, Cockshutt AM, Vandenhecke JM-R, Huot Y (2013) The impact of light pollution on diel changes in the photophysiology of *Microcystis aeruginosa*. *J Plankton Res* 36:286–291
- Pun CSJ, So CW (2011) Night-sky brightness monitoring in Hong Kong: a city-wide light pollution assessment. *Environ Mon Assess* 184:2537–2557
- Reed JR, Sincok JL, Hailman JP (1985) Light attraction in endangered procellariiform birds: reduction by shielding upward radiation. *Auk* 102:377–383
- Rich C, Longcore T (eds) (2006) *Ecological consequences of artificial night lighting*. Island Press, Washington
- Riley WD, Bendall B, Ives MJ, Edmonds NJ, Maxwell DL (2012) Street lighting disrupts the diel migratory pattern of wild Atlantic salmon, *Salmo salar* L., smolts leaving their natal stream. *Aquaculture* 330–333:74–81
- Riley WD, Davison PI, Maxwell DL, Bendall B (2013) Street lighting delays and disrupts the dispersal of Atlantic salmon (*Salmo salar*) fry. *Biol Conserv* 158:140–146
- Rodrigues P, Aubrecht C, Gil A, Longcore T, Elvidge C (2012) Remote sensing to map influence of light pollution on Cory's shearwater in Sao Miguel Island, Azores Archipelago. *Eur J Wildl Res* 58:147–155
- Rodríguez A, Rodríguez B (2009) Attraction of petrels to artificial lights in the Canary Islands: effects of the moon phase and age class. *Ibis* 151:299–310
- Rodríguez A, Rodríguez B, Curbelo AJ, Pérez A, Marrero S, Negro JJ (2012a) Factors affecting mortality of shearwaters stranded by light pollution. *Anim Conserv* 15:519–526
- Rodríguez A, Rodríguez B, Lucas MP (2012b) Trends in numbers of petrels attracted to artificial lights suggest population declines in Tenerife, Canary Islands. *Ibis* 154:167–172
- Rudloe A (1980) The breeding behavior and patterns of movement of horseshoe crabs, *Limulus polyphemus*, in the vicinity of breeding beaches in Apalachee Bay, Florida. *Estuaries* 3:177–183
- Rydell J (2006) Bats and their insect prey at streetlights. In: Rich C, Longcore T (eds) *Ecological consequences of artificial night lighting*. Island Press, Washington, pp 43–60
- Ryer CH, Stoner AW, Iseri PJ, Spencer ML (2009) Effects of simulated underwater vehicle lighting on fish behavior. *Mar Ecol Prog Ser* 391:97–106
- Saikkonen K, Taulavuori K, Hyvönen T, Gundel PE, Hamilton CE, Vänninen I, Nissinen A, Helander M (2012) Climate change-driven range shifts filtered by photoperiodism. *Nat Clim Change* 2:239–242
- Salmon M, Tolbert MG, Painter DP, Goff M, Reiners R (1995) Behavior of loggerhead sea turtles on an urban beach. II. Hatchling orientation. *J Herpetol* 29:568–576
- Santos CD, Miranda AC, Granadeiro JP, Lourenco PM, Saraiva S, Palmeirim JM (2010) Effects of artificial illumination on the nocturnal foraging of waders. *Acta Oecol* 36:166–172
- Schwartz CC, Cain SL, Podruzny S, Cherry S, Frattaroli L (2010) Contrasting activity patterns of sympatric and allopatric black and grizzly bears. *J Wildl Manage* 74:1628–1638

- Schwimmer H, Metzger A, Pilosof Y, Szyf M, Machnes ZM, Fares F, Harel O, Haim A (2014) Light at night and melatonin have opposite effects on breast cancer tumors in mice assessed by growth rates and global DNA methylation. *Chronobiol Int* 31:144–150
- Sharma VK, Chandrashekar MK, Nongkynrih P (1997) Daylight and artificial light phase response curves for the circadian rhythm in locomotor activity of the field mouse *Mus booduga*. *Biol Rhythm Res* 28(Suppl 1):39–40
- Small C, Cohen JE (2004) Continental physiography, climate, and the global distribution of human population. *Curr Anthropol* 45:269–277
- Small C, Elvidge CD (2011) Mapping decadal change in anthropogenic night light. *Proc Environ Sci* 7:353–358
- Small C, Elvidge CD (2013) Night on Earth: mapping decadal changes of anthropogenic night light in Asia. *Int J Appl Earth Observ Geoinf* 22:40–52
- Somers-Yeates R, Hodgson D, McGregor PK, Spalding A, French-Constant RH (2013) Shedding light on moths: shorter wavelengths attract noctuids more than geometrids. *Biol Lett* 9:20130376
- Stark H, Brown SS, Wong KW, Stutz J, Elvidge CD, Pollack IB, Ryerson TB, Dube WP, Wagner NL, Parrish DD (2011) City lights and urban air. *Nat Geosci* 4:730–731
- Stockli R (2013) NASA's Earth Observatory—cloud fraction imagery using data provided by the MODIS Atmospheric Science Team, NASA Goddard Space Flight Centre, MA. http://neo.sci.gsfc.nasa.gov/view.php?datasetId=MYDAL2_M_CLD_FR. Accessed November 2013
- Stone EL, Jones G, Harris S (2009) Street lighting disturbs commuting bats. *Curr Biol* 19:1123–1127
- Stone EL, Jones G, Harris S (2012) Conserving energy at a cost to biodiversity? Impacts of LED lighting on bats. *Glob Change Biol* 18:2458–2465
- Stutte GW (2009) Light-emitting diodes for manipulating the phytochrome apparatus. *HortScience* 44:231–234
- Sutton PC (2003) A scale adjusted measure of “urban sprawl” using nighttime satellite imagery. *Remote Sens Environ* 86:353–369
- Tanner J (1996) Seasonality and lunar periodicity in the reproduction of pocilloporid corals. *Coral Reefs* 15:59–66
- Telfer TC, Sincock JL, Byrd GV, Reed JR (1987) Attraction of Hawaiian seabirds to lights: conservation efforts and effects of moon phase. *Wildl Soc Bull* 15:406–413
- Threlfall CG, Law B, Banks PB (2013) The urban matrix and artificial light restricts the nightly ranging behaviour of Gould's long-eared bat (*Nyctophilus gouldi*). *Austral Ecol* 38:921–930
- UNESCO (2009) Starlight reserves and world heritage: scientific, cultural and environmental values. UNESCO, Paris
- Urbanski J, Mogi M, O'Donnell D, DeCotiis M, Toma T, Armbruster P (2012) Rapid adaptive evolution of photoperiodic response during invasion and range expansion across a climatic gradient. *Am Nat* 179:490–500
- US Department of energy (2012) Light at night: the latest science. http://apps1.eere.energy.gov/buildings/publications/pdfs/ssl/ssl_whitepaper_nov2010.pdf
- Van Tichelen P, Geerken T, Jansen B, Vanden Bosch M, Van Hoof V, Vanhooydonck L, Vercalsteren A (2007) Final report lot 9: Public street lighting. <http://www.eup4light.net/assets/pdffiles/Final/VITOEuPStreetLightingFinal.pdf>
- Vivien-Roels B, Pévet P (1993) Melatonin: presence and formation in invertebrates. *Experientia* 49:642–647
- Vollsmes AV, Eriksen AB, Otterholt E, Kvaal K, Oxaal U, Futsaether CM (2009) Visible foliar injury and infrared imaging show that daylength affects short-term recovery after ozone stress in *Trifolium subterraneum*. *J Exp Bot* 60:3677–3686
- Wahr JM (1988) The Earth's rotation. *Annu Rev Earth Planet Sci* 16:231–249
- Wells JW (1963) Coral growth and geochronometry. *Nature* 197:948–950
- Widder EA, Robison BH, Reisenbichler KR, Haddock SHD (2005) Using red light for in situ observations of deep-sea fishes. *Deep-Sea Res I* 52:2077–2085
- Wiltschko W, Munro U, Ford H, Wiltschko R (1993) Red light disrupts magnetic orientation of migratory birds. *Nature* 364:525–527
- World Resources Institute (2007) Available at: <http://earthtrends.wri.org/index.php>
- Yurk H, Trites AW (2000) Experimental attempts to reduce predation by harbor seals on out-migrating juvenile salmonids. *Trans Am Fish Soc* 129:1360–1366
- Zubaidat AE, Ben-Shlomo R, Haim A (2007) Thermoregulatory and endocrine responses to light pulses in short-day acclimated social voles (*Microtus socialis*). *Chronobiol Int* 24:269–288