## University of Warsaw

Faculty of Mathematics, Informatics and Mechanics

### **Adam Starak**

Student no. 361021

# Title in English

> Supervisor: dr Michał Pilipczuk Instytut Informatyki

Supe	rvisor'	s sta	tement
Dupe	1 4 12 01	o ota	$c_{CIIICII}$

Hereby I confirm that the presented thesis was prepared under my supervision and that it fulfils the requirements for the degree of Master of Computer Science.

Date Supervisor's signature

#### Author's statement

Hereby I declare that the presented thesis was prepared by me and none of its contents was obtained by means that are against the law.

The thesis has never before been a subject of any procedure of obtaining an academic degree.

Moreover, I declare that the present version of the thesis is identical to the attached electronic version.

Date Author's signature

#### Abstract

W pracy przedstawiono prototypową implementację blabalizatora różnicowego bazującą na teorii fetorów  $\sigma$ - $\rho$  profesora Fifaka. Wykorzystanie teorii Fifaka daje wreszcie możliwość efektywnego wykonania blabalizy numerycznej. Fakt ten stanowi przełom technologiczny, którego konsekwencje trudno z góry przewidzieć.

#### Keywords

parameterized algorithm

Thesis domain (Socrates-Erasmus subject area codes)

11.3 Informatyka

#### Subject classification

D. SoftwareD.127. BlabalgorithmsD.127.6. Numerical blabalysis

Tytuł pracy w języku polskim

Tytuł po polsku

# Contents

In	troduction	5
1.	Basic definitions	
	1.1. Structures	
	1.2. Parameterized complexity	7
	1.3. Graph decomposition	7
2.	Spanning Star Forest Problem	Ć
	2.1. Obtaining a solution	10
	2.2. Spanning Star Forest parameterized by the number of stars	10
Bi	ibliografia	11

# Introduction

Blabalizator różnicowy jest podstawowym narzędziem blabalii fetorycznej. Dlatego naukowcy z całego świata prześcigają się w próbach efektywnej implementacji. Opracowana przez prof. Fifaka teoria fetorów  $\sigma$ - $\rho$  otwiera w tej dziedzinie nowe możliwości. Wykorzystujemy je w niniejszej pracy.

# Chapter 1

# Basic definitions

#### 1.1. Structures

A simple graph G is a pair (V, E) where V denotes a set of vertices and E denotes a set of undirected edges. Let  $deg_G(v)$  denote a degree of vertex v in graph G. Let  $G \setminus \{v\}$  be the abbreviation for  $G' = (V(G) \setminus \{v\}, E(G) \setminus \{(u, v) : u \in V(G)\})$ . A tree T is a graph where two vertices are connected by excatly one path. A spanning tree T of a graph G is a graph which includes all of the vertices of G, with minimum possible number of edges. A star S is a tree of size at least 2 for which at most 1 vertex has a degree greater than 1.

### 1.2. Parameterized complexity

**Definition 1.1.** Parameterized problem

**Definition 1.2.** FPT algorithm

Definition 1.3. Kernel

**Definition 1.4.** Kernelization algorithm

## 1.3. Graph decomposition

**Definition 1.5.** Path decomposition and pathwidth

**Definition 1.6.** Tree decomposition and treewidth

**Definition 1.7.** Nice tree decomposition

## Chapter 2

# Spanning Star Forest Problem

For a given graph G, we say that S is a  $Spanning\ Star\ Forest$  if every connected component C is a star. In the  $Spanning\ Star\ Forest\ Problem$  given a graph G, the objective is to determine whether there exists a  $Spanning\ Star\ Forest$ .

It turns out that the problem formulated in such a way is relatively simple. Although, various parametrizations described in this paper make it more complex. The following lemma easily clarifies all the concerns.

**Lemma 2.1.** A graph G has a Spanning Star Forest if and only if it does not contain any isolated vertices.

*Proof.* If G has a Spanning Star Forest S, then trivially  $\forall_{v \in V(G)} \ 1 \leq deg_S(v) \leq deg_G(v)$ . Thus, none of the vertices is isolated.

For the opposite direction, we prove the lemma by induction on |V(G)|. Assume |V(G)| = 2. The statement trivially holds because a graph representing an edge is a correct Spanning Star Forest. Let |V(G)| > 2 and let v be an arbitrary vertex. From the inductive assumption, let S be a Spanning Star Forest of a graph  $G \setminus \{v\}$ , u be a vertex such that  $(u, v) \in E(G)$  and w be a vertex such that  $w \in N_S(u)$ . Consider the 3 following cases:

- 1. Let  $deg_S(u) > 1$ . Then,  $S' = (V(S) \cup \{v\}, E(S) \cup \{(u,v)\})$  is a correct solution for graph G.
- 2. Let  $deg_S(u) = deg_S(w) = 1$ . Then,  $S' = (V(S) \cup v, E(S) \cup (u, v))$  is a correct solution for graph G.
- 3. Let  $deg_S(w) > 1$ . The,  $S' = (V(S) \cup \{v\}, (E(S) \cup \{(u,v)\}) \setminus \{(u,w)\})$  is a correct solution for graph G.

Observe that in graph G there are no isolated vertices. Thus, one can always extend a solution inductively.

Application of Lemma 2.1 yields the following result for Spanning Star Forest Problem:

**Theorem 2.1.** Decision version of Spanning Star Forest Problem can be solved in linear time.

*Proof.* Given an input G = (V, E) we answer YES if  $\forall_{v \in V(G)} \deg_G(v) \neq 0$  and NO otherwise.

### 2.1. Obtaining a solution

In this section the focus will be set on obtaining an arbitrary solution for a given instance of the *Spanning Star Forest Problem*.

**Theorem 2.2.** A solution for a Spanning Star Forest Problem can be found in linear time.

### 2.2. Spanning Star Forest parameterized by the number of stars

In the Spanning Star Forest Problem parameterized by the number of stars, given a graph G and a natural number k, the objective is to determine whether there exists a Spanning Star Forest S such that the number of components is less than k.

It is natural to ask whether one can find a solution that minimizes the number of connected components. Even though the problem looks slightly different than the previous one, *Spanning Star Forest* parameterized by the number of stars is NP-Complete. The following theorem proves the statement:

**Theorem 2.3.** Spanning Star Forest Parameterized by the number of stars is NP-Complete.

**Lemma 2.2.** There exists a reduction from Spanning Star Forest parameterized by the number of stars to Dominating Set.

# **Bibliography**

- [Bea65] Juliusz Beaman, Morbidity of the Jolly function, Mathematica Absurdica, 117 (1965) 338-9.
- [Blar16] Elizjusz Blarbarucki, O pewnych aspektach pewnych aspektów, Astrolog Polski, Zeszyt 16, Warszawa 1916.
- [Fif00] Filigran Fifak, Gizbert Gryzogrzechotalski, O blabalii fetorycznej, Materiały Konferencji Euroblabal 2000.
- [Fif01] Filigran Fifak, O fetorach  $\sigma$ - $\rho$ , Acta Fetorica, 2001.
- [Głomb04] Gryzybór Głombaski, Parazytonikacja blabiczna fetorów nowa teoria wszystkiego, Warszawa 1904.
- [Hopp96] Claude Hopper, On some  $\Pi$ -hedral surfaces in quasi-quasi space, Omnius University Press, 1996.
- [Leuk00] Lechoslav Leukocyt, Oval mappings ab ovo, Materiały Białostockiej Konferencji Hodowców Drobiu, 2000.
- [Rozk93] Josip A. Rozkosza, *O pewnych własnościach pewnych funkcji*, Północnopomorski Dziennik Matematyczny 63491 (1993).
- [Spy59] Mrowclaw Spyrpt, A matrix is a matrix is a matrix, Mat. Zburp., 91 (1959) 28–35.
- [Sri64] Rajagopalachari Sriniswamiramanathan, Some expansions on the Flausgloten Theorem on locally congested lutches, J. Math. Soc., North Bombay, 13 (1964) 72–6.
- [Whi25] Alfred N. Whitehead, Bertrand Russell, *Principia Mathematica*, Cambridge University Press, 1925.
- [Zen69] Zenon Zenon, Użyteczne heurystyki w blabalizie, Młody Technik, nr 11, 1969.