

# Spanning Star Forest Problem

Adam Starak

May 15, 2019

# Star definition

## Definition


A **star** is a tree of size at least 2 for which at most one vertex has a degree greater than 1.

# Star definition

## Definition

A **star** is a tree of size at least 2 for which at most one vertex has a degree greater than 1.

## Examples





# Star definition

## Definition

A **star** is a tree of size at least 2 for which at most one vertex has a degree greater than 1.

## Examples






# Star definition

## Definition

A **star** is a tree of size at least 2 for which at most one vertex has a degree greater than 1.

## Examples




# Spanning Star Forest Problem


## Definition

Given a graph  $G$  decide whether it has a *Spanning Star Forest*.

# Example




# Example





# Example



# Spanning Star Forest Problem

## Lemma

$G$  contains a Spanning Star Forest if and only if it does not contain any isolated vertices.

# Spanning Star Forest Problem

## Lemma

$G$  contains a Spanning Star Forest if and only if it does not contain any isolated vertices.

**Forward implication.** Trivial, Every vertex belongs to a tree of size at least 2. Thus, it's degree is at least 1.

# Backward implication

Suppose  $G$  has no isolated vertices.

# Backward implication

Suppose  $G$  has no isolated vertices.

Induction by the number of vertices in  $G$ .

# Backward implication

Suppose  $G$  has no isolated vertices.

Induction by the number of vertices in  $G$ .

Let  $n = 2$ .


# Backward implication

Suppose  $G$  has no isolated vertices.

Induction by the number of vertices in  $G$ .

Let  $n = 2$ .

Graph is a correct Spanning Star Forest.



# Backward implication

Suppose  $G$  has no isolated vertices.

Induction by the number of vertices in  $G$ .

Inductive step.



# Backward implication

Suppose  $G$  has no isolated vertices.

Induction by the number of vertices in  $G$ .

Inductive step.

Assume the condition holds for all graphs of size at most  $n$ .

# Backward implication

Suppose  $G$  has no isolated vertices.

Induction by the number of vertices in  $G$ .

Inductive step.

Assume the condition holds for all graphs of size at most  $n$ .

Suppose there does not exist a vertex  $v$  such that  $G \setminus \{v\}$  has no isolated vertices.

# Backward implication

Suppose  $G$  has no isolated vertices.




Induction by the number of vertices in  $G$ .

Inductive step.

Assume the condition holds for all graphs of size at most  $n$ .

Suppose there does not exist a vertex  $v$  such that  $G \setminus \{v\}$  has no isolated vertices.

Graph  $G$  is a correct Spanning Star Forest.



# Backward implication

Suppose  $G$  has no isolated vertices.

Induction by the number of vertices in  $G$ .

Inductive step.

Assume the condition holds for all graphs of size at most  $n$ .

Suppose there exists a vertex  $v$  such that  $G \setminus \{v\}$  has no isolated vertices.

# Backward implication

Suppose  $G$  has no isolated vertices.

Induction by the number of vertices in  $G$ .

Inductive step.

Assume the condition holds for all graphs of size at most  $n$ .

Suppose there exists a vertex  $v$  such that  $G \setminus \{v\}$  has no isolated vertices.

Let  $S$  be a solution for a graph  $G \setminus \{v\}$ .

# Backward implication

Suppose  $G$  has no isolated vertices.

Induction by the number of vertices in  $G$ .

Inductive step.

Assume the condition holds for all graphs of size at most  $n$ .

Suppose there exists a vertex  $v$  such that  $G \setminus \{v\}$  has no isolated vertices.


Let  $S$  be a solution for a graph  $G \setminus \{v\}$ .

Now, let's try to add vertex  $v$  to the solution  $S$ .

# Backward implication

Suppose  $G$  has no isolated vertices.

Induction by the number of vertices in  $G$ .




Inductive step.

Assume the condition holds for all graphs of size at most  $n$ .

Suppose there exists a vertex  $v$  such that  $G \setminus \{v\}$  has no isolated vertices.

Let  $S$  be a solution for a graph  $G \setminus \{v\}$ .


Now, let's try to add vertex  $v$  to the solution  $S$ .



# Backward implication

Suppose  $G$  has no isolated vertices.

Induction by the number of vertices in  $G$ .




Inductive step.

Assume the condition holds for all graphs of size at most  $n$ .

Suppose there exists a vertex  $v$  such that  $G \setminus \{v\}$  has no isolated vertices.

Let  $S$  be a solution for a graph  $G \setminus \{v\}$ .

Now, let's try to add vertex  $v$  to the solution  $S$ .





# Backward implication

Suppose  $G$  has no isolated vertices.

Induction by the number of vertices in  $G$ .


Inductive step.

Assume the condition holds for all graphs of size at most  $n$ .

Suppose there exists a vertex  $v$  such that  $G \setminus \{v\}$  has no isolated vertices.

Let  $S$  be a solution for a graph  $G \setminus \{v\}$ .

Now, let's try to add vertex  $v$  to the solution  $S$ .



# Backward implication

Suppose  $G$  has no isolated vertices.

Induction by the number of vertices in  $G$ .


Inductive step.

Assume the condition holds for all graphs of size at most  $n$ .

Suppose there exists a vertex  $v$  such that  $G \setminus \{v\}$  has no isolated vertices.

Let  $S$  be a solution for a graph  $G \setminus \{v\}$ .

Now, let's try to add vertex  $v$  to the solution  $S$ .



# Backward implication

Suppose  $G$  has no isolated vertices.

Induction by the number of vertices in  $G$ .


Inductive step.

Assume the condition holds for all graphs of size at most  $n$ .

Suppose there exists a vertex  $v$  such that  $G \setminus \{v\}$  has no isolated vertices.

Let  $S$  be a solution for a graph  $G \setminus \{v\}$ .

Now, let's try to add vertex  $v$  to the solution  $S$ .



# Backward implication

Suppose  $G$  has no isolated vertices.

Induction by the number of vertices in  $G$ .


Inductive step.

Assume the condition holds for all graphs of size at most  $n$ .

Suppose there exists a vertex  $v$  such that  $G \setminus \{v\}$  has no isolated vertices.

Let  $S$  be a solution for a graph  $G \setminus \{v\}$ .

Now, let's try to add vertex  $v$  to the solution  $S$ .



# Parametrization by the number of stars

## Definition

Given a pair  $(G, k)$  find a Spanning Star Forest  $S$  such that  $S$  has at most  $k$  connected components.

# Parametrization by the number of stars

## Definition

Given a pair  $(G, k)$  find a Spanning Star Forest  $S$  such that  $S$  has at most  $k$  connected components.

## Dominating Set

Given a pair  $(G, k)$  find a set  $D \subseteq V(G)$  such that  $|D| \leq k$  and every node is either in  $D$  or adjacent to  $D$ .

# Parametrization by the number of stars

## Definition

Given a pair  $(G, k)$  find a Spanning Star Forest  $S$  such that  $S$  has at most  $k$  connected components.

## Dominating Set

Given a pair  $(G, k)$  find a set  $D \subseteq V(G)$  such that  $|D| \leq k$  and every node is either in  $D$  or adjacent to  $D$ .

The problem is NP-Complete

# NP-completeness - membership in NP

Trivial. Given  $S$  check if it is a Spanning Star Forest and if it has at most  $k$  connected components.



## Construction:

**Construction:** Let  $(G, k)$  be an instance of Dominating Set Problem.

**Construction:** Let  $(G, k)$  be an instance of Dominating Set Problem.

We create  $G'$  as follows: for every isolated vertex  $v$  introduce a vertex  $v'$  and an edge  $(v, v')$ .

**Construction:** Let  $(G, k)$  be an instance of Dominating Set Problem.

We create  $G'$  as follows: for every isolated vertex  $v$  introduce a vertex  $v'$  and an edge  $(v, v')$ .

## Lemma

*$(G, k)$  has a solution if and only if  $(G', k)$  has one.*

**Construction:** Let  $(G, k)$  be an instance of Dominating Set Problem.

We create  $G'$  as follows: for every isolated vertex  $v$  introduce a vertex  $v'$  and an edge  $(v, v')$ .

## Lemma

*$(G, k)$  has a solution if and only if  $(G', k)$  has one.*

**Backward implication.**

Set of centers represents a correct dominating set of  $G$ .

# NP-completeness: Forward implication

**Forward implication.**

# NP-completeness: Forward implication

## Forward implication.

Without a loss of generality, if there exists a solution for  $(G, k)$ , then there exists a solution of minimal size. Let  $D$  be such a solution.

# NP-completeness: Forward implication

## Forward implication.

Without a loss of generality, if there exists a solution for  $(G, k)$ , then there exists a solution of minimal size. Let  $D$  be such a solution.

We create solution  $S$  as follows: For every  $v \in V(G') \setminus D$  add an edge  $(v, u)$  to the solution where  $u \in D$ .



# NP-completeness: Forward implication

## Forward implication.

Without a loss of generality, if there exists a solution for  $(G, k)$ , then there exists a solution of minimal size. Let  $D$  be such a solution.

We create solution  $S$  as follows: For every  $v \in V(G') \setminus D$  add an edge  $(v, u)$  to the solution where  $u \in D$ .

Assume  $S$  contains an isolated vertex. Trivially,  $v \in D$ .

# NP-completeness: Forward implication

## Forward implication.

Without a loss of generality, if there exists a solution for  $(G, k)$ , then there exists a solution of minimal size. Let  $D$  be such a solution.

We create solution  $S$  as follows: For every  $v \in V(G') \setminus D$  add an edge  $(v, u)$  to the solution where  $u \in D$ .

Assume  $S$  contains an isolated vertex. Trivially,  $v \in D$ .




# NP-completeness: Forward implication

## Forward implication.

Without a loss of generality, if there exists a solution for  $(G, k)$ , then there exists a solution of minimal size. Let  $D$  be such a solution.

We create solution  $S$  as follows: For every  $v \in V(G') \setminus D$  add an edge  $(v, u)$  to the solution where  $u \in D$ .

Assume  $S$  contains an isolated vertex. Trivially,  $v \in D$ .




# NP-completeness: Forward implication

## Forward implication.

Without a loss of generality, if there exists a solution for  $(G, k)$ , then there exists a solution of minimal size. Let  $D$  be such a solution.

We create solution  $S$  as follows: For every  $v \in V(G') \setminus D$  add an edge  $(v, u)$  to the solution where  $u \in D$ .

Assume  $S$  contains an isolated vertex. Trivially,  $v \in D$ .




# NP-completeness: Forward implication

## Forward implication.

Without a loss of generality, if there exists a solution for  $(G, k)$ , then there exists a solution of minimal size. Let  $D$  be such a solution.

We create solution  $S$  as follows: For every  $v \in V(G') \setminus D$  add an edge  $(v, u)$  to the solution where  $u \in D$ .

Assume  $S$  contains an isolated vertex. Trivially,  $v \in D$ .



# Reduction from SSF to Dom-Set

## Construction

Given  $(G, k)$  transform the instance as following:


$$(G', k') = \begin{cases} (G, 0), & \text{if } G \text{ contains an isolated vertex.} \\ (G, k), & \text{otherwise.} \end{cases}$$

# Spanning Star Forest Extension Problem

## Definition

Given a graph  $G$  and a set of edges  $F \subseteq E(G)$ , find a Spanning Star Forest  $S$  such that  $F \subseteq E(S)$ .

## Example




# Instance normalization

- (R1) If  $G$  has an isolated vertex, it is a no instance.




# Instance normalization

- (R1) If  $G$  has an isolated vertex, it is a no instance.
- (R2) If  $G$  has a path of size 3 made from isolated edges, then it is a no instance



# Instance normalization


- (R1) If  $G$  has an isolated vertex, it is a no instance.
- (R2) If  $G$  has a path of size 3 made from isolated edges, then it is a no instance
- (R3) If  $G$  has a path of size 2 made from isolated edges, then remove all the vertices adjacent to the center.



# Instance normalization 2

$$G_P = \{v : u, v \notin V(F) \text{ and } \exists(u, v) \in E(G)\}.$$


$$G_{NP} = G \setminus G_P.$$



# Instance normalization 2

$$G_P = \{v : u, v \notin V(F) \text{ and } \exists(u, v) \in E(G)\}.$$

$$G_{NP} = G \setminus G_P.$$




# Instance normalization 2

$$G_P = \{v : u, v \notin V(F) \text{ and } \exists(u, v) \in E(G)\}.$$

$$G_{NP} = G \setminus G_P.$$

$G_P$  has always a SSF because it does not contain any isolated edges and for all  $v \in V(G_P)$   $\deg_{G_P}(v) \neq 0$ .



# Instance normalization 3

$$G_P = \{v : u, v \notin V(F) \text{ and } \exists(u, v) \in E(G)\}.$$

$$G_{NP} = G \setminus G_P.$$

## Lemma

*$(G, F)$  has a solution if and only if  $(G_{NP}, F)$  has one.*

# Instance normalization 3

$$G_P = \{v : u, v \notin V(F) \text{ and } \exists(u, v) \in E(G)\}.$$

$$G_{NP} = G \setminus G_P.$$

## Lemma

*$(G, F)$  has a solution if and only if  $(G_{NP}, F)$  has one.*

**Backward implication.**

# Instance normalization 3

$$G_P = \{v : u, v \notin V(F) \text{ and } \exists(u, v) \in E(G)\}.$$

$$G_{NP} = G \setminus G_P.$$

## Lemma

*$(G, F)$  has a solution if and only if  $(G_{NP}, F)$  has one.*

### Backward implication.

Let  $S$  be a solution for  $G_{NP}$ .



# Instance normalization 3

$$G_P = \{v : u, v \notin V(F) \text{ and } \exists(u, v) \in E(G)\}.$$

$$G_{NP} = G \setminus G_P.$$

## Lemma

*$(G, F)$  has a solution if and only if  $(G_{NP}, F)$  has one.*

### Backward implication.

Let  $S$  be a solution for  $G_{NP}$ .

Partition  $G$  into  $(G_P, G_{NP})$ .

# Instance normalization 3

$$G_P = \{v : u, v \notin V(F) \text{ and } \exists(u, v) \in E(G)\}.$$

$$G_{NP} = G \setminus G_P.$$

## Lemma

*$(G, F)$  has a solution if and only if  $(G_{NP}, F)$  has one.*

### Backward implication.

Let  $S$  be a solution for  $G_{NP}$ .

Partition  $G$  into  $(G_P, G_{NP})$ .

Find a solution  $S'$  for  $G_P$ .

# Instance normalization 3

$$G_P = \{v : u, v \notin V(F) \text{ and } \exists(u, v) \in E(G)\}.$$

$$G_{NP} = G \setminus G_P.$$

## Lemma

*$(G, F)$  has a solution if and only if  $(G_{NP}, F)$  has one.*

### Backward implication.

Let  $S$  be a solution for  $G_{NP}$ .

Partition  $G$  into  $(G_P, G_{NP})$ .

Find a solution  $S'$  for  $G_P$ .

$S \cup S'$  is a solution for  $G$ .

# Instance normalization 3


$$G_P = \{v : u, v \notin V(F) \text{ and } \exists(u, v) \in E(G)\}.$$

$$G_{NP} = G \setminus G_P.$$

## Lemma

*$(G, F)$  has a solution if and only if  $(G_{NP}, F)$  has one.*

**Forward implication.**



# SSFE is NP-complete

# SSFE is NP-complete

## Membership in NP

Trivial. Given a solution  $S$  check if all isolated edges are included and if it is a correct SSF.

# SSFE - Hardness

**Reduction from 3-SAT:** Given formula  $\phi$  do the following.




**Reduction from 3-SAT:** Given formula  $\phi$  do the following.

$$(x_1 \vee x_2 \vee \neg x_4) \wedge (\neg x_1 \vee x_3 \vee \neg x_5) \wedge (x_4 \vee x_5 \vee \neg x_1)$$

**Reduction from 3-SAT:** Given formula  $\phi$  do the following.

- 1 For each variable  $x_i$  introduce vertices  $v_{x_i}$ ,  $v_{\neg x_i}$ .


$$(x_1 \vee x_2 \vee \neg x_4) \wedge (\neg x_1 \vee x_3 \vee \neg x_5) \wedge (x_4 \vee x_5 \vee \neg x_1)$$



**Reduction from 3-SAT:** Given formula  $\phi$  do the following.

- 1 For each variable  $x_i$  introduce vertices  $v_{x_i}, v_{\neg x_i}$ .
- 2 For each  $v_{x_i}, v_{\neg x_i}$  introduce an isolated edge.


$$(x_1 \vee x_2 \vee \neg x_4) \wedge (\neg x_1 \vee x_3 \vee \neg x_5) \wedge (x_4 \vee x_5 \vee \neg x_1)$$



**Reduction from 3-SAT:** Given formula  $\phi$  do the following.

- 1 For each variable  $x_i$  introduce vertices  $v_{x_i}, v_{\neg x_i}$ .
- 2 For each  $v_{x_i}, v_{\neg x_i}$  introduce an isolated edge.
- 3 For each clause  $c_i$  introduce vertex  $v_{c_i}$ .


$$(x_1 \vee x_2 \vee \neg x_4) \wedge (\neg x_1 \vee x_3 \vee \neg x_5) \wedge (x_4 \vee x_5 \vee \neg x_1)$$



**Reduction from 3-SAT:** Given formula  $\phi$  do the following.

- 1 For each variable  $x_i$  introduce vertices  $v_{x_i}, v_{\neg x_i}$ .
- 2 For each  $v_{x_i}, v_{\neg x_i}$  introduce an isolated edge.
- 3 For each clause  $c_i$  introduce vertex  $v_{c_i}$ .
- 4 If literal  $l$  exists in clause  $c$  introduce an edge between them.

$$(x_1 \vee x_2 \vee \neg x_4) \wedge (\neg x_1 \vee x_3 \vee \neg x_5) \wedge (x_4 \vee x_5 \vee \neg x_1)$$



# SSFE reverse reduction

## Theorem

There exists a reduction from 3-SAT to SSFE.

# SSFE reverse reduction

## Theorem

There exists a reduction from 3-SAT to SSFE.

## Corollary

3-SAT and SSFE are equally hard



# Parametrizations

# Parametrizations

Number of isolated edges  $|F|$ .

# Parametrizations

Number of isolated edges  $|F|$ .

Number of vertices  $|V(G) \setminus V(F)|$ .

# Parametrizations

Number of isolated edges  $|F|$ .

Number of vertices  $|V(G) \setminus V(F)|$ .

Treewidth.

Number of isolated edges  $|F|$

# Number of isolated edges $|F|$

## Simple algorithm

Branch on every isolated edge and check whether chosen vertices form a Spanning Star forest.

# Number of isolated edges $|F|$

## Simple algorithm

Branch on every isolated edge and check whether chosen vertices form a Spanning Star forest.

Complexity:  $\mathcal{O}^*(2^{|F|})$ .

# Number of isolated edges $|F|$

## Simple algorithm

Branch on every isolated edge and check whether chosen vertices form a Spanning Star forest.

Complexity:  $\mathcal{O}^*(2^{|F|})$ .

## SETH

let  $\delta_q$  be the infimum of the set of constants  $c$  for which there exists an algorithm solving q-SAT in time  $\mathcal{O}^*(2^{cn})$ . Then:

$$\lim_{q \rightarrow \infty} \delta_q = 1$$



SSFE parameterized by  $|F|$  doesn't invoke kernel.

SSFE parameterized by  $|F|$  doesn't invoke kernel.

## Proving nonexistence of a kernel

Input:  $(x_1, k), (x_2, k), \dots, (x_t, k)$  where  $(x_i, k)$  - instance of a given problem  $Q$ .

# SSFE parameterized by $|F|$ doesn't invoke kernel.

## Proving nonexistence of a kernel

Input:  $(x_1, k), (x_2, k), \dots, (x_t, k)$  where  $(x_i, k)$  - instance of a given problem  $Q$ .

Question: Does there exist an instance  $(x', k)$  such that satisfies the following conditions:

SSFE parameterized by  $|F|$  doesn't invoke kernel.

## Proving nonexistence of a kernel

Input:  $(x_1, k), (x_2, k), \dots, (x_t, k)$  where  $(x_i, k)$  - instance of a given problem  $Q$ .

Question: Does there exist an instance  $(x', k)$  such that satisfies the following conditions:

$$(1) \ k' \leq \text{poly}(\max |x_i, k| + \log(t))$$

# SSFE parameterized by $|F|$ doesn't invoke kernel.

## Proving nonexistence of a kernel

Input:  $(x_1, k), (x_2, k), \dots, (x_t, k)$  where  $(x_i, k)$  - instance of a given problem  $Q$ .

Question: Does there exist an instance  $(x', k)$  such that satisfies the following conditions:

$$(1) \ k' \leq \text{poly}(\max |x_i, k| + \log(t))$$

$$(2) \ (x', k') \in Q \text{ if and only if } (x_i, k) \in Q \text{ for some } i \leq t.$$

# SSFE parameterized by $|F|$ doesn't invoke kernel.

## Proving nonexistence of a kernel

Input:  $(x_1, k), (x_2, k), \dots, (x_t, k)$  where  $(x_i, k)$  - instance of a given problem  $Q$ .

Question: Does there exist an instance  $(x', k)$  such that satisfies the following conditions:

- (1)  $k' \leq \text{poly}(\max |x_i, k| + \log(t))$
- (2)  $(x', k') \in Q$  if and only if  $(x_i, k) \in Q$  for some  $i \leq t$ .

For a more general idea, read chapter 15 from Parameterized Algorithms.

# SSFE parameterized by $|F|$ doesn't invoke kernel.

## Proving nonexistence of a kernel

Input:  $(x_1, k), (x_2, k), \dots, (x_t, k)$  where  $(x_i, k)$  - instance of a given problem  $Q$ .

Question: Does there exist an instance  $(x', k)$  such that satisfies the following conditions:

- (1)  $k' \leq \text{poly}(\max |x_i, k| + \log(t))$
- (2)  $(x', k') \in Q$  if and only if  $(x_i, k) \in Q$  for some  $i \leq t$ .

For a more general idea, read chapter 15 from Parameterized Algorithms.

## Example

Given instances  $(G_1, k), (G_2, k), \dots, (G_t, k)$  of  $k$ -Path problem return  $(\bigcup_{i=1}^t G_i, k)$

Proof on the board.



SSFE parameterized by  $|V(G)| \setminus |V(F)|$

# SSFE parameterized by $|V(G)| \setminus |V(F)|$

## Quadratic kernel

If there exists a vertex  $v \in V(G) \setminus V(F)$  such that  $\deg_G(v) > k$ , then remove  $v$ .

# SSFE parameterized by $|V(G)| \setminus |V(F)|$

## Quadratic kernel

If there exists a vertex  $v \in V(G) \setminus V(F)$  such that  $\deg_G(v) > k$ , then remove  $v$ .

**Proof:**

# SSFE parameterized by $|V(G)| \setminus |V(F)|$

## Quadratic kernel

If there exists a vertex  $v \in V(G) \setminus V(F)$  such that  $\deg_G(v) > k$ , then remove  $v$ .

### **Proof:**

Assume there exists a SSF  $S$  in  $G \setminus \{v\}$ .

# SSFE parameterized by $|V(G)| \setminus |V(F)|$

## Quadratic kernel

If there exists a vertex  $v \in V(G) \setminus V(F)$  such that  $\deg_G(v) > k$ , then remove  $v$ .

### **Proof:**

Assume there exists a SSF  $S$  in  $G \setminus \{v\}$ .

Then, we "set" at most  $k$  centers.

# SSFE parameterized by $|V(G)| \setminus |V(F)|$

## Quadratic kernel

If there exists a vertex  $v \in V(G) \setminus V(F)$  such that  $\deg_G(v) > k$ , then remove  $v$ .

### Proof:

Assume there exists a SSF  $S$  in  $G \setminus \{v\}$ .

Then, we "set" at most  $k$  centers.

In graph  $G$   $\deg_G(v) > k$ . So, there exists isolated edge  $(u, w)$  such that  $\deg_S(u) = \deg_S(w) = 1$  and either  $(v, u) \in E(G)$  or  $(v, w) \in E(G)$ .

# SSFE parameterized by $|V(G)| \setminus |V(F)|$

## Quadratic kernel

If there exists a vertex  $v \in V(G) \setminus V(F)$  such that  $\deg_G(v) > k$ , then remove  $v$ .

### Proof:

Assume there exists a SSF  $S$  in  $G \setminus \{v\}$ .

Then, we "set" at most  $k$  centers.

In graph  $G$   $\deg_G(v) > k$ . So, there exists isolated edge  $(u, w)$  such that  $\deg_S(u) = \deg_S(w) = 1$  and either  $(v, u) \in E(G)$  or  $(v, w) \in E(G)$ .

Number of edges:  $\leq 2k^2$

# SSFE parameterized by $|V(G)| \setminus |V(F)|$

## Quadratic kernel

If there exists a vertex  $v \in V(G) \setminus V(F)$  such that  $\deg_G(v) > k$ , then remove  $v$ .

### Proof:

Assume there exists a SSF  $S$  in  $G \setminus \{v\}$ .

Then, we "set" at most  $k$  centers.

In graph  $G$   $\deg_G(v) > k$ . So, there exists isolated edge  $(u, w)$  such that  $\deg_S(u) = \deg_S(w) = 1$  and either  $(v, u) \in E(G)$  or  $(v, w) \in E(G)$ .

Number of edges:  $\leq 2k^2$

Number of edges:  $\leq k + 2k^2$ .



# SSFE parameterized by $tw(G)$

**Dynamic table.**  $dp[t, f]$  where  $t$  is node from a tree decomposition and  $f : X_t \rightarrow \{true, false\}$  defined as follows:

$$f(v) = \begin{cases} true, & v \in V(F) \text{ and } v \text{ is a center.} \\ true, & v \notin V(F) \text{ and } v \text{ is a part of a Star Tree.} \\ false, & \text{otherwise.} \end{cases}$$

$$f_{v \rightarrow p}(u) = \begin{cases} p, & \text{if } u=v. \\ f(u), & \text{otherwise.} \end{cases}$$

# SSFE parameterized by $tw(G)$

**Leaf node.** An empty graph is a correct Spanning Star Forest.

$$dp[t, f] = true$$

# SSFE parameterized by $tw(G)$

**Introduce vertex**  $v \in V(F)$ .

# SSFE parameterized by $tw(G)$

**Introduce vertex**  $v \in V(F)$ .

We assign *false* if both vertices of an isolated vertex  $(u, v)$  have the same value.

# SSFE parameterized by $tw(G)$

**Introduce vertex**  $v \in V(F)$ .

We assign *false* if both vertices of an isolated vertex  $(u, v)$  have the same value.


$$dp[t, f_{v \rightarrow p}] = \begin{cases} \text{false}, & \text{if } u \in X_t \text{ and } f(u) = p. \\ dp[t', f], & \text{otherwise.} \end{cases}$$

# SSFE parameterized by $tw(G)$

**Introduce vertex**  $v \in V(F)$ .

We assign *false* if both vertices of an isolated vertex  $(u, v)$  have the same value.

$$dp[t, f_{v \rightarrow p}] = \begin{cases} \text{false}, & \text{if } u \in X_t \text{ and } f(u) = p. \\ dp[t', f], & \text{otherwise.} \end{cases}$$




# SSFE parameterized by $tw(G)$

**Introduce vertex**  $v \in V(F)$ .

We assign *false* if both vertices of an isolated vertex  $(u, v)$  have the same value.

$$dp[t, f_{v \rightarrow p}] = \begin{cases} \text{false}, & \text{if } u \in X_t \text{ and } f(u) = p. \\ dp[t', f], & \text{otherwise.} \end{cases}$$




# SSFE parameterized by $tw(G)$

**Introduce vertex**  $v \in V(F)$ .

We assign *false* if both vertices of an isolated vertex  $(u, v)$  have the same value.

$$dp[t, f_{v \rightarrow p}] = \begin{cases} \text{false}, & \text{if } u \in X_t \text{ and } f(u) = p. \\ dp[t', f], & \text{otherwise.} \end{cases}$$






# SSFE parameterized by $tw(G)$

**Introduce vertex**  $v \in V(F)$ .

We assign *false* if both vertices of an isolated vertex  $(u, v)$  have the same value.

$$dp[t, f_{v \rightarrow p}] = \begin{cases} \text{false}, & \text{if } u \in X_t \text{ and } f(u) = p. \\ dp[t', f], & \text{otherwise.} \end{cases}$$



# SSFE parameterized by $tw(G)$

**Introduce vertex**  $v \notin V(F)$ .

# SSFE parameterized by $tw(G)$

**Introduce vertex**  $v \notin V(F)$ .

Vertex is isolated. Thus, the only correct possibility is to assign 0.

# SSFE parameterized by $tw(G)$

**Introduce vertex**  $v \notin V(F)$ .

Vertex is isolated. Thus, the only correct possibility is to assign 0.

$$dp[t, f_{v \rightarrow p}] = \begin{cases} dp[t', f], & \text{if } p = 0. \\ \text{false}, & \text{otherwise.} \end{cases}$$

# SSFE parameterized by $tw(G)$

**Introduce edge**  $(u, v) \in F$

# SSFE parameterized by $tw(G)$

**Introduce edge**  $(u, v) \in F$

We can skip this part. We assigned proper values during vertex introduction.

# SSFE parameterized by $tw(G)$

**Forget**  $v \in V(F)$

# SSFE parameterized by $tw(G)$

**Forget**  $v \in V(F)$

The vertex is covered anyway. We get the maximal value.



# SSFE parameterized by $tw(G)$

**Forget**  $v \in V(F)$

The vertex is covered anyway. We get the maxial value.

$$dp[t, f|_v] = dp[t', f_{v \rightarrow 0}] \vee dp[t', f_{v \rightarrow 1}]$$

# SSFE parameterized by $tw(G)$

**Forget**  $v \notin V(F)$

# SSFE parameterized by $tw(G)$

**Forget**  $v \notin V(F)$

The vertex needs to be covered. So, we pass the value if and only if 1 is assigned to vertex  $v$ .

# SSFE parameterized by $tw(G)$

**Forget**  $v \notin V(F)$

The vertex needs to be covered. So, we pass the value if and only if 1 is assigned to vertex  $v$ .

$$dp[t, f|_v] = \begin{cases} dp[t', f], & \text{if } f(v)=1. \\ false, & \text{otherwise.} \end{cases}$$

# SSFE parameterized by $tw(G)$

**Introduce edge**  $(u, v) \notin F, v \notin V(F)$

# SSFE parameterized by $tw(G)$

**Introduce edge**  $(u, v) \notin F, v \notin V(F)$

We change the value of vertex  $v$  from 0 to 1 if and only if  $u$  is a center ( $f(u) = 1$ ):

# SSFE parameterized by $tw(G)$

**Introduce edge**  $(u, v) \notin F, v \notin V(F)$

We change the value of vertex  $v$  from 0 to 1 if and only if  $u$  is a center ( $f(u) = 1$ ):

$$dp[t, f_{v \rightarrow 1}] = \begin{cases} dp[t', f_{v \rightarrow 0}] \vee dp[t', f_{v \rightarrow 1}], & \text{if } f(u) = 1 \\ dp[t', f_{v \rightarrow 1}], & \text{otherwise.} \end{cases}$$
$$dp[t, f_{v \rightarrow 0}] = dp[t', f_{v \rightarrow 0}]$$


# SSFE parameterized by $tw(G)$

**Introduce edge**  $(u, v) \notin F, v \notin V(F)$


We change the value of vertex  $v$  from 0 to 1 if and only if  $u$  is a center ( $f(u) = 1$ ):

$$dp[t, f_{v \rightarrow 1}] = \begin{cases} dp[t', f_{v \rightarrow 0}] \vee dp[t', f_{v \rightarrow 1}], & \text{if } f(u) = 1 \\ dp[t', f_{v \rightarrow 1}], & \text{otherwise.} \end{cases}$$
$$dp[t, f_{v \rightarrow 0}] = dp[t', f_{v \rightarrow 0}]$$

**Before**



**After**






# SSFE parameterized by $tw(G)$

**Introduce edge**  $(u, v) \notin F, v \notin V(F)$


We change the value of vertex  $v$  from 0 to 1 if and only if  $u$  is a center ( $f(u) = 1$ ):

$$dp[t, f_{v \rightarrow 1}] = \begin{cases} dp[t', f_{v \rightarrow 0}] \vee dp[t', f_{v \rightarrow 1}], & \text{if } f(u) = 1 \\ dp[t', f_{v \rightarrow 1}], & \text{otherwise.} \end{cases}$$
$$dp[t, f_{v \rightarrow 0}] = dp[t', f_{v \rightarrow 0}]$$

**Before**



**After**




# SSFE parameterized by $tw(G)$

**Introduce edge**  $(u, v) \notin F, v \notin V(F)$


We change the value of vertex  $v$  from 0 to 1 if and only if  $u$  is a center ( $f(u) = 1$ ):

$$dp[t, f_{v \rightarrow 1}] = \begin{cases} dp[t', f_{v \rightarrow 0}] \vee dp[t', f_{v \rightarrow 1}], & \text{if } f(u) = 1 \\ dp[t', f_{v \rightarrow 1}], & \text{otherwise.} \end{cases}$$
$$dp[t, f_{v \rightarrow 0}] = dp[t', f_{v \rightarrow 0}]$$

**Before**



**After**




# SSFE parameterized by $tw(G)$

**Introduce edge**  $(u, v) \notin F, v \notin V(F)$


We change the value of vertex  $v$  from 0 to 1 if and only if  $u$  is a center ( $f(u) = 1$ ):

$$dp[t, f_{v \rightarrow 1}] = \begin{cases} dp[t', f_{v \rightarrow 0}] \vee dp[t', f_{v \rightarrow 1}], & \text{if } f(u) = 1 \\ dp[t', f_{v \rightarrow 1}], & \text{otherwise.} \end{cases}$$
$$dp[t, f_{v \rightarrow 0}] = dp[t', f_{v \rightarrow 0}]$$

**Before**



**After**





# SSFE parameterized by $tw(G)$

**Join nodes**  $t_1, t_2$



# SSFE parameterized by $tw(G)$

Join nodes  $t_1, t_2$



# SSFE parameterized by $tw(G)$


Join nodes  $t_1, t_2$




$$\forall v \in V(F) \cap X_t \quad f(v) = f_1(v) = f_2(v)$$

# SSFE parameterized by $tw(G)$


Join nodes  $t_1, t_2$   
 $(t_1)$



$(t_2)$




$(t)$




# SSFE parameterized by $tw(G)$


Join nodes  $t_1, t_2$   
 $(t_1)$



$(t_2)$




$(t)$






# SSFE parameterized by $tw(G)$


Join nodes  $t_1, t_2$   
 $(t_1)$



$(t_2)$



$(t)$



$$C_2 = \forall v((V(G) \setminus V(F)) \cap X_t) \ f(v) = f_1(v) \vee f_2(v)$$

# SSFE parameterized by $tw(G)$

# SSFE parameterized by $tw(G)$

$$dp[t, f] = \begin{cases} \bigvee_{f_1, f_2} dp[t_1, f_1] \wedge dp[t_2, f_2], & \text{if } C_1 \text{ and } C_2 \text{ holds.} \\ false, & \text{otherwise.} \end{cases}$$

## SSFE parameterized by $tw(G)$

$$dp[t, f] = \begin{cases} \bigvee_{f_1, f_2} dp[t_1, f_1] \wedge dp[t_2, f_2], & \text{if } C_1 \text{ and } C_2 \text{ holds.} \\ false, & \text{otherwise.} \end{cases}$$

There are  $2^{|X_t|}$  functions. Each can be calculated naively in time  $\mathcal{O}^*(2^{|X_t|})$ .

# SSFE parameterized by $tw(G)$

$$dp[t, f] = \begin{cases} \bigvee_{f_1, f_2} dp[t_1, f_1] \wedge dp[t_2, f_2], & \text{if } C_1 \text{ and } C_2 \text{ holds.} \\ false, & \text{otherwise.} \end{cases}$$

There are  $2^{|X_t|}$  functions. Each can be calculated naively in time  $\mathcal{O}^*(2^{|X_t|})$ . Applying cover product, we can calculate all of the functions  $f$  in time  $\mathcal{O}^*(2^{|X_t|})$

## SSFE parameterized by $tw(G)$

$$dp[t, f] = \begin{cases} \bigvee_{f_1, f_2} dp[t_1, f_1] \wedge dp[t_2, f_2], & \text{if } C_1 \text{ and } C_2 \text{ holds.} \\ false, & \text{otherwise.} \end{cases}$$

There are  $2^{|X_t|}$  functions. Each can be calculated naively in time  $\mathcal{O}^*(2^{|X_t|})$ . Applying cover product, we can calculate all of the functions  $f$  in time  $\mathcal{O}^*(2^{|X_t|})$ .

### SSFE parameterized by $tw(G)$

There exists an algorithm solving SSFE parameterized by treewidth in time  $\mathcal{O}^*(2^{tw(G)})$ .

## SSFE parameterized by $tw(G)$

$$dp[t, f] = \begin{cases} \bigvee_{f_1, f_2} dp[t_1, f_1] \wedge dp[t_2, f_2], & \text{if } C_1 \text{ and } C_2 \text{ holds.} \\ false, & \text{otherwise.} \end{cases}$$

There are  $2^{|X_t|}$  functions. Each can be calculated naively in time  $\mathcal{O}^*(2^{|X_t|})$ . Applying cover product, we can calculate all of the functions  $f$  in time  $\mathcal{O}^*(2^{|X_t|})$ .

## SSFE parameterized by $tw(G)$

There exists an algorithm solving SSFE parameterized by treewidth in time  $\mathcal{O}^*(2^{tw(G)})$ .

## Lower bound

Assuming SETH, there does not exist an algorithm solving SSFE parameterized by  $tw(G)$  in time  $\mathcal{O}^*((2 - \epsilon)^{tw(G)})$ .