

Spanning Star Forest Problem

Adam Starak

May 12, 2019

Star definition

Definition


A star is a tree of size at least 2 for which at most one vertex has a degree greater than 1.

Star definition

Definition

A star is a tree of size at least 2 for which at most one vertex has a degree greater than 1.

Examples





Star definition

Definition

A star is a tree of size at least 2 for which at most one vertex has a degree greater than 1.

Examples






Star definition

Definition

A star is a tree of size at least 2 for which at most one vertex has a degree greater than 1.

Examples




Spanning Star Forest Problem


Definition

Given a graph G decide whether it has a *Spanning Star Forest*.


Example



Example



Example



Spanning Star Forest Problem

Lemma

G contains a Spanning Star Forest if and only if it does not contain any isolated vertices.

Spanning Star Forest Problem

Lemma

G contains a Spanning Star Forest if and only if it does not contain any isolated vertices.

Forward implication. Trivial, Every vertex belongs to a tree of size at least 2. Thus, it's degree is at least 1.

Backward implication

Suppose G has no isolated vertices.

Backward implication

Suppose G has no isolated vertices.

Induction by the number of vertices in G .

Backward implication

Suppose G has no isolated vertices.

Induction by the number of vertices in G .

Let $n = 2$.


Backward implication

Suppose G has no isolated vertices.

Induction by the number of vertices in G .

Let $n = 2$.

Graph is a correct Spanning Star Forest.



Backward implication

Suppose G has no isolated vertices.

Induction by the number of vertices in G .

Inductive step.

Backward implication

Suppose G has no isolated vertices.

Induction by the number of vertices in G .

Inductive step.

Assume the condition holds for all graphs of size at most n .

Backward implication

Suppose G has no isolated vertices.

Induction by the number of vertices in G .

Inductive step.

Assume the condition holds for all graphs of size at most n .

Suppose there does not exist a vertex v such that $G \setminus \{v\}$ has no isolated vertices.

Backward implication

Suppose G has no isolated vertices.




Induction by the number of vertices in G .

Inductive step.

Assume the condition holds for all graphs of size at most n .

Suppose there does not exist a vertex v such that $G \setminus \{v\}$ has no isolated vertices.

Graph G is a correct Spanning Star Forest.



Backward implication

Suppose G has no isolated vertices.

Induction by the number of vertices in G .

Inductive step.

Assume the condition holds for all graphs of size at most n .

Suppose there exists a vertex v such that $G \setminus \{v\}$ has no isolated vertices.

Backward implication

Suppose G has no isolated vertices.

Induction by the number of vertices in G .

Inductive step.

Assume the condition holds for all graphs of size at most n .

Suppose there exists a vertex v such that $G \setminus \{v\}$ has no isolated vertices.

Let S be a solution for a graph $G \setminus \{v\}$.

Backward implication

Suppose G has no isolated vertices.

Induction by the number of vertices in G .

Inductive step.

Assume the condition holds for all graphs of size at most n .

Suppose there exists a vertex v such that $G \setminus \{v\}$ has no isolated vertices.


Let S be a solution for a graph $G \setminus \{v\}$.

Now, let's try to add vertex v to the solution S .

Backward implication

Suppose G has no isolated vertices.

Induction by the number of vertices in G .




Inductive step.

Assume the condition holds for all graphs of size at most n .

Suppose there exists a vertex v such that $G \setminus \{v\}$ has no isolated vertices.

Let S be a solution for a graph $G \setminus \{v\}$.


Now, let's try to add vertex v to the solution S .



Backward implication

Suppose G has no isolated vertices.

Induction by the number of vertices in G .




Inductive step.

Assume the condition holds for all graphs of size at most n .

Suppose there exists a vertex v such that $G \setminus \{v\}$ has no isolated vertices.

Let S be a solution for a graph $G \setminus \{v\}$.

Now, let's try to add vertex v to the solution S .



Backward implication

Suppose G has no isolated vertices.

Induction by the number of vertices in G .


Inductive step.

Assume the condition holds for all graphs of size at most n .

Suppose there exists a vertex v such that $G \setminus \{v\}$ has no isolated vertices.

Let S be a solution for a graph $G \setminus \{v\}$.

Now, let's try to add vertex v to the solution S .



Backward implication

Suppose G has no isolated vertices.

Induction by the number of vertices in G .


Inductive step.

Assume the condition holds for all graphs of size at most n .

Suppose there exists a vertex v such that $G \setminus \{v\}$ has no isolated vertices.

Let S be a solution for a graph $G \setminus \{v\}$.

Now, let's try to add vertex v to the solution S .



Backward implication

Suppose G has no isolated vertices.

Induction by the number of vertices in G .


Inductive step.

Assume the condition holds for all graphs of size at most n .

Suppose there exists a vertex v such that $G \setminus \{v\}$ has no isolated vertices.

Let S be a solution for a graph $G \setminus \{v\}$.

Now, let's try to add vertex v to the solution S .



Backward implication

Suppose G has no isolated vertices.

Induction by the number of vertices in G .


Inductive step.

Assume the condition holds for all graphs of size at most n .

Suppose there exists a vertex v such that $G \setminus \{v\}$ has no isolated vertices.

Let S be a solution for a graph $G \setminus \{v\}$.

Now, let's try to add vertex v to the solution S .



Parametrization by the number of stars

Definition

Given a pair (G, k) find a Spanning Star Forest S such that S has at most k connected components.

Parametrization by the number of stars

Definition

Given a pair (G, k) find a Spanning Star Forest S such that S has at most k connected components.

Dominating Set

Given a pair (G, k) find a set $D \subseteq V(G)$ such that $|D| \leq k$ and every node is either in D or adjacent to D .

Parametrization by the number of stars

Definition

Given a pair (G, k) find a Spanning Star Forest S such that S has at most k connected components.

Dominating Set

Given a pair (G, k) find a set $D \subseteq V(G)$ such that $|D| \leq k$ and every node is either in D or adjacent to D .

The problem is NP-Complete

NP-completeness - membership in NP

Trivial. Given S check if it is a Spanning Star Forest and if it has at most k connected components.

Construction:

Construction: Let (G, k) be an instance of Dominating Set Problem.

Construction: Let (G, k) be an instance of Dominating Set Problem.

We create G' as follows: for every isolated vertex v introduce a vertex v' and an edge (v, v') .

Construction: Let (G, k) be an instance of Dominating Set Problem.

We create G' as follows: for every isolated vertex v introduce a vertex v' and an edge (v, v') .

Lemma

(G, k) has a solution if and only if (G', k) has one.

Construction: Let (G, k) be an instance of Dominating Set Problem.

We create G' as follows: for every isolated vertex v introduce a vertex v' and an edge (v, v') .

Lemma

(G, k) has a solution if and only if (G', k) has one.

Backward implication.

Set of centers represents a correct dominating set of G .

NP-completeness: Forward implication

Forward implication.

NP-completeness: Forward implication

Forward implication.

Without a loss of generality, if there exists a solution for (G, k) , then there exists a solution of minimal size. Let D be such a solution.

NP-completeness: Forward implication

Forward implication.

Without a loss of generality, if there exists a solution for (G, k) , then there exists a solution of minimal size. Let D be such a solution.

We create solution S as follows: For every $v \in V(G') \setminus D$ add an edge (v, u) to the solution where $u \in D$.

NP-completeness: Forward implication

Forward implication.

Without a loss of generality, if there exists a solution for (G, k) , then there exists a solution of minimal size. Let D be such a solution.

We create solution S as follows: For every $v \in V(G') \setminus D$ add an edge (v, u) to the solution where $u \in D$.

Assume S contains an isolated vertex. Trivially, $v \in D$.

NP-completeness: Forward implication

Forward implication.

Without a loss of generality, if there exists a solution for (G, k) , then there exists a solution of minimal size. Let D be such a solution.

We create solution S as follows: For every $v \in V(G') \setminus D$ add an edge (v, u) to the solution where $u \in D$.

Assume S contains an isolated vertex. Trivially, $v \in D$.




NP-completeness: Forward implication

Forward implication.

Without a loss of generality, if there exists a solution for (G, k) , then there exists a solution of minimal size. Let D be such a solution.

We create solution S as follows: For every $v \in V(G') \setminus D$ add an edge (v, u) to the solution where $u \in D$.

Assume S contains an isolated vertex. Trivially, $v \in D$.




NP-completeness: Forward implication

Forward implication.

Without a loss of generality, if there exists a solution for (G, k) , then there exists a solution of minimal size. Let D be such a solution.

We create solution S as follows: For every $v \in V(G') \setminus D$ add an edge (v, u) to the solution where $u \in D$.

Assume S contains an isolated vertex. Trivially, $v \in D$.




NP-completeness: Forward implication

Forward implication.

Without a loss of generality, if there exists a solution for (G, k) , then there exists a solution of minimal size. Let D be such a solution.

We create solution S as follows: For every $v \in V(G') \setminus D$ add an edge (v, u) to the solution where $u \in D$.

Assume S contains an isolated vertex. Trivially, $v \in D$.



Reduction from SSF to Dom-Set

Construction

Given (G, k) transform the instance as following:


$$(G', k') = \begin{cases} (G, 0), & \text{if } G \text{ contains an isolated vertex.} \\ (G, k), & \text{otherwise.} \end{cases}$$

Spanning Star Forest Extension Problem

Definition

Given a graph G and a set of edges $F \subseteq E(G)$, find a Spanning Star Forest S such that $F \subseteq E(S)$.

Example




Instance normalization

- (R1) If G has an isolated vertex, it is a no instance.


Instance normalization

- (R1) If G has an isolated vertex, it is a no instance.
- (R2) If G has a path of size 3 made from isolated edges, then it is a no instance



Instance normalization


- (R1) If G has an isolated vertex, it is a no instance.
- (R2) If G has a path of size 3 made from isolated edges, then it is a no instance
- (R3) If G has a path of size 2 made from isolated edges, then remove all the vertices adjacent to the center.



Instance normalization 2

$$G_P = \{v : u, v \notin V(F) \text{ and } \exists(u, v) \in E(G)\}.$$


$$G_{NP} = G \setminus G_P.$$



Instance normalization 2

$$G_P = \{v : u, v \notin V(F) \text{ and } \exists(u, v) \in E(G)\}.$$

$$G_{NP} = G \setminus G_P.$$




Instance normalization 2

$$G_P = \{v : u, v \notin V(F) \text{ and } \exists(u, v) \in E(G)\}.$$

$$G_{NP} = G \setminus G_P.$$

G_P has always a SSF because it does not contain any isolated edges and for all $v \in V(G_P)$ $\deg_{G_P}(v) \neq 0$.



Instance normalization 3

$$G_P = \{v : u, v \notin V(F) \text{ and } \exists(u, v) \in E(G)\}.$$

$$G_{NP} = G \setminus G_P.$$

Lemma

(G, F) has a solution if and only if (G_{NP}, F) has one.

Instance normalization 3

$$G_P = \{v : u, v \notin V(F) \text{ and } \exists(u, v) \in E(G)\}.$$

$$G_{NP} = G \setminus G_P.$$

Lemma

(G, F) has a solution if and only if (G_{NP}, F) has one.

Backward implication.

Instance normalization 3

$$G_P = \{v : u, v \notin V(F) \text{ and } \exists(u, v) \in E(G)\}.$$

$$G_{NP} = G \setminus G_P.$$

Lemma

(G, F) has a solution if and only if (G_{NP}, F) has one.

Backward implication.

Let S be a solution for G_{NP} .

Instance normalization 3

$$G_P = \{v : u, v \notin V(F) \text{ and } \exists(u, v) \in E(G)\}.$$

$$G_{NP} = G \setminus G_P.$$

Lemma

(G, F) has a solution if and only if (G_{NP}, F) has one.

Backward implication.

Let S be a solution for G_{NP} .

Partition G into (G_P, G_{NP}) .

Instance normalization 3

$$G_P = \{v : u, v \notin V(F) \text{ and } \exists(u, v) \in E(G)\}.$$

$$G_{NP} = G \setminus G_P.$$

Lemma

(G, F) has a solution if and only if (G_{NP}, F) has one.

Backward implication.

Let S be a solution for G_{NP} .

Partition G into (G_P, G_{NP}) .

Find a solution S' for G_P .

Instance normalization 3

$$G_P = \{v : u, v \notin V(F) \text{ and } \exists(u, v) \in E(G)\}.$$

$$G_{NP} = G \setminus G_P.$$

Lemma

(G, F) has a solution if and only if (G_{NP}, F) has one.

Backward implication.

Let S be a solution for G_{NP} .

Partition G into (G_P, G_{NP}) .

Find a solution S' for G_P .

$S \cup S'$ is a solution for G .

Instance normalization 3


$$G_P = \{v : u, v \notin V(F) \text{ and } \exists(u, v) \in E(G)\}.$$

$$G_{NP} = G \setminus G_P.$$

Lemma

(G, F) has a solution if and only if (G_{NP}, F) has one.

Forward implication.



SSFE is NP-complete

SSFE is NP-complete

Membership in NP

Trivial. Given a solution S check if all isolated edges are included and if it is a correct SSF.

SSFE - Hardness

Reduction from 3-SAT: Given formula ϕ do the following.


Reduction from 3-SAT: Given formula ϕ do the following.

$$(x_1 \vee x_2 \vee \neg x_4) \wedge (\neg x_1 \vee x_3 \vee \neg x_5) \wedge (x_4 \vee x_5 \vee \neg x_1)$$

Reduction from 3-SAT: Given formula ϕ do the following.

- 1 For each variable x_i introduce vertices v_{x_i} , $v_{\neg x_i}$.


$$(x_1 \vee x_2 \vee \neg x_4) \wedge (\neg x_1 \vee x_3 \vee \neg x_5) \wedge (x_4 \vee x_5 \vee \neg x_1)$$



Reduction from 3-SAT: Given formula ϕ do the following.

- 1 For each variable x_i introduce vertices $v_{x_i}, v_{\neg x_i}$.
- 2 For each $v_{x_i}, v_{\neg x_i}$ introduce an isolated edge.


$$(x_1 \vee x_2 \vee \neg x_4) \wedge (\neg x_1 \vee x_3 \vee \neg x_5) \wedge (x_4 \vee x_5 \vee \neg x_1)$$



Reduction from 3-SAT: Given formula ϕ do the following.

- 1 For each variable x_i introduce vertices $v_{x_i}, v_{\neg x_i}$.
- 2 For each $v_{x_i}, v_{\neg x_i}$ introduce an isolated edge.
- 3 For each clause c_i introduce vertex v_{c_i} .


$$(x_1 \vee x_2 \vee \neg x_4) \wedge (\neg x_1 \vee x_3 \vee \neg x_5) \wedge (x_4 \vee x_5 \vee \neg x_1)$$



Reduction from 3-SAT: Given formula ϕ do the following.

- 1 For each variable x_i introduce vertices $v_{x_i}, v_{\neg x_i}$.
- 2 For each $v_{x_i}, v_{\neg x_i}$ introduce an isolated edge.
- 3 For each clause c_i introduce vertex v_{c_i} .
- 4 If literal l exists in clause c introduce an edge between them.

$$(x_1 \vee x_2 \vee \neg x_4) \wedge (\neg x_1 \vee x_3 \vee \neg x_5) \wedge (x_4 \vee x_5 \vee \neg x_1)$$



SSFE reverse reduction

Theorem

There exists a reduction from 3-SAT to SSFE.

SSFE reverse reduction

Theorem

There exists a reduction from 3-SAT to SSFE.

Corollary

3-SAT and SSFE are equally hard

Parametrizations

Parametrizations

Number of isolated edges $|F|$.

Parametrizations

Number of isolated edges $|F|$.

Number of vertices $|V(G) \setminus V(F)|$.

Parametrizations

Number of isolated edges $|F|$.

Number of vertices $|V(G) \setminus V(F)|$.

Treewidth.

Number of isolated edges $|F|$

Number of isolated edges $|F|$

Simple algorithm

Branch on every isolated edge and check whether chosen vertices form a Spanning Star forest.

Number of isolated edges $|F|$

Simple algorithm

Branch on every isolated edge and check whether chosen vertices form a Spanning Star forest.

Complexity: $\mathcal{O}^*(2^{|F|})$.

Number of isolated edges $|F|$

Simple algorithm

Branch on every isolated edge and check whether chosen vertices form a Spanning Star forest.

Complexity: $\mathcal{O}^*(2^{|F|})$.

SETH

let δ_q be the infimum of the set of constants c for which there exists an algorithm solving q-SAT in time $\mathcal{O}^*(2^{cn})$. Then:

$$\lim_{q \rightarrow \infty} \delta_q = 1$$

SSFE parameterized by $|F|$ doesn't invoke kernel.

SSFE parameterized by $|F|$ doesn't invoke kernel.

Proving nonexistence of a kernel

Input: $(x_1, k), (x_2, k), \dots, (x_t, k)$ where (x_i, k) - instance of a given problem Q .

SSFE parameterized by $|F|$ doesn't invoke kernel.

Proving nonexistence of a kernel

Input: $(x_1, k), (x_2, k), \dots, (x_t, k)$ where (x_i, k) - instance of a given problem Q .

Question: Does there exist an instance (x', k) such that satisfies the following conditions:

SSFE parameterized by $|F|$ doesn't invoke kernel.

Proving nonexistence of a kernel

Input: $(x_1, k), (x_2, k), \dots, (x_t, k)$ where (x_i, k) - instance of a given problem Q .

Question: Does there exist an instance (x', k) such that satisfies the following conditions:

$$(1) \ k' \leq \text{poly}(\max |x_i, k| + \log(t))$$

SSFE parameterized by $|F|$ doesn't invoke kernel.

Proving nonexistence of a kernel

Input: $(x_1, k), (x_2, k), \dots, (x_t, k)$ where (x_i, k) - instance of a given problem Q .

Question: Does there exist an instance (x', k) such that satisfies the following conditions:

(1) $k' \leq \text{poly}(\max |x_i, k| + \log(t))$

(2) $(x', k') \in Q$ if and only if $(x_i, k) \in Q$ for some $i \leq t$.

SSFE parameterized by $|F|$ doesn't invoke kernel.

Proving nonexistence of a kernel

Input: $(x_1, k), (x_2, k), \dots, (x_t, k)$ where (x_i, k) - instance of a given problem Q .

Question: Does there exist an instance (x', k) such that satisfies the following conditions:

$$(1) \ k' \leq \text{poly}(\max |x_i, k| + \log(t))$$

$$(2) \ (x', k') \in Q \text{ if and only if } (x_i, k) \in Q \text{ for some } i \leq t.$$

For a more general idea, read chapter 15 from Parameterized Algorithms.

SSFE parameterized by $|F|$ doesn't invoke kernel.

Proving nonexistence of a kernel

Input: $(x_1, k), (x_2, k), \dots, (x_t, k)$ where (x_i, k) - instance of a given problem Q .

Question: Does there exist an instance (x', k) such that satisfies the following conditions:

$$(1) \ k' \leq \text{poly}(\max |x_i, k| + \log(t))$$

$$(2) \ (x', k') \in Q \text{ if and only if } (x_i, k) \in Q \text{ for some } i \leq t.$$

For a more general idea, read chapter 15 from Parameterized Algorithms.

Example

Given instances $(G_1, k), (G_2, k), \dots, (G_t, k)$ of k -Path problem return $(\bigcup_{i=1}^t G_i, k)$

Proof on the board.

SSFE parameterized by $|V(G)| \setminus |V(F)|$

SSFE parameterized by $|V(G)| \setminus |V(F)|$

Quadratic kernel

If there exists a vertex $v \in V(G) \setminus V(F)$ such that $\deg_G(v) > k$, then remove v .

SSFE parameterized by $|V(G)| \setminus |V(F)|$

Quadratic kernel

If there exists a vertex $v \in V(G) \setminus V(F)$ such that $\deg_G(v) > k$, then remove v .

Proof:

SSFE parameterized by $|V(G)| \setminus |V(F)|$

Quadratic kernel

If there exists a vertex $v \in V(G) \setminus V(F)$ such that $\deg_G(v) > k$, then remove v .

Proof:

Assume there exists a SSF S in $G \setminus \{v\}$.

SSFE parameterized by $|V(G)| \setminus |V(F)|$

Quadratic kernel

If there exists a vertex $v \in V(G) \setminus V(F)$ such that $\deg_G(v) > k$, then remove v .

Proof:

Assume there exists a SSF S in $G \setminus \{v\}$.

Then, we "set" at most k centers.

SSFE parameterized by $|V(G)| \setminus |V(F)|$

Quadratic kernel

If there exists a vertex $v \in V(G) \setminus V(F)$ such that $\deg_G(v) > k$, then remove v .

Proof:

Assume there exists a SSF S in $G \setminus \{v\}$.

Then, we "set" at most k centers.

In graph G $\deg_G(v) > k$. So, there exists isolated edge (u, w) such that $\deg_S(u) = \deg_S(w) = 1$ and either $(v, u) \in E(G)$ or $(v, w) \in E(G)$.

SSFE parameterized by $|V(G)| \setminus |V(F)|$

Quadratic kernel

If there exists a vertex $v \in V(G) \setminus V(F)$ such that $\deg_G(v) > k$, then remove v .

Proof:

Assume there exists a SSF S in $G \setminus \{v\}$.

Then, we "set" at most k centers.

In graph G $\deg_G(v) > k$. So, there exists isolated edge (u, w) such that $\deg_S(u) = \deg_S(w) = 1$ and either $(v, u) \in E(G)$ or $(v, w) \in E(G)$.

Number of edges: $\leq 2k^2$

SSFE parameterized by $|V(G)| \setminus |V(F)|$

Quadratic kernel

If there exists a vertex $v \in V(G) \setminus V(F)$ such that $\deg_G(v) > k$, then remove v .

Proof:

Assume there exists a SSF S in $G \setminus \{v\}$.

Then, we "set" at most k centers.

In graph G $\deg_G(v) > k$. So, there exists isolated edge (u, w) such that $\deg_S(u) = \deg_S(w) = 1$ and either $(v, u) \in E(G)$ or $(v, w) \in E(G)$.

Number of edges: $\leq 2k^2$

Number of edges: $\leq k + 2k^2$.