

# Functions of Several Random Variables (Ch. 5.5)

Dason Kurkiewicz

Iowa State University

June 19, 2013

# Outline

## Functions of Several Random Variables

## Expectations and variances of linear combinations

## The Central Limit Theorem

## Approximating the Mean and Variance of a Function

Functions of  
Several Random  
Variables (Ch. 5.5)

Dason Kurkiewicz

Functions of  
Several Random  
Variables

Expectations and  
variances of linear  
combinations

The Central Limit  
Theorem

Approximating the  
Mean and Variance  
of a Function

# Functions of several random variables

Functions of  
Several Random  
Variables (Ch. 5.5)

Dason Kurkiewicz

Functions of  
Several Random  
Variables

Expectations and  
variances of linear  
combinations

The Central Limit  
Theorem

Approximating the  
Mean and Variance  
of a Function

- ▶ We often consider functions of random variables of the form:

$$U = g(X, Y, \dots, Z)$$

where  $X, Y, \dots, Z$  are random variables.

- ▶  $U$  is itself a random variable.

## Example: connecting steel parts

- Suppose that a steel plate with nominal thickness .15 in. is to rest in a groove of nominal width .155 in., machined on the surface of a steel block.

Relative Frequency Distribution of Plate Thicknesses

Plate Thickness (in.)	Relative Frequency
.148	.4
.149	.3
.150	.3

Relative Frequency Distribution of Slot Widths

Slot Width (in.)	Relative Frequency
.153	.2
.154	.2
.155	.4
.156	.2

- $X$  = plate thickness
- $Y$  = slot width
- $U = Y - X$ , the “wiggle room” of the plate

# The distributions of $X$ , $Y$ , and $U$

The Probability Function for the  
Clearance  $U = Y - X$

Marginal and Joint Probabilities for  $X$  and  $Y$

$y \setminus x$	.148	.149	.150	$f_Y(y)$
.156	.08	.06	.06	.2
.155	.16	.12	.12	.4
.154	.08	.06	.06	.2
.153	.08	.06	.06	.2
$f_X(x)$	.4	.3	.3	

$u$	$f(u)$
.003	.06
.004	.12 = .06 + .06
.005	.26 = .08 + .06 + .12
.006	.26 = .08 + .12 + .06
.007	.22 = .16 + .06
.008	.08

- Determining the distribution of  $U$  is difficult in the continuous case.

# Outline

Functions of  
Several Random  
Variables (Ch. 5.5)

Dason Kurkiewicz

Functions of  
Several Random  
Variables

Expectations and  
variances of linear  
combinations

The Central Limit  
Theorem

Approximating the  
Mean and Variance  
of a Function

Functions of Several Random Variables

Expectations and variances of linear combinations

The Central Limit Theorem

Approximating the Mean and Variance of a Function

# Expectations and variances of linear combinations

Functions of  
Several Random  
Variables (Ch. 5.5)

Dason Kurkiewicz

Functions of  
Several Random  
Variables

Expectations and  
variances of linear  
combinations

The Central Limit  
Theorem

Approximating the  
Mean and Variance  
of a Function

- ▶  $X_1, X_2, \dots, X_n$  are independent random variables and

$$Y = a_0 + a_1X_1 + a_2X_2 + \cdots + a_nX_n$$

then:

$$\begin{aligned} E(Y) &= E(a_0 + a_1X_1 + a_2X_2 + \cdots + a_nX_n) \\ &= a_0 + a_1E(X_1) + a_2E(X_2) + \cdots + a_nE(X_n) \end{aligned}$$

$$\begin{aligned} \text{Var}(Y) &= \text{Var}(a_0 + a_1X_1 + a_2X_2 + \cdots + a_nX_n) \\ &= a_1^2 \cdot \text{Var}(X_1) + a_2^2 \cdot \text{Var}(X_2) + \cdots + a_n^2 \cdot \text{Var}(X_n) \end{aligned}$$

# Your turn: linear combinations

- Say we have two independent random variables  $X$  and  $Y$  with  $E(X) = 3.3$ ,  $Var(X) = 1.91$ ,  $E(Y) = 25$ , and  $Var(Y) = 65$ .
- Find:

$$E(3 + 2X - 3Y)$$

$$E(-4X + 3Y)$$

$$E(-4X - 6Y)$$

$$Var(3 + 2X - 3Y)$$

$$Var(2X - 5Y)$$

$$Var(-4X - 6Y)$$



## Answers: linear combinations

Functions of  
Several Random  
Variables (Ch. 5.5)

Dason Kurkiewicz

Functions of  
Several Random  
Variables

Expectations and  
variances of linear  
combinations

The Central Limit  
Theorem

Approximating the  
Mean and Variance  
of a Function

$$\begin{aligned}E(3 + 2X - 3Y) &= 3 + 2E(X) - 3E(Y) \\&= 3 + 2 \cdot 3.3 - 3 \cdot 25 \\&= -65.4\end{aligned}$$

$$\begin{aligned}E(-4X + 3Y) &= -4E(X) + 3E(Y) \\&= -4 \cdot 3.3 + 3 \cdot 25 \\&= 61.8\end{aligned}$$

$$\begin{aligned}E(-4X - 6Y) &= -4 \cdot E(X) - 6 \cdot E(Y) \\&= -4 \cdot 3.3 - 6 \cdot 25 \\&= -163.2\end{aligned}$$

## Answers: linear combinations

Functions of  
Several Random  
Variables (Ch. 5.5)

Dason Kurkiewicz

Functions of  
Several Random  
Variables

Expectations and  
variances of linear  
combinations

The Central Limit  
Theorem

Approximating the  
Mean and Variance  
of a Function

$$\begin{aligned} \text{Var}(3 + 2X - 3Y) &= 2^2 \cdot \text{Var}(X) + (-3)^2 \text{Var}(Y) \\ &= 4 \cdot 1.91 + 9 \cdot 65 \\ &= 592.64 \end{aligned}$$

$$\begin{aligned} \text{Var}(2X - 5Y) &= 2^2 \cdot \text{Var}(X) + (-5)^2 \text{Var}(Y) \\ &= 4 \cdot 1.91 + 25 \cdot 65 \\ &= 1632.64 \end{aligned}$$

$$\begin{aligned} \text{Var}(-4X - 6Y) &= (-4)^2 \cdot \text{Var}(X) + (-6)^2 \text{Var}(Y) \\ &= 16 \cdot 1.91 + 36 \cdot 65 \\ &= 2370.56 \end{aligned}$$

# Your turn: more linear combinations

- ▶ Say  $X \sim \text{Binomial}(n = 10, p = 0.5)$  and  $Y \sim \text{Poisson}(\lambda = 3)$ .
- ▶ Calculate:

$$E(5 + 2X - 7Y)$$

$$\text{Var}(5 + 2X - 7Y)$$

## Answer: more linear combinations

- First, note that:

$$E(X) = np = 10 \cdot 0.5 = 5$$

$$E(Y) = \lambda = 3$$

$$\text{Var}(X) = np(1 - p) = 10(0.5)(1 - 0.5) = 2.5$$

$$\text{Var}(Y) = \lambda = 3$$

Now, we can calculate:

$$\begin{aligned} E(5 + 2X - 7Y) &= 5 + 2E(X) - 7E(Y) \\ &= 5 + 2 \cdot 5 - 7 \cdot 3 \\ &= -6 \end{aligned}$$

$$\begin{aligned} \text{Var}(5 + 2X - 7Y) &= 2^2 \cdot \text{Var}(X) + (-7)^2 \cdot \text{Var}(Y) \\ &= 4 \cdot 2.5 + 49 \cdot 3 \\ &= 157 \end{aligned}$$

# iid random variables.

- ▶ **Identically Distributed:** Random variables  $X_1, X_2, \dots, X_n$  are identically distributed if they have the same probability distribution.
- ▶ **“iid”:** Random variables  $X_1, X_2, \dots, X_n$  are iid if they are **I**ndependent and **I**dentically **D**istributed.

# Your turn: averages of iid random variables

- ▶  $X_1, X_2, \dots, X_n$  are iid with expectation  $\mu$  and variance  $\sigma^2$ .
- ▶ Derive:

$$E(\bar{X})$$

$$\text{Var}(\bar{X})$$

where:

$$\bar{X} = \frac{X_1 + X_2 + \dots + X_n}{n}$$

the mean of the  $X_i$ 's.

# Answers: averages of iid random variables

Functions of  
Several Random  
Variables (Ch. 5.5)

Dason Kurkiewicz

Functions of  
Several Random  
Variables

Expectations and  
variances of linear  
combinations

The Central Limit  
Theorem

Approximating the  
Mean and Variance  
of a Function

$$\begin{aligned} E(\bar{X}) &= E\left(\frac{X_1 + X_2 + \cdots + X_n}{n}\right) \\ &= \frac{1}{n}E(X_1) + \frac{1}{n}E(X_2) + \cdots + \frac{1}{n}E(X_n) \\ &= \underbrace{\frac{1}{n}\mu + \frac{1}{n}\mu + \cdots + \frac{1}{n}\mu}_{n \text{ times}} \\ &= n \cdot \frac{1}{n}\mu \\ &= \boxed{\mu} \end{aligned}$$

► Remember  $E(\bar{X}) = \mu$ : it's an important result.

# Answers: averages of iid random variables

Functions of  
Several Random  
Variables (Ch. 5.5)

Dason Kurkiewicz

Functions of  
Several Random  
Variables

Expectations and  
variances of linear  
combinations

The Central Limit  
Theorem

Approximating the  
Mean and Variance  
of a Function

$$\begin{aligned} \text{Var}(\bar{X}) &= \text{Var}\left(\frac{X_1 + X_2 + \cdots + X_n}{n}\right) \\ &= \left(\frac{1}{n}\right)^2 \text{Var}(X_1) + \left(\frac{1}{n}\right)^2 \text{Var}(X_2) + \cdots + \left(\frac{1}{n}\right)^2 \cdot \text{Var}(X_n) \\ &= \underbrace{\frac{1}{n^2}\sigma^2 + \frac{1}{n^2}\sigma^2 + \cdots + \frac{1}{n^2}\sigma^2}_{n \text{ times}} \\ &= n \cdot \frac{1}{n^2}\sigma^2 \\ &= \boxed{\frac{\sigma^2}{n}} \end{aligned}$$

- Remember  $\text{Var}(\bar{X}) = \frac{\sigma^2}{n}$ : it's another important result.



## Example: length of seeds

- ▶ A botanist has collected a sample of 10 seeds and measures the length of each.
- ▶ The seed lengths  $X_1, X_2, \dots, X_{10}$  are supposed to be iid with mean  $\mu = 5$  mm and variance  $\sigma^2 = 2$  mm<sup>2</sup>.

$$E(\bar{X}) = \mu = 5$$

$$\text{Var}(\bar{X}) = \sigma^2/n = 2/10 = 0.2$$

# Outline

Functions of Several Random Variables

Expectations and variances of linear combinations

The Central Limit Theorem

Approximating the Mean and Variance of a Function

Functions of  
Several Random  
Variables (Ch. 5.5)

Dason Kurkiewicz

Functions of  
Several Random  
Variables

Expectations and  
variances of linear  
combinations

The Central Limit  
Theorem

Approximating the  
Mean and Variance  
of a Function

# The Central Limit Theorem

Functions of  
Several Random  
Variables (Ch. 5.5)

Dason Kurkiewicz

Functions of  
Several Random  
Variables

Expectations and  
variances of linear  
combinations

The Central Limit  
Theorem

Approximating the  
Mean and Variance  
of a Function

- ▶ If  $X_1, X_2, \dots, X_n$  are *any* iid random variables with mean  $\mu$  and variance  $\sigma^2 < \infty$  then as  $n \rightarrow \infty$

$$\bar{X} \approx \text{Normal} \left( \mu, \frac{\sigma^2}{n} \right)$$

- ▶ The Central Limit Theorem (CLT) one of the most important and useful results in statistics.

## Example: tool serial numbers

- ▶  $W_1$  = last digit of the serial number observed next Monday at 9 AM
- ▶  $W_2$  = last digit of the serial number the Monday after at 9 AM
- ▶  $W_1$  and  $W_2$  are independent with pmf:



$$f(w) = \begin{cases} 0.1 & w = 0, 1, 2, \dots, 9 \\ 0 & \text{otherwise} \end{cases}$$

- ▶  $\bar{W} = \frac{1}{2}(W_1 + W_2)$  has the pmf:


The Probability Function for  $\bar{W}$  for  $n = 2$

$\bar{w}$	$f(\bar{w})$	$\bar{w}$	$f(\bar{w})$	$\bar{w}$	$f(\bar{w})$	$\bar{w}$	$f(\bar{w})$	$\bar{w}$	$f(\bar{w})$
0.0	.01	2.0	.05	4.0	.09	6.0	.07	8.0	.03
0.5	.02	2.5	.06	4.5	.10	6.5	.06	8.5	.02
1.0	.03	3.0	.07	5.0	.09	7.0	.05	9.0	.01
1.5	.04	3.5	.08	5.5	.08	7.5	.04		

## Example: tool serial numbers




- What if  $\bar{W} = \frac{1}{8}(W_1 + W_2 + \cdots + W_8)$ , the average of 8 days of initial serial numbers?



## Example: excess sale time

- ▶  $\bar{S}$  = sample mean excess sale time (over a 7.5 s threshold) for 100 stamp sales.
- ▶ Each individual excess sale time should have an  $\text{Exp}(\alpha = 16.5 \text{ s})$  distribution. That means:
  - ▶  $E(\bar{S}) = \alpha = 16.5 \text{ s}$
  - ▶  $SD(\bar{S}) = \sqrt{\text{Var}(\bar{S})} = \sqrt{\frac{\alpha^2}{100}} = 1.65 \text{ s}$
  - ▶ By the Central Limit Theorem,  $\bar{S} \approx N(16.5, 1.65^2)$
- ▶ We want to approximate  $P(\bar{S} > 17)$ .

The approximate probability distribution of  $\bar{S}$  is normal with mean 16.5 and standard deviation 1.65



## Example: excess sale time

$$\begin{aligned}P(\bar{S} > 17) &= P\left(\frac{\bar{S} - 16.5}{1.65} > \frac{17 - 16.5}{1.65}\right) \\&\approx P(Z > 0.303) \quad (Z \sim N(0, 1)) \\&= 1 - P(Z \leq 0.303) \\&= 1 - \Phi(0.303) \\&= 1 - 0.62 \quad \text{from the standard normal table} \\&= 0.38\end{aligned}$$

## Example: net weight of baby food jars

Functions of  
Several Random  
Variables (Ch. 5.5)

Dason Kurkiewicz

Functions of  
Several Random  
Variables

Expectations and  
variances of linear  
combinations

The Central Limit  
Theorem

Approximating the  
Mean and Variance  
of a Function

- ▶ Individual jar weights are iid with unknown mean  $\mu$  and standard deviation  $\sigma = 1.6$  g
- ▶  $\bar{V}$  = sample mean weight of  $n$  jars  $\approx N\left(\mu, \frac{1.6^2}{n}\right)$ .
- ▶ We want to find  $\mu$ . One way to hone in on  $\mu$  is to find  $n$  such that:

$$P(\mu - 0.3 < \bar{V} < \mu + 0.3) = 0.8$$

That way, our measured value of  $\bar{V}$  is likely to be close to  $\mu$ .



## Example: net weight of baby food jars

Functions of  
Several Random  
Variables (Ch. 5.5)

Dason Kurkiewicz

Functions of  
Several Random  
Variables

Expectations and  
variances of linear  
combinations

The Central Limit  
Theorem

Approximating the  
Mean and Variance  
of a Function

$$\begin{aligned} 0.8 &= P(\mu - 0.3 < \bar{V} < \mu + 0.3) \\ &= P\left(\frac{-0.3}{1.6/\sqrt{n}} < \frac{\bar{V} - \mu}{1.6/\sqrt{n}} < \frac{0.3}{1.6/\sqrt{n}}\right) \\ &\approx P(-0.19\sqrt{n} < Z < 0.19\sqrt{n}) \quad (\text{by CLT}) \\ &= 1 - 2\Phi(-0.19\sqrt{n}) \quad (\text{look at the } N(0,1) \text{ pdf}) \end{aligned}$$

$$\begin{aligned} \Phi^{-1}(0.1) &= -0.19\sqrt{n} \\ n &= \frac{\Phi^{-1}(0.1)^2}{(-0.19)^2} \\ &= \frac{(-1.28)^2}{(-0.19)^2} \quad (\text{standard normal table}) \\ &= 46.10 \end{aligned}$$

► Hence, we'll need a sample size of  $n = 47$ .

## Example: cars

- ▶ Suppose a bunch of cars pass through certain stretch of road. Whenever a car comes, you look at your watch and record the time.
- ▶ Let  $X_i$  be the time (in hours) between when the  $i$ 'th car comes and the  $(i+1)$ 'th car comes,  $i = 1, \dots, 44$ . Suppose you know:

$$X_1, X_2, \dots, X_{44} \sim \text{iid } f(x) = e^{-x} \quad x \geq 0$$

- ▶ Find the probability that the average time gap between cars exceeds 1.05 hours.

## Example: cars

$$\mu = E(X_1)$$

$$= \int_{-\infty}^{\infty} xf(x)dx$$

$$= \int_0^{\infty} xe^{-x}dx$$

$$= -e^{-x}(x+1)|_0^{\infty} \quad \text{integration by parts}$$

$$= 1$$

## Example: cars

$$\begin{aligned}E(X_1^2) &= \int_{-\infty}^{\infty} x^2 f(x) dx \\&= \int_0^{\infty} x^2 e^{-x} dx \\&= -e^{-x}(x^2 + 2x + 2)|_0^{\infty} \quad \text{integration by parts} \\&= 2 \\\sigma^2 &= \text{Var}(X_1) \\&= E(X_1^2) - E^2(X_1) \\&= 2 - 1^2 \\&= 1\end{aligned}$$

## Example: cars

$$\begin{aligned}\bar{X} &\sim \text{approx. } N(\mu, \sigma^2/n) \\ &= N(1, 1/44)\end{aligned}$$

Thus:


$$\frac{\bar{X} - 1}{\sqrt{1/44}} \sim N(0, 1)$$

## Example: cars


Now, we're ready to approximate:

$$\begin{aligned}P(\bar{X} > 1.05) &= P\left(\frac{\bar{X} - 1}{\sqrt{1/44}} > \frac{1.05 - 1}{\sqrt{1/44}}\right) \\&= P\left(\frac{\bar{X} - 1}{\sqrt{1/44}} > 0.332\right) \\&\approx P(Z > 0.332) \\&= 1 - P(Z \leq 0.332) \\&= 1 - \Phi(0.332) \\&= 1 - 0.630 = 0.370\end{aligned}$$

# Example: cars



# Example: cars





# Example: cars

Functions of  
Several Random  
Variables (Ch. 5.5)


Dason Kurkiewicz

Functions of  
Several Random  
Variables

Expectations and  
variances of linear  
combinations

The Central Limit  
Theorem

Approximating the  
Mean and Variance  
of a Function



# Outline

Functions of Several Random Variables

Expectations and variances of linear combinations

The Central Limit Theorem

Approximating the Mean and Variance of a Function

Functions of  
Several Random  
Variables (Ch. 5.5)

Dason Kurkiewicz

Functions of  
Several Random  
Variables

Expectations and  
variances of linear  
combinations

The Central Limit  
Theorem

Approximating the  
Mean and Variance  
of a Function

## Approximating $E(U)$ and $Var(U)$ when determining $f_U(u)$ is too hard

Functions of  
Several Random  
Variables (Ch. 5.5)

Dason Kurkiewicz

Functions of  
Several Random  
Variables

Expectations and  
variances of linear  
combinations

The Central Limit  
Theorem

Approximating the  
Mean and Variance  
of a Function


- ▶ If  $X, Y, \dots, Z$  are independent,  $g$  is well-behaved, and the variances  $Var(X), Var(Y), \dots, Var(Z)$  are small enough, then  $U = g(X, Y, \dots, Z)$  has:

$$E(U) \approx g(E(X), E(Y), \dots, E(Z))$$

$$Var(U) \approx \left(\frac{\partial g}{\partial x}\right)^2 Var(X) + \left(\frac{\partial g}{\partial y}\right)^2 Var(Y) + \dots + \left(\frac{\partial g}{\partial z}\right)^2 Var(Z)$$

- ▶ These formulas are often called the **propagation of error formulas**.

## Example: an electric circuit



- ▶  $R$  is the total resistance of the circuit.
- ▶  $R_1$ ,  $R_2$ , and  $R_3$  are the resistances of resistors 1, 2, and 3, respectively.
- ▶  $E(R_i) = 100$ ,  $\text{Var}(R_i) = 2$ ,  $i = 1, 2, 3$ .

$$R = g(R_1, R_2, R_3) = R_1 + \frac{R_2 R_3}{R_2 + R_3}$$

## Example: an electric circuit

Functions of  
Several Random  
Variables (Ch. 5.5)

Dason Kurkiewicz

Functions of  
Several Random  
Variables

Expectations and  
variances of linear  
combinations

The Central Limit  
Theorem

Approximating the  
Mean and Variance  
of a Function

$$E(R) \approx g(100, 100, 100) = 100 + \frac{(100)(100)}{100 + 100} = 150\Omega$$

$$\frac{\partial g}{\partial r_1} = 1$$

$$\frac{\partial g}{\partial r_2} = \frac{(r_2 + r_3)r_3 - r_2r_3}{(r_2 + r_3)^2} = \frac{r_3^2}{(r_2 + r_3)^2}$$

$$\frac{\partial g}{\partial r_3} = \frac{(r_2 + r_3)r_2 - r_2r_3}{(r_2 + r_3)^2} = \frac{r_2^2}{(r_2 + r_3)^2}$$

$$\begin{aligned}\text{Var}(R) &\approx (1)^2(2)^2 + \left(\frac{(100)^2}{(100 + 100)^2}\right)^2 (2)^2 + \left(\frac{(100)^2}{(100 + 100)^2}\right)^2 (2)^2 \\ &= 4.5\end{aligned}$$

$$\text{SD}(R)\sqrt{4.5} \approx 2.12\Omega$$