

Joint Distributions and Independence (Ch. 5.4)

Dason Kurkiewicz

Iowa State University

June 18, 2013

Outline

Joint Distributions
and Independence
(Ch. 5.4)

Dason Kurkiewicz

The Discrete Case

Joint Distributions
Marginal Distributions
Conditional
Distributions
Independence

The Continuous Case

The Discrete Case

Joint Distributions
Marginal Distributions
Conditional Distributions
Independence

The Continuous Case

Example: bearings

- ▶ Consider multiple random variables at the same time.
- ▶ Suppose you're manufacturing ring bearings (nominal inner diameter 1.00 in) on rods (nominal diameter 0.99 in). Let:
 - ▶ X = the inside diameter of the next ring bearing
 - ▶ Y = rod diameter where the ring is located
- ▶ We might want to know probabilities like

$$P(X < Y)$$

since if $X < Y$, the assembly cannot be made.

Example: bearings

- ▶ A **joint probability function** for discrete random variables X and Y is a nonnegative function $f(x, y)$ such that:

$$f(x, y) = P(X = x \text{ and } Y = y)$$

as a distribution, $f \geq 0$ and:

$$\sum_{x,y} f(x, y) = 1$$

- ▶ For the discrete case, it is useful to give $f(x, y)$ in a table.
- ▶ Example: suppose:
 - ▶ X = torque required to loosen bolt #3 in the next apparatus.
 - ▶ Y = torque for bolt #4.

where all torques are rounded to the nearest integer.

Example: torque (blank entries are 0)

$f(x, y)$ for the Bolt Torque Problem

$y \backslash x$	11	12	13	14	15	16	17	18	19	20
20								2/34	2/34	1/34
19							2/34			
18			1/34	1/34			1/34	1/34	1/34	
17					2/34	1/34	1/34	2/34		
16				1/34	2/34	2/34			2/34	
15	1/34	1/34			3/34					
14					1/34			2/34		
13					1/34					

- ▶ $P(X = 18 \text{ and } Y = 17) = \frac{2}{34}$
- ▶ $P(X = 14 \text{ and } Y = 19) = 0$

Your turn: torque

$f(x, y)$ for the Bolt Torque Problem

$y \backslash x$	11	12	13	14	15	16	17	18	19	20
20								2/34	2/34	1/34
19							2/34			
18			1/34	1/34			1/34	1/34	1/34	
17					2/34	1/34	1/34	2/34		
16				1/34	2/34	2/34			2/34	
15	1/34	1/34			3/34					
14					1/34			2/34		
13					1/34					

Calculate:

1. $P(X \geq Y)$
2. $P(|X - Y| \leq 1)$

Answers: torque

$y \backslash x$	11	12	13	14	15	16	17	18	19	20
20										*
19									*	*
18								*	*	*
17							*	*	*	*
16						*	*	*	*	*
15					*	*	*	*	*	*
14				*	*	*	*	*	*	*
13			*	*	*	*	*	*	*	*

Combinations of bolt 3
and bolt 4 torques with $x \geq y$

Answers: torque

$$\begin{aligned}P(X \geq Y) &= \sum_{x \geq y} f(x, y) \\&= f(20, 20) + f(20, 19) + f(20, 18) + \cdots + f(13, 13)\end{aligned}$$

Dropping all the $f(x, y)$ values that equal 0:

$$\begin{aligned}&= f(15, 13) + f(15, 14) + f(15, 15) + f(16, 16) \\&+ f(17, 17) + f(18, 14) + f(18, 17) + f(18, 18) \\&+ f(19, 16) + f(19, 18) + f(20, 20) \\&\frac{1}{34} + \frac{1}{34} + \frac{3}{34} + \frac{2}{34} + \cdots + \frac{1}{34} = \frac{17}{34}\end{aligned}$$

Answers: torque

$y \backslash x$	11	12	13	14	15	16	17	18	19	20
20									*	*
19								*	*	*
18							*	*	*	
17						*	*	*		
16					*	*	*			
15				*	*	*				
14			*	*	*					
13		*	*	*						

Combinations of bolt 3
and bolt 4 torques with $|x - y| \leq 1$

Answers: torque

$$\begin{aligned}P(X \geq Y) &= \sum_{x \geq y} f(x, y) \\&= f(13, 13) + f(14, 13) + f(14, 14) + \cdots + f(20, 20)\end{aligned}$$

Dropping all the $f(x, y)$ values that equal 0:

$$\begin{aligned}&= f(15, 14) + f(15, 15) + f(15, 16) + f(16, 16) \\&+ f(16, 17) + f(17, 17) + f(17, 18) + f(18, 17) \\&+ f(18, 18) + f(19, 18) + f(19, 20) + f(20, 20) \\&= \frac{18}{34}\end{aligned}$$

- ▶ The **marginal distributions** of X and Y , which have joint pmf $f(x, y)$, are:

$$f_X(x) = \sum_y f(x, y)$$

$$f_Y(y) = \sum_x f(x, y)$$

- ▶ $f_X(x)$ is just the ordinary, univariate pmf of X .

Your turn: torque

- Calculate the marginal pmfs of X and Y

$f(x, y)$ for the Bolt Torque Problem

$y \backslash x$	11	12	13	14	15	16	17	18	19	20
20								$2/34$	$2/34$	$1/34$
19							$2/34$			
18			$1/34$	$1/34$			$1/34$	$1/34$	$1/34$	
17					$2/34$	$1/34$	$1/34$	$2/34$		
16				$1/34$	$2/34$	$2/34$			$2/34$	
15	$1/34$	$1/34$			$3/34$					
14					$1/34$			$2/34$		
13					$1/34$					

Answers: torque

- ▶ Take the column sums to calculate f_X at each x .
- ▶ Take the row sums to calculate f_Y at each y .

x	$f_X(x)$	y	$f_Y(y)$
11	1/34	13	5/34
12	1/34	14	2/34
13	1/34	15	5/34
14	2/34	16	6/34
15	9/34	17	7/34
16	3/34	18	7/34
17	4/34	19	3/34
18	7/34	20	1/34
19	5/34		
20	1/34		

Answers: torque

- It is customary to write the marginal pmfs in the margins of the table of the joint pmf.

Joint and Marginal Probabilities for X and Y

$y \setminus x$	11	12	13	14	15	16	17	18	19	20	$f_Y(y)$
20								2/34	2/34	1/34	5/34
19							2/34				2/34
18			1/34	1/34			1/34	1/34	1/34		5/34
17					2/34	1/34	1/34	2/34			6/34
16				1/34	2/34	2/34			2/34		7/34
15	1/34	1/34			3/34						5/34
14					1/34			2/34			3/34
13					1/34						1/34
$f_X(x)$	1/34	1/34	1/34	2/34	9/34	3/34	4/34	7/34	5/34	1/34	

- ▶ The **conditional distribution** of Y given $X = x$ is a function, $f_{Y|X=x}$, given by:

$$f_{Y|X=x}(y) = \frac{f(x, y)}{f_X(x)}$$

- ▶ To make sense of conditional distributions, return to the torque example...

Example: torque

Joint and Marginal Probabilities for X and Y

y \ x	11	12	13	14	15	16	17	18	19	20	$f_Y(y)$
20								2/34	2/34	1/34	5/34
19							2/34	0			2/34
18			1/34	1/34			1/34	1/34	1/34		5/34
17					2/34	1/34	1/34	2/34			6/34
16				1/34	2/34	2/34		0	2/34		7/34
15	1/34	1/34			3/34			0			5/34
14					1/34			2/34			3/34
13					1/34			0			1/34
$f_X(x)$	1/34	1/34	1/34	2/34	9/34	3/34	4/34	7/34	5/34	1/34	

- ▶ For example, $f_{Y|X=18}(20) = \frac{2/34}{7/34} = 2/7$. That makes sense because:
 - ▶ Since $f_X(18) = 7/34$, we expect roughly 7 out of every 34 cases to have $X = 18$.
 - ▶ Since $f_{X,Y}(18, 20) = 2/34$, we expect roughly 2 of those 7 cases to also have $Y = 20$.

The Discrete Case

Joint Distributions
Marginal Distributions
Conditional Distributions
Independence

The Continuous Case

Example: torque

y	13	14	15	16	17	18	19	20
$f_{X,Y}(18, y)$	2/34	0	1/34	2/34	0	0	2/34	0
$f_{Y X=18}(y)$	2/7	0	1/7	2/7	0	0	2/7	0

- ▶ $\sum_{y=13}^{20} f_{X,Y}(18, y) = f_X(18) = 7/34$
- ▶ $\sum_{y=13}^{20} f_{Y|X=18}(y) = 1$
- ▶ The conditional distribution, $f_{Y|X=18}$ is the renormalized column of the joint distribution corresponding to $X = 18$.

Your turn: torque

Joint and Marginal Probabilities for X and Y

$y \backslash x$	11	12	13	14	15	16	17	18	19	20	$f_Y(y)$
20								2/34	2/34	1/34	5/34
19							2/34				2/34
18			1/34	1/34			1/34	1/34	1/34		5/34
17					2/34	1/34	1/34	2/34			6/34
16				1/34	2/34	2/34			2/34		7/34
15	1/34	1/34			3/34						5/34
14					1/34			2/34			3/34
13					1/34						1/34
$f_X(x)$	1/34	1/34	1/34	2/34	9/34	3/34	4/34	7/34	5/34	1/34	

► Calculate:

1. $f_{Y|X=15}(y)$
2. $f_{Y|X=20}(y)$
3. $f_{X|Y=18}(x)$

The Discrete Case

Joint Distributions
Marginal Distributions
Conditional Distributions
Independence

The Continuous Case

Answers: torque

1.

y	13	14	15	16	17	18	19	20
$f_{Y X=15}(y)$	1/9	1/9	3/9	2/9	2/9	0	0	0

2.

y	13	14	15	16	17	18	19	20
$f_{Y X=20}(y)$	0	0	0	0	0	0	0	1

3.

x	11	12	13	14	15	16	17	18	19	20
$f_{X Y=18}(x)$	0	0	1/5	1/5	0	0	1/5	1/5	1/5	0

Given a set of marginal distributions, there are many possible joint distributions.

- What do you notice about each of the following joint distributions?

Distribution 1

$y \backslash x$	1	2	3	
3	.4	0	0	.4
2	0	.4	0	.4
1	0	0	.2	.2
	.4	.4	.2	

Distribution 2

$y \backslash x$	1	2	3	
3	.16	.16	.08	.4
2	.16	.16	.08	.4
1	.08	.08	.04	.2
	.4	.4	.2	

Given a set of marginal distributions, there are many possible joint distributions.

- What do you notice about each of the following joint distributions?

Distribution 1

$y \backslash x$	1	2	3	
3	.4	0	0	.4
2	0	.4	0	.4
1	0	0	.2	.2
	.4	.4	.2	

Distribution 2

$y \backslash x$	1	2	3	
3	.16	.16	.08	.4
2	.16	.16	.08	.4
1	.08	.08	.04	.2
	.4	.4	.2	

1. Given $X = x$, you know what Y has to be (and vice versa).
2. Each $P(X = x, Y = y)$ is just $P(X = x) \cdot P(Y = y)$; i.e., X and Y have no influence on each other.

A look at distribution 2

$y \backslash x$	1	2	3	
3	.16	.16	.08	.4
2	.16	.16	.08	.4
1	.08	.08	.04	.2
	.4	.4	.2	

- ▶ Among just the cases when $X = 1$:
 - ▶ $Y = 3$ every 16 out of $(16 + 16 + 8) = 40$ times: i.e., with probability $\frac{16}{40} = 0.4$
 - ▶ Same with $Y = 2$
 - ▶ $Y = 1$ every 8 out of $(16 + 16 + 8) = 40$ times: i.e., with probability 0.2
- ▶ So pmf of Y given $X = 1$ is the same as the marginal pmf of Y .

Independence

- ▶ Discrete random variables X and Y are independent (written $X \perp\!\!\!\perp Y$) if for all x and y ,

$$P(Y = y \mid X = x) = P(Y = y)$$

where \mid means “given”.

- ▶ If $X \perp\!\!\!\perp Y$, then:

$$P(Y = y \text{ and } X = x) = P(X = x) \cdot P(Y = y)$$

$$f(x, y) = f_X(x) \cdot f_Y(y)$$

- ▶ If X and Y are not only independent but also have the same marginal distribution, then they are **independent and identically distributed**, abbreviated **iid**.

Outline

Joint Distributions
and Independence
(Ch. 5.4)

Dason Kurkiewicz

The Discrete Case

Joint Distributions

Marginal Distributions

Conditional Distributions

Independence

The Discrete Case

Joint Distributions

Marginal Distributions

Conditional
Distributions

Independence

The Continuous Case

The Continuous Case

- ▶ A **joint probability density function** (pdf) for two continuous random variables X and Y is a nonnegative function with:

$$\int \int f(x, y) dx dy = 1$$

$$P((X, Y) \in R) = \int \int_R f(x, y) dx dy$$

where R is some region of \mathbb{R}^2 .

Example: sales desk


- ▶ S = true excess time (over a 7.5 s threshold) required to complete the next sale
- ▶ R = excess time measured with a stopwatch

$$f(s, r) = \begin{cases} \frac{1}{16.5} e^{-s/16.5} \frac{1}{\sqrt{2\pi(0.25)}} e^{-(r-s)^2/2(0.25)} & s > 0 \\ 0 & \text{otherwise} \end{cases}$$

$f(s, r)$ is valid.

$$\begin{aligned}\int \int f(s, r) ds dr &= \int_0^\infty \int_{-\infty}^\infty \frac{1}{16.5\sqrt{2\pi(0.25)}} e^{-(s/16.5) - ((r-s)^2/2(0.25))} dr ds \\&= \int_0^\infty \frac{1}{1.65} e^{-s/16.5} \left\{ \int_{-\infty}^\infty \frac{1}{\sqrt{2\pi(0.25)}} e^{-(r-s)^2/2(0.25)} dr \right\} ds \\&= \int_0^\infty \frac{1}{16.5} e^{-s/16.5} ds \\&= 1\end{aligned}$$


A look at $f(s, r)$



Checking for measurement bias: $P(\text{measured excess time} > \text{actual excess time})$

$$\begin{aligned} P(R > S) &= \int \int_{r > s} f(s, r) ds \, dr \\ &= \int_0^\infty \int_s^\infty f(s, r) dr \, ds \\ &= \int_0^\infty \frac{1}{16.5} e^{-s/16.5} \left\{ \int_s^\infty \frac{1}{\sqrt{2\pi}(0.25)} e^{-(r-s)^2/2(0.25)} dr \right\} ds \\ &= \int_0^\infty \frac{1}{16.5} e^{-s/16.5} \left\{ \frac{1}{2} \right\} ds \\ &= \frac{1}{2} \end{aligned}$$


Checking for measurement bias: region of integration



Probability of taking too long

$$\begin{aligned}P(S > 20) &= \int \int_{s > 20} f(s, r) dr \, ds \\&= \int_{20}^{\infty} \int_{-\infty}^{\infty} f(s, r) dr \, ds \\&= \int_{20}^{\infty} \frac{1}{16.5} e^{-s/16.5} \left\{ \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi(0.25)}} e^{-(r-s)^2/s(0.25)} \right\} ds \\&= \int_{20}^{\infty} e^{-s/16.5} ds \\&= e^{-20/16.5} \\&\approx 0.30\end{aligned}$$

Probability of taking too long: region of integration



Continuous marginal and conditional distributions

Joint Distributions
and Independence
(Ch. 5.4)

Dason Kurkiewicz

The Discrete Case
Joint Distributions
Marginal Distributions
Conditional
Distributions
Independence

The Continuous
Case

- ▶ For continuous random variables X and Y , the **marginal distribution** of X is:

$$f_X(x) = \int_{-\infty}^{\infty} f(x, y) dy$$

- ▶ The **conditional distribution** of Y given $X = x$ is:

$$f_{Y|X=x}(y) = \frac{f(x, y)}{f_X(x)}$$