
Bayesian Statistics II — Hierarchical Models

ECON 8310: Business Forecasting — Lecture 11, Part 2

Department of Economics

University of Nebraska at Omaha

September 1, 2026

Lecture Outline

- 1 Which Series May Be Pooled
- 2 Partial Pooling
- 3 Fitting It, and the Trap
- 4 Reading the Result
- 5 Key Takeaways

Which Series May Be Pooled

Before you pool anything, ask what the groups have in common.

The Question

A concrete one, and you have met it before. **Does demand rise in weeks when SNAP benefits are disbursed, and by how much — store by store?** Homework 3's memo had you stock up for SNAP weeks store by store, which assumed those differences are real. Today we check.

But one question comes before the model. SNAP is a *food* assistance program. Measure the raw SNAP-week effect on log units in all thirty series, grouped by category:

Category	Mean effect	Spread across stores
FOODS (10 series)	+0.089	0.058
HOBBIES (10 series)	+0.006	0.010
HOUSEHOLD (10 series)	+0.011	0.014

These are not thirty versions of the same thing. FOODS responds; the other two barely move. Pooling all thirty would tell the model they are interchangeable.

Exchangeability: the Assumption Under Every Hierarchy

Partial pooling rests on one assumption, worth naming before we use it.

[
Exchangeable] Groups are **exchangeable** when, before seeing the data, you have no reason to expect one to differ from another in a particular direction. Swapping their labels would not change what you believe.

The ten FOODS series qualify: same category, different stores, and you could not say in advance which store's SNAP effect should be largest. FOODS against HOBBIES does **not** — you know before looking that a food-assistance program moves food.

The setup for today. We work with the **ten FOODS series**, and deliberately thin **five** of them down to **8 weeks**. Five series are data-rich, five are data-poor. That imbalance is the whole point.

The lab pools all thirty anyway, and measures what it costs.

Complete Pooling and No Pooling

Within that exchangeable set there are two obvious approaches, and each fails in a different direction.

	What it does	How it fails
Complete pooling	One SNAP effect for all ten series. <code>b = pm.Normal("b", 0, 1)</code>	Denies that stores differ at all. A genuinely different store is averaged away.
No pooling	A separate effect per series, estimated independently. <code>shape=10</code>	A series with 8 weeks gets an estimate built from 8 weeks. Mostly noise, reported as a finding.

You have used both already without naming them. Homework 3's random forest pooled all thirty series into one model with store dummies — close to complete pooling. Homework 1 fitted ARIMA to one series at a time — no pooling. The choice has always been made by convenience; partial pooling makes it a modeling decision and lets the data settle it.

Partial Pooling

Let each series have its own effect — drawn from a shared distribution the data also estimates.

The Hierarchy Is One Extra Line

Give each series its own effect b_j , but say where those effects come from:

$$b_j \sim \mathcal{N}(\mu_b, \sigma_b), \quad j = 1, \dots, 10$$

μ_b is the typical SNAP effect across FOODS, and σ_b is **how much stores genuinely differ**. Crucially, both are estimated from the data rather than assumed.

$\sigma_b \rightarrow 0$	The model concludes stores do not differ, and reproduces complete pooling.
σ_b large	The model concludes they differ a lot, and approaches no pooling.
In between	Each b_j is pulled toward μ_b — <i>how far</i> depending on how much data that series has.

That last row is the whole idea. A series with 277 weeks barely moves; one with 8 weeks moves a long way. Nobody chose those weights — they fall out of the model.

Why This Is the Right Answer, Not a Compromise

Partial pooling sounds like splitting the difference between two bad options. It is better than that, and the reason is worth stating.

When a series has 8 observations, its unpooled estimate is mostly sampling noise. The hierarchy's pull toward μ_b is not a fudge — it is the correct response to a noisy measurement. You already believe this in another form: it is why you would not conclude a coin is biased after three flips.

This is regularization again, for the sixth time. A hierarchical prior shrinks per-group estimates toward a common value exactly as Ridge shrinks coefficients toward zero — and here the shrinkage strength σ_b is *estimated* rather than tuned by cross-validation.

GAM smoothing penalty (L3) → tree pruning (L4) → `reg_lambda` (L5) → Ridge and LASSO (L6) → weight decay (L7) → hierarchical priors (today).

Fitting It, and the Trap

The obvious code does not work, and the reason is geometry rather than statistics.

The Model in PyMC

Non-centered hierarchical model

```
with pm.Model() as model:

    mu_b = pm.Normal("mu_b", 0, 0.5)      # typical SNAP effect
    sd_b = pm.HalfNormal("sd_b", 0.25)    # how much stores differ
    z     = pm.Normal("z", 0, 1, shape=J)  # non-centered
    b     = pm.Deterministic("b", mu_b + z * sd_b)

    a = pm.Normal("a", y.mean(), 1, shape=J) # per-series level
    s = pm.Exponential("s", 1.0)

    pm.Normal("obs", mu=a[idx] + b[idx]*snap, sigma=s, observed=y)
```

`idx` maps each row to its series, so `b[idx]` picks out the right effect for every observation. That indexing pattern is how every hierarchical model in PyMC is written.

Note what is *not* here. No decision about how much to pool, no weighting scheme, no threshold for “enough data.” Those all come out of estimating σ_b .

The Trap: Write It the Obvious Way and It Breaks

The natural way to write the hierarchy is `b = pm.Normal("b", mu_b, sd_b, shape=J)` — the *centered* form, which reads exactly like the equation. On our data it produces:

	Centered	Non-centered	Want
Divergences	127	1	0
$\hat{R}$	1.044	1.002	< 1.01
ESS (bulk)	413	1,728	> 400

Why: when σ_b is small, the b_j must crowd into a narrow band, and the posterior takes on a funnel shape — wide where σ_b is large, pinched where it is small. A sampler tuned for the wide part cannot navigate the neck, and reports divergences.

The alarms do not fail alike. ESS passes at 413; $\hat{R}$ fails at 1.044, the sort of near-miss people talk themselves out of. **127 divergences against a threshold of zero admits no such conversation.**

BMCP §4.6.1, Posterior Geometry Matters — the single most common failure in applied hierarchical modeling.

Reading the Result

Shrinkage, measured — and checked against data the model never saw.

Shrinkage, Measured

Fit both models. Compare each series' estimated SNAP effect before and after pooling.

Series group	Unpooled spread	After pooling	Mean move
Thin — 8 weeks (5 series)	0.108	0.010	0.120
Full — 277 weeks (5 series)	0.055	0.037	0.015

The data-poor series moved 8.2 times further toward the group mean than the data-rich ones. Nobody told the model which series were thin. It inferred that from how much each series had to say.

Read the two right-hand columns together. The thin series' apparent spread of 0.108 nearly vanishes, while the full series **keep most of theirs** ($0.055 \rightarrow 0.037$). The hierarchy did not flatten everything: it discarded spread that 8 weeks could not support and kept spread that 277 weeks could.

That is the claim to be skeptical of, so the next slide tests it.

Checked Against Data the Model Never Saw

We thinned those five series ourselves, so the weeks we removed are still there. Compare each estimate against the SNAP effect actually observed in the held-out weeks.

Thin series	Unpooled	Hierarchical	Held-out truth
CA_1_FOODS	0.286	0.097	0.087
CA_3_FOODS	0.105	0.082	0.102
TX_1_FOODS	0.052	0.075	0.082
TX_3_FOODS	0.219	0.091	0.072
WI_2_FOODS	0.338	0.103	0.207
RMSE against truth	0.126	0.049	—

Partial pooling was two and a half times more accurate, beating the unpooled fit on four series out of five. Those estimates were not just noisy — they were confidently wrong, and the confidence came from 8 weeks of data.

Borrowing strength is not a figure of speech: the thin series were estimated better because the other five existed.

What the Model Actually Concluded

The estimated hierarchy: $\mu_b = 0.077$ with a 94% HDI of $[0.016, 0.150]$, and $\sigma_b = 0.066$.

In words: a SNAP week lifts FOODS units by roughly **8%**, and stores genuinely differ around that by about **7 percentage points**. Both numbers are answers, and the second one is the one no point estimate would have given you.

It does not license the store-level policy the unpooled fit suggested. An analyst reading the unpooled numbers would report uplifts from 5% to 40% and stock accordingly. The held-out weeks say that range was mostly noise — but $\sigma_b = 0.066$ says it is not *all* noise either.

The defensible recommendation: plan on an 8% FOODS uplift everywhere, allow real but modest store variation around it, and do not act on a store's own estimate until that store has the weeks to support one.

Key Takeaways

What hierarchy buys, and what it costs.

Lecture 11 Part 2: Key Takeaways

1. **Ask what may be pooled before you pool it.** Groups must be **exchangeable**. FOODS and HOBBIES are not, and no amount of good sampling fixes a hierarchy built over the wrong set.
2. **Complete pooling** denies groups differ; **no pooling** lets a group with 8 observations report noise as a finding. You have used both without naming them.
3. **Partial pooling** gives each group its own parameter drawn from a shared distribution, and estimates how much groups genuinely differ.
4. **Shrinkage is automatic and proportionate.** Our thin series moved $8.1\times$ further toward the group mean than data-rich ones, with nobody specifying that — and on held-out weeks it was $2.6\times$ more accurate.
5. This is **regularization for the sixth time** — and the only version where the shrinkage strength is estimated rather than tuned.
6. **Write it non-centered.** The obvious form gave 127 divergences, the reparameterized form 1 — same model, different geometry.

Next: Bayesian linear regression — coefficients as distributions, and what you can ask them.

References I
