



The t Distribution



- Normal distribution

a symmetric continuous distribution, has two parameters, the *mean* and *std*

- Normal distribution

a symmetric continuous distribution, has two parameters, the ***mean*** and ***std***


- Standard Normal distribution

a Normal distribution with mean = 0 and std = 1




The ***Standard*** Normal Distribution

- It is symmetric about zero with a fixed standard deviation = 1




The t Distribution, aka the *Student's t* Distribution

- A symmetric distribution centered at zero




The t Distribution, aka the *Student's t* Distribution

- A symmetric distribution centered at zero
- Has a single parameter called the *degrees of freedom* or *df*



The t Distribution, aka the *Student's t* Distribution

- A symmetric distribution centered at zero
- Has a single parameter called the *degrees of freedom* or *df*



The t Distribution

- Does not have any standalone business application
- Used as an interim tool to calculate confidence intervals and hypothesis testing

The t Distribution


- Does not have any standalone business application
- Used as an interim tool to calculate confidence intervals and hypothesis testing
- Has an associated *probability density function*

The t Distribution

- Does not have any standalone business application
- Used as an interim tool to calculate confidence intervals and hypothesis testing
- Has an associated *probability density function*
- Associated Excel functions are the **=T.DIST** and **=T.INV**




The t Distribution, T.DIST function






The t Distribution, T.DIST function






The t Distribution, T.DIST function






The t Distribution, T.DIST function






The t Distribution, T.DIST function






The t Distribution, T.DIST function






The t Distribution, T.DIST function






The t Distribution, T.DIST function






The t Distribution, T.DIST function






The t Distribution, T.DIST function






The t Distribution, T.DIST function






The t Distribution, T.DIST function






The t Distribution, T.DIST function






The t Distribution, T.DIST function






The t Distribution, T.DIST function






The t Distribution, T.DIST function






The t Distribution, T.DIST function






The t Distribution, T.DIST function






The t Distribution, T.DIST function






The t Distribution, T.DIST function




The t Distribution, T.DIST function






The t Distribution, T.DIST function






The t Distribution, T.DIST function






The t Distribution, T.DIST function






The t Distribution, T.DIST function






The t Distribution, T.DIST function






The t Distribution, T.DIST function







The t Distribution, T.DIST function



The t Distribution, T.DIST function




The t Distribution, T.DIST function






The t Distribution, T.DIST function







The t Distribution, T.DIST function




The t Distribution, T.DIST function




The t Distribution, T.DIST function




The t Distribution, T.DIST function




The t Distribution, T.DIST function




The t Distribution, T.DIST function




The t Distribution, T.DIST function




The t Distribution, T.DIST function



The t Distribution, T.DIST function




The t Distribution, T.DIST function





The t Distribution, T.DIST function





The t Distribution, T.DIST function

Syntax:

`=T.DIST(x, df, TRUE)`

The t Distribution, T.DIST function

Syntax:

`=T.DIST(x, df, TRUE)`

`=T.DIST(1.3, 21, TRUE)`

The t Distribution, T.DIST function


Syntax:

`=T.DIST(x, df, TRUE)`

`=T.DIST(1.3, 21, TRUE)`




The t Distribution, T.DIST function






The t Distribution, T.DIST function



The t Distribution, T.DIST function



The t Distribution, T.DIST function

Syntax:

=T.DIST(x, df, TRUE)

We will **never** use the T.DIST function
with the “FALSE” option



RICE
JONES GRADUATE
SCHOOL OF BUSINESS

The t Distribution, *degrees of freedom*



The t Distribution, *degrees of freedom*

- It is the parameter of the t-distribution



The t Distribution, *degrees of freedom*

- It is the parameter of the t-distribution
- A mathematical constraint imposed by the data

The t Distribution, *degrees of freedom*

- It is the parameter of the t-distribution
- A mathematical constraint imposed by the data
 - linked to size of data being used
 - larger set of data would have greater degrees of freedom
 - as degrees of freedom increase the t-distribution becomes closer and closer to the standard normal distribution