

STAT 145 Course Pack

UWL Department of Mathematics & Statistics

1.1 Introduction to Data

Scientists seek to answer questions using rigorous methods and careful observations. These observations — collected from the likes of field notes, surveys, and experiments — form the backbone of a statistical investigation and are called **data**. Statistics is the study of how best to collect, analyze, and draw conclusions from data. In this first chapter, we focus on both the properties of data and on the collection of data. We will introduce the building blocks of statistical thinking: observations, variables, data frames, and the types of questions that statistics can help us answer.

Key Concepts

- Recognize that a data set consists of cases and variables
- Identify variables as either categorical or quantitative
- Determine explanatory and response variables where appropriate
- Determine whether a study is observational or experimental

Why we collect data

All statistical analyses start with a question or an idea. This question or idea is posed about a *population*, in particular a population *parameter*. The main purpose of collecting data is to learn about this unknown population parameter. We often cannot measure data for every individual in a population because it would cost too much money or take too long, so a *sample*, or subset of the population, is collected instead. Using the information in the sample, a *statistic* is calculated and used to estimate the corresponding population parameter. This process of using a sample statistic to learn about a population parameter is called *statistical inference*. One of the main goals of this class is to introduce you to the procedures for making statistical inferences in a variety of settings.


A *population* is a well-defined collection of *all* individuals or objects of interest.

A *parameter* is a number that describes a population and is almost always unknown.

A *sample* is a subset of the population on which we measure data.

A *statistic* is a number that describes a sample and is almost always known.

Inference is the process of drawing conclusions about a *population parameter* from a *sample statistic*.



Data Basics

After data is gathered, it needs a place to be stored. In statistics, this is called a *data set*, or *data frame*. Datasets can be as simple as an Excel spreadsheet or as complicated as a major database. It all depends on the complexity of the question.

Structure: Cases and variables

In general, when data is stored in a table, each row of the table represents an *observation*, or a *case*, in the dataset. Each column of the table represents a *variable* in the dataset.

Note: Many datasets use the first column as an identifier. This would not be considered a variable.

A *data set*, or *data frame* is used to store the set of measurements taken on individual units.

Observations, or *Cases*, are the subjects/objects that we obtain information about. These are most generally known as *experimental units*.

A *Variable* is a characteristic that is measured and recorded for each case.

Class Example 1.1.1 Distinguishing between cases and variables

Let's get to know your classmates. Find two people near you and ask them the following questions:

- What is your first name?
- What is your major?
- How tall are you?
- How many credit hours are you taking this semester?
- How often do you drink caffeine (never, once a week, few times a week, everyday)?

Name	Major	Height	Credit Hours	Caffeine

How many cases are there?

How many variables are there?

Types of Variables

Before we can know how to answer a question, we need to think about the kinds of variables we have recorded. Variables that split cases into groups, called *levels*, are called *categorical*, or *qualitative*, variables. Variables that are numerical quantities are called *quantitative* variables.

Categorical variables can further be subdivided:

- Nominal variables* have levels with no natural ordering. The variable *major* from Example 1.1 is nominal, there is no reason to rank one major over another.
- Ordinal variables* have levels with a meaningful order. The variable *caffeine* has a natural ordering.

Quantitative variables can further be subdivided:

- Continuous variables* can take any value within a range, including decimals. The variable *height* is continuous.
- Discrete variables* can only take counting values (typically whole numbers: 0, 1, 2, ...). The variable *credit hours* is discrete.

Note: In most of this course, we will treat categorical variables as nominal (un-

Categorical, or *Qualitative* variables divide the cases into groups, placing each case into exactly one of two or more non-numerical, non-overlapping categories.

Quantitative variables take on numerical values for each case, often by measuring or counting. Numerical operations like adding and averaging make sense for quantitative variables.

ordered) and quantitative variables as continuous. The distinction between nominal and ordinal, and between discrete and continuous becomes more important in advanced courses when special methods are used to take advantage of the ordering.

The ability to identify whether a variable is categorical or quantitative is important because different types of variables are summarized in different ways.

Class Example 1.1.2 *Computer ownership at UWL*

UWL would like to estimate the proportion of students who own a computer. The university asks 500 students.

- Identify the cases and variable(s) recorded.
- What is the sample?
- What is the population?
- Is the variable categorical or quantitative?
- Identify an unmeasured quantitative variable of interest.
- Is the result of this study a statistic or parameter?

Explanatory and Response Variables

In this class, we will discuss two ways in which data are used to answer questions. The first is looking only at a single variable, and the second is to determine if there is a relationship between two variables. When two variables show some connection or pattern with one another, they are called *associated* variables (or related variables). If two variables are not associated — there is no evident relationship between them — they are said to be *independent*.

In studies that look for an association, many times we want to know if one variable explains the behavior of another variable. In this kind of study, the *explanatory* variable would explain the behavior in the *response* variable.

Associated variables show some connection or pattern with one another.

Independent variables have no evident relationship between them.

An *explanatory* variable is one that helps us understand or predict the value of another variable.

The *response* variable is the variable that we are trying to understand.

Class Example 1.1.3 *Explanatory and response variables*

For the following research questions, identify the explanatory variable and the response variable.

- Does eating yogurt cause people to lose weight?

- Does meditation help reduce stress?

- Is hyperactivity in children affected by sugar consumption?

Class Example 1.1.4 *2023 Hollywood Movies*

Here is a small part of a dataset that includes information on all 136 movies to come out of Hollywood in 2023. “Rating” is audience rating on a 100-point scale, “Budget” is budget in millions of dollars, and “Opening” is opening weekend gross, in millions of dollars.

Title	Lead Studio	Rating	Genre	Budget	Opening
Oppenheimer	Atlas	93	Biography	\$100	\$82
Barbie	Warner Bros	88	Comedy	\$145	\$162
Spider-Man: Across the Spider-Verse	Sony	95	Action	\$100	\$120.5
Guardians of the Galaxy Vol. 3	Disney	82	Sci-fi	\$250	\$118
The Holdovers	Focus	97	Drama	\$13	\$2.7
Killers of the Flower Moon	Paramount	93	Drama	\$200	\$23

a) What are the cases in this dataset?

b) What are the variables in this dataset? For each variable, identify whether it is categorical or quantitative.

c) Name a question we might ask about this dataset that is about

- a single variable
- a relationship between two of the variables

Observational Studies and Experiments

There are two primary types of data collection: experiments and observational studies. Understanding the difference is critical, because the type of study determines what conclusions we can draw.

If the researcher is only observing what happened, the study is an *observational study*. In statistics, the word observational does not mean that the researcher is watching the participant do every part of the study. Instead, observational refers to the fact that the researcher takes a dataset and makes observations about the data in the set. If the researcher is actively involved with assigning different values of the explanatory variable to the participants, then the study is an *experiment*. In general, observational studies can provide evidence of a naturally occurring *association* between variables, but they cannot by themselves show a *causal* connection. This is because in observational studies, researchers cannot control for all the other variables that might be influencing the outcome.

Warning: Association does not imply causation.

In an *experiment* the researcher actively controls one or more of the explanatory variables.

In an *observational study* the researcher simply observes the values as they naturally exist.

Two variables are *associated* if values of one variable tend to be related to the values of the other variable.

Two variables are *causally associated* if changing the value of one variable influences the value of the other variable.

Class Example 1.1.5 Experimental or Observational Studies

Determine whether each of the following is an experiment or an observational study:

- a) A pharmaceutical company randomly assigns 200 patients to receive either a new drug or a placebo and measures their cholesterol levels after 6 months.

- b) A university surveys its graduates to determine whether those who participated in study-abroad programs earn higher salaries five years after graduation.
- c) A teacher assigns students in her morning class to use a new learning app and students in her afternoon class to study using traditional methods, then compares exam scores.

Summary

This chapter introduced you to the world of data. Here are the key ideas:

- *Data* are observations collected about the world around us. *Statistics* is the science of collecting, analyzing, and drawing conclusions from data.
- A *data set*, or *data frame*, organizes data so that each row represents an *observation* and each column represents a *variable*.
- Variables come in two fundamental types: *quantitative* (continuous or discrete) and *categorical* (nominal or ordinal).
- The *explanatory* variable is the variable we think might influence the *response* variable.
- *Experiments* randomly assign subjects to treatments and can establish causation. *Observational* studies merely observe what happens and can identify associations, but generally cannot establish causation.
- Association does not imply causation — this is one of the most important principles in statistics.

Many of the ideas from this chapter will be revisited as we move on to doing end-to-end data analyses. In the next chapter, you will learn about how we can design studies to collect the data we need to make conclusions with the desired scope of inference.

1.2 Data Collection and Study Design

*Before digging into the details of working with data, we pause to think about how data come to be. If data are to be used to draw broad conclusions, then it is important to understand who or what the data represent. One important aspect is **sampling** — knowing how the observational units were selected from a larger group allows us to generalize back to the population from which the data were drawn. Additionally, by understanding the structure of the study, we can separate causal relationships from mere associations. A good question to ask before analyzing any dataset is: How were these observations collected? You will learn a lot about the data by understanding its source.*

Key Concepts

- Distinguish between a sample and a population
- Recognize when it is appropriate to use sample data to make inferences about the population
- Critically examine the way a sample is selected, identifying possible sources of bias
- Recognize that random sampling is a powerful way to avoid sampling bias
- Identify other potential sources of bias that may arise in studies on humans
- Distinguish between an observational study and a randomized experiment
- Recognize that only randomized experiments can lead to claims of causation
- Distinguish between a randomized comparative experiment and a matched pairs experiment
- Identify potential confounding variables in a study

Motivation

A state in the upper Midwest is currently considering a bill that would provide amnesty to underage persons who call for help from the authorities with an overly intoxicated person. One of the state senators who represents a district that contains a university is hesitant to initially support the bill, so she has commissioned a study to learn about how the residents of her district feel about the bill. The senator will select from one of the three study options.

Study 1: “Would you support a bill that prevents authorities from giving a ticket to an underage person who has asked for help with an overly intoxicated person?” Participants would be recruited by purchasing ads on the local radio stations.

Study 2: “Would you support a bill that has the potential to allow underage people to drink with no fear of retribution?” Participants would be a random sample of all taxpayers in the district gathered by door-to-door interviews.

Study 3: “Would you support a bill that helps reduce the chances of undergraduate death due to alcohol poisoning by guaranteeing that underage people who request emergency medical assistance for someone who is very intoxicated will not receive disciplinary sanctions from the university?” Participants would be a random sample of all property owners in the district contacted to fill out a survey via email.

- a) Rank the three studies based on the question they ask. Briefly justify your ranking.

- b) Rank the three studies based on the sampling method. Briefly justify your ranking.

- c) If you could mix the questions and sampling methods, which combination do you think would be the best?

A *population* includes **all** individuals or objects of interest.

A *census* occurs when every case in the population is measured.

The *sampling frame* is a list of the names of all subjects or objects in the population.

Data are collected from a *sample*, which is a subset of the population.


Statistical inference is the process of using data from a sample to gain information about the population.

Populations and Samples

The first step in conducting research is to identify the question to be investigated. A clearly stated research question helps identify what subjects or cases should be studied and what variables are important, or the *population*. It is also important to consider how data are collected so that they are reliable and help achieve the research goals.

One way we could find an answer is to look at every case by taking a *census*. However, most of the time this is not feasible. Taking a census could take months and is often very costly. Sometimes items are destroyed in the sampling process, such as measuring the tensile strength of a wire or the length of life of a light bulb or battery. Collecting a census in these cases would be foolish. So, most of the time, researchers take a *sample* from the population and use what they find from the sample to describe the population.

In order to make sure that all units in the population are included, a *sampling frame* is compiled and used to select the subjects for the sample. This process of using a sample to describe the population is called *statistical inference*, and the accuracy of the inference can be greatly impacted by the quality of the sample and how well the sample reflects the characteristics of the population.



Class Example 1.2.1 Population Identification

For the following research questions, identify the target population and what represents an individual case

- a) What is the average mercury content in swordfish in the Atlantic Ocean?
- b) Over the last five years, what is the average time to complete a degree for UW–La Crosse undergrads?

- c) Does a new drug reduce the number of deaths in patients with severe heart disease?

Parameters and Statistics

In most statistical analyses, the research question boils down to understanding a numerical summary — perhaps a quantity you already know (like the average) or one you will learn about in this course.

A numerical summary can be calculated on either the sample, called a *sample statistic*, or the entire population, called a *population parameter*. However, measuring every unit in the population is usually impossible. So we calculate the summary from a sample and use it to estimate the corresponding population value.

A *parameter* is a number that describes a population and is almost always unknown.

A *statistic* is a number that describes a sample and is almost always known.


Class Example 1.2.2 Parameter and Statistic Identification

For the following research questions, identify the population parameter and the sample statistic.

- a) A sample of 60 swordfish have an average of 0.97 ppm
- b) An average of 4.37 years is reported from 25 UWL students as the time to complete their degree.
- c) After 1000 patients started taking a new drug for severe heart disease, 28% were reported to have died within 5 years.


Sampling from a Population

We might try to estimate the time to graduation for UW–La Crosse undergraduates by collecting a sample of graduates. All graduates in the last five years represent the *population*, and graduates who are selected for review are collectively called the *sample*. In general, we always seek to *randomly* select a sample from a population. The most basic type of random selection is equivalent to how raffles are conducted. For example, we could write each graduate's name on a raffle ticket and draw 10 tickets. The selected names would represent a random sample of 10 graduates.



If we ask a student who happens to be majoring in nutrition to select several graduates for the study, they might pick a disproportionate number of graduates from health-related fields. When selecting samples by hand, we run the risk of picking a *biased* sample, even if our bias is unintentional.

Bias occurs when the results from the sample differ from the population.



If someone is permitted to pick and choose exactly which individuals are included in the sample, it is entirely possible that the sample will overrepresent that person's interests — even if the bias is unintentional. Sampling randomly helps address this problem.

The act of taking a simple random sample helps minimize bias. However, bias can crop up in other ways. For example, asking sensitive questions may result in *response bias* because participants do not feel comfortable answering the question truthfully. If a lot of people do not respond to a survey, it will suffer from *non-response bias*. Finally, if the population from which the sample is drawn excludes groups of people,

Response bias is the tendency for people to not answer questions honestly.

Non-response bias occurs when the response rate is very low.

Undercoverage occurs when part of the population is not considered for inclusion in the sample.

then the sample will suffer from *undercoverage*.

Class Example 1.2.3 Wording bias

A survey is to be conducted using a random sample of citizens in a town, asking them if they support raising taxes to increase funding for the public school.

a) Write the question in a way that is likely to bias the results toward more *yes* answers.

b) Write the question in a way that is likely to bias the results toward more *no* answers.

Sampling Methods


Good Ways to Obtain a Sample

There are a number of appropriate ways to randomly select your sample. The most basic type of sampling is the *simple random sample* (SRS). Taking a SRS avoids sampling bias and tends to produce a good representative sample of the population. However, there are many times when we can't conduct a simple random sample. *Stratified random sampling* occurs when the population is divided into homogeneous, non-overlapping groups (called *strata*) and a simple random sample is selected from each group. *Cluster random sampling* occurs when the population is divided into many non-overlapping groups (called *clusters*), a simple random sample of clusters is obtained, and data is collected from every case in the cluster.

A *simple random sample* is obtained when every sample of a given size is equally likely to be the one selected.

Stratified random sampling occurs when the population is divided into homogeneous, non-overlapping groups (called *strata*) and a simple random sample is selected from each group.

Cluster random sampling divides the population into many non-overlapping groups (called *clusters*), randomly selects a subset of clusters, and collects data from every case in the cluster.



Class Example 1.2.4 Picking a good sampling method

Which type of sampling method is most appropriate in the following scenarios?

a) Researchers want to test a new math curriculum for third graders. They only have the budget to try the curriculum in 10 elementary schools.

b) Processor chips are continuously manufactured at a production plant. To ensure that they meet size specifications so that they will fit inside an iPhone, manufacturers want to test 100 chips throughout the day.

- c) A player's agent wants to know the average salary for each position of major league baseball.

The table below summarizes the key features of each sampling method:

Method	How it works	Best used when...
SRS	Every case has an equal chance of selection	Population is relatively homogeneous and acc
Stratified	Divide into similar groups (strata), then SRS within each	Known subgroups differ on the outcome of in
Cluster	Randomly select entire groups (clusters)	Clusters are internally diverse but similar to e
Multistage	Randomly select clusters, then SRS within each	Same as cluster, but clusters are large

Random sampling caution: Random is not the same as haphazard. We cannot obtain a random sample by haphazardly picking a sample on our own. We must use a formal random sampling method such as technology or drawing names out of a hat.

Class Activity: Sampling from Gettysburg Address Worksheet

Bad Ways to Take a Sample

Without a random sample, we may encounter *sampling bias* which occurs when the sample does not accurately represent the population of interest. One common sampling technique that will nearly always be biased is *voluntary response sampling*. With this kind of sampling, a researcher asks people to volunteer to be a part of a study. Another common type of sampling that produces biased results is the *convenience sample*. In a convenience sample, a researcher simply uses the most convenient group available as the sample.

Sampling bias occurs when the sample does not accurately represent the population of interest.

Voluntary response sampling occurs when a researcher asks people to volunteer to be participants instead of randomly selecting possible participants.

Convenience sampling happens when a researcher just uses the

Class Example 1.2.5 Driving with a pet on your lap

Over 30,000 people participated in an online poll on cnn.com conducted in April 2012 asking "Have you ever driven with a pet on your lap?" We see that 34% of the participants answered yes and 66% answered no.

- What is the sample?
- Is the variable in this study quantitative or categorical?
- Can we conclude that 34% of all drivers have driven with a pet on their lap?
- Explain why is not appropriate to generalize these results to all drivers, or even to all drivers who visit cnn.com. What is the problem with this method of data collection?

Experiments

Well-designed *experiments* incorporate four key principles:

1. **Controlling.** Researchers assign treatments to cases and do their best to *control* any other differences between the groups. For example, when patients take a drug in pill form, some patients might take the pill with a sip of water while others drink a full glass. To control for the effect of water consumption, a doctor might instruct every patient to drink a 12-ounce glass of water with the pill.
2. **Randomization.** Researchers randomly assign subjects to treatment groups to account for variables that cannot be controlled. For example, some patients may be more susceptible to a disease than others due to their dietary habits. Randomly assigning patients to treatment or control groups helps even out such differences across groups.
3. **Replication.** The more cases researchers observe, the more accurately they can estimate the effect of the explanatory variable on the response. In a single study, we **replicate** by collecting a sufficiently large sample. At a minimum, we want multiple subjects per treatment group. Replication also refers to repeating an entire study to verify earlier findings. (The *replication crisis* refers to the ongoing problem in several scientific disciplines where past findings have failed to be replicated in new studies.)
4. **Blocking.** Researchers sometimes know or suspect that variables other than the treatment influence the response. They may first group individuals based on this variable into *blocks* and then randomize cases within each block to the treatment groups. For instance, in studying the effect of a drug on heart attacks, researchers might first split patients into low-risk and high-risk blocks, then randomly assign half the patients from each block to the treatment and half to the control group.

To demonstrate the differences between some of the more commonly used experimental designs, consider the following study. A running shoe company wants to know if the type of shoe impacts the speed of runners. One aspect of the shoe that they are considering is the heel-toe drop, or the difference in height between the heel of the shoe to the toe of the shoe. Minimalist shoes reduce the heel-toe drop to 0 – 4 mm, whereas a more traditional shoe has an 8 – 12 mm heel-toe drop. This company wants to know if a lower heel-toe drop will cause runners to run more quickly, so they recruit 100 marathon runners of varying ability. For the study, the runner will run 5k, and their time will be used to establish their pace.


Randomized comparative design

In a *randomized comparative design*, we take all the participants in the study and randomize them to a different treatment group.

In an *experiment* the researcher actively controls one or more of the explanatory variables.


In a *randomized experiment* the value of the explanatory variable for each unit is determined randomly, before the response variable is measured.

In a *randomized comparative design*, we randomly assign cases to different treatment groups and then compare results on the response variable(s).



Block design

In a *block design*, we first group participants by some logical grouping that we think might have an impact on the response in addition to the variable we are studying.



In a *block design*, cases are first split into groups (called *blocks*), and within each group the value of the explanatory variable for each unit is determined randomly.

Class Example 1.2.6 *Picking the blocking factor*

Why might it make sense to group the runners into professionals vs amateurs?

Matched pairs design

In a *matched pairs design*, each participant is assigned to each of two treatment settings, responses are measured for each treatment, and the differences in responses are computed for each participant. Because two measurements are obtained for each individual, the measurements are *dependent* upon each other. The two treatments often come in the form of “Before vs. After” or “Method A vs. Method B.” In the “Method A vs. Method B” type of pairing, it is important to randomize the order of the treatments to avoid a confounding *order effect*.

In a *matched pairs* design, cases get both treatments (or cases are paired) and we examine differences in the response between the two treatments.

Class Example 1.2.7 *Describing the matched pairs running shoe experiment*

Thinking about the matched pairs design for the experiment about running shoes

a) What are the variables in the study? Are they categorical or quantitative?

b) What are the explanatory and response variables?

c) Explain how the treatments would be assigned.

d) Why might a matched pairs design be beneficial in this experiment?

Reducing Bias in Human Experiments

For an experiment, it is important to be able to determine if the treatment actually is making a difference, or if we are seeing results that would have happened regardless of the treatment. The purpose of a *control group* is to establish what we would expect to happen regardless of the treatment. Often times, in order to make sure that people adhere to the study guidelines, we use a *placebo* that is designed to look like the treatment.

The placebo makes it much easier to conduct *single-blinded* experiments or *double-blinded* experiments. Not all randomized experiments make use of control groups, placebos, or blinding; however, in studies involving humans, these techniques are often used to help make effective comparisons between groups.

The *control group* does not receive a treatment.

A *placebo* is a fake treatment that is designed to look like the treatment but has no therapeutic value.

In a *single-blind* study, participants do not know if they are in the treatment or control group.

In a *double-blind* study, neither the participants nor the people interacting with the participants knows if they are in the treatment or control group.

Class Example 1.2.8 *Depression and Prozac*

Researchers at a prominent Midwestern university would like to determine if Prozac is effective in reducing depression as determined by patient responses to a questionnaire. The researchers recruited 50 individuals


suffering from depression. Describe a completely randomized double-blind experiment using a placebo to test this association is a causal association.

Confounding Variables

A *confounding variable* is a variable that is associated with both the explanatory variable and the response variable. Because of this dual association, a confounding variable makes it impossible to determine whether the explanatory variable truly caused the observed response.

A *confounding variable* is a third variable that is associated with both the explanatory and response variable, but not necessarily of interest in the study.

Consider a classic example: ice cream sales and drowning rates both increase during summer months. If we looked only at the association between ice cream sales and drownings, we might (absurdly) conclude that ice cream causes drowning. The confounding variable is temperature (or season) — warm weather drives both increased ice cream consumption and increased swimming, which leads to more drowning incidents.



Confounding diagram.

Summary

In this chapter, we explored how data are collected and why the method of collection matters for the conclusions we can draw.

Key concepts:

- The *population* is the entire group of interest; a *sample* is a subset from which we collect data. A *census* collects data on every member of the population but is usually impractical.
- Good sampling methods include *simple random sampling (SRS)*, *stratified sampling*, and *cluster sampling*. Each has advantages depending on the population's structure and practical constraints.
- *Bias* can enter through non-random sampling (selection bias), non-response, or convenience sampling.
- In an *experiment*, researchers actively assign treatments to subjects. In an *observational study*, researchers simply observe without intervening.
- *Confounding variables* are associated with both the explanatory and response

variables and can create misleading associations.

- *Blinding* (single and double) and placebos help reduce bias in human experiments.

1.3 Exploring Categorical Data

This chapter focuses on **categorical data** — variables whose values are categories or labels rather than numbers. We begin with visualizations for a single categorical variable (bar charts, pie charts, waffle charts), then move to tools for exploring the relationship between two categorical variables (contingency tables and stacked bar charts). We close with principles for creating effective, honest data visualizations.

Key Concepts

- Display information from a categorical variable in a table or a graph
- Use information about a categorical variable to find and report a proportion, using correct notation
- Display information about a relationship between two categorical variables in a contingency table
- Use a contingency table to find proportions

Technology for Chapter

StatLens Use the [One Categorical Variable](#) explorer for bar charts and pie charts. Use the [Two Categorical Variables](#) explorer for contingency tables and stacked bar charts.

jamovi Use Analyses>Exploration>Descriptives for bar charts for one categorical variable. Use Analyses>Frequencies>Independent Samples for two categorical variables.

One Categorical Variable

When we want to summarize one categorical variable, we use a *proportion* which tells us what fraction of the sample is in each group. In this course, we will use p to represent the population proportion, and $\hat{p}$ (p-hat) to represent the sample proportion. The sample proportion is given by

$$\hat{p} = \frac{x}{n}$$

where x is the number of cases in a given category and n is the total number of cases.

Numerical displays

A single categorical variable is often summarized using a *frequency table* to display the number of observations that fall into each category. An alternative to the frequency table is the *relative frequency table* which displays the proportion of the sample that is in each category.

The *proportion* (or *relative frequency*) in a category is found by dividing the number of cases in that category by the total number of cases.

A *frequency table* is a table that displays the count for each category.

A *relative frequency table* is a table that displays the proportion of the sample in each category.

Class Example 1.3.1 Hair Color

The table below gives the frequency of hair color for a random sample of 55 UWL students. Find the proportions of students with each hair color and use them fill in the relative frequency column.

	Frequency	Relative Frequency
Brown	30	
Blonde	14	
Red	2	
Black	5	
Colorful	3	
Total	54	

Frequency table for hair color.

Graphical summaries

The standard graphical summaries for a categorical variable are the *bar chart* and the *pie chart*. A bar chart is the most common way to display the distribution of a single categorical variable. Each category gets its own bar, and the height (or length) of the bar represents either the count or the proportion of observations in that category. A pie chart divides a circle into slices, where each slice represents a category and the size of the slice corresponds to the proportion of observations in that category.

A *bar chart* includes one bar for each category, and the height of the bar is the frequency the category appears.

In a *pie chart*, the proportions correspond to the areas of sectors of a circle.

Class Example 1.3.2 *Hair Color revisited*

Sketch a bar chart and a pie chart for the hair color of random sample of 55 UWL students.

Pie charts can give a quick high-level overview, especially when a variable has only a few levels or when each level represents a simple fraction (one-half, one-quarter, etc.). However, bar charts are almost always easier to read because our eyes are better at comparing lengths than angles or areas.

Two Categorical Variables

Sometimes our goal is to examine whether there is a relationship between two categorical variables. To investigate these relationships, we use a *contingency table*, or *two-way table*. In a two-way table, the rows represent the groups of the first category, and the columns represent the groups of the second category. Each cell of the table contains the number of cases that are in both the row and the column category.

A *contingency table* summarizes the data for two categorical variables.

The rows represent the levels of one variables, the columns represent the levels of the other.

Class Example 1.3.3 *Relationship status and class level*


169 college students were asked about their relationship status and class year (lower=freshman/sophomore, upper=junior/senior). The results are given in the table.

	Upper	Lower	Total
In a relationship	32	10	42
It's complicated	12	7	19
Single	63	45	108
Total	107	62	169

Contingency table for relationship status and class year.

- What proportion of students in this sample are in a relationship?
- What proportion of lower class students in this sample are in a relationship?
- What proportion of upper class students in this sample are in a relationship?
- Using $\hat{p}_U$ to represent the proportion of upper class students in a relationship and $\hat{p}_L$ to represent the proportion of lower class students in a relationship, find the difference in proportions $\hat{p}_U - \hat{p}_L$
- What proportion of the people who are in a relationship in this sample are upper class students?
- What proportion of the people in this sample are upper class students and in a relationship?
- Answers c), e) and f) are all ways of representing upper class students in a relationship. Why are they different numbers?

Contingency tables can be displayed graphically using an extension of the bar chart. There are two common variants: a stacked bar chart and a side-by-side bar chart.



Class Example 1.3.4 *Handedness and occupation*

In a study of handedness in occupations, 10 of 118 psychiatrists were left-handed, 26 of 148 architects were left-handed, 5 of 132 orthopedic surgeons were left-handed, and 16 of 105 lawyers were left-handed.

a) Make a two-way table of this relationship.

b) What proportion of the people in the sample are left-handed?

c) What proportion of left-handed people are architects?

d) Is your answer in part c) the same or different than the proportion of architects who are left-handed?

Effective Data Visualization

Graphs can powerfully communicate ideas directly and quickly. However, there are times when a visualization conveys a message that is inaccurate or misleading. The following are guiding principles for creating clear, honest, and effective visualizations.

- *Keep it simple:* Colors should be used purposefully and can be used to draw attention
 - Colors used only for decoration can be distracting and can confuse readers
- *Tell a story:* Include annotations (text, lines or labels that provide context beyond the raw data)
- *Order matters:* The arrangement of categories in a bar chart can help or hinder understanding
 - Alphabetical order is rarely the most informative choice
- *Make labels readable*
- *Select meaningful colors:* Default or rainbow color schemes are not always the best choice

Summary

This chapter introduced tools for exploring categorical data.

-For a single categorical variable, *bar charts* are the primary visualization, with *pie charts* reserved for simple cases.

-For two categorical variables, *contingency tables* organize counts, and stacked and side-by-side bar charts provide visual comparisons.

-Effective visualizations follow principles of simplicity, purposeful use of color, meaningful ordering, and accessibility.

1.4 Exploring Quantitative Data

*In the previous chapter we explored categorical data. Now we turn to **quantitative data** — variables that take numerical values where arithmetic makes sense (heights, incomes, interest rates). This chapter teaches you to both see and measure distributions. We introduce dot plots and histograms alongside the mean and median, box plots alongside the IQR, and the standard deviation alongside histogram shapes. By the end, you will be able to describe any distribution in terms of its shape, center, spread, and unusual features.*

Key Concepts

- Understand the difference between displays of categorical and quantitative data
- Use a dotplot or histogram to describe the shape of a distribution
- Calculate the mean and the median for a set of data values, with appropriate notation
- Identify the approximate locations of the mean and the median on a dotplot or histogram
- Explain how outliers and skewness affect the values for the mean and median
- Recognize the uses and meaning of the standard deviation
- Interpret a five number summary
- Use the range, the interquartile range, and the standard deviation as measures of spread
- Describe the advantages and disadvantages of the different measures of spread
- Identify outliers in a dataset based on the IQR method
- Use a boxplot to describe data for a single quantitative variable
- Use the empirical rule to estimate the middle 68%, 95%, or 99.7% for bell-shaped distributions
- Use technology to compute summary statistics for a quantitative variable

Technology for Chapter

StatLens Use the [One Quantitative Variable](#) to visualize distributions as histograms, dot plots, or box plots, and compute means, medians, standard deviations, and quartiles for any dataset.


jamovi Use Analyses>Exploration>Descriptives to visualize distributions as histograms or box plots, and compute means, medians, standard deviations, and quartiles for any dataset.

Dot Plots and Histograms

Dot Plots

A *dot plot* is the simplest display for a single quantitative variable. Each observation is represented by a dot placed above its value on a number line. When observations share the same value (or are rounded to the same value), the dots stack vertically.

Consider the interest rates on 50 loans from a lending platform. In a dot plot, each of the 50 loans is a single dot. The red triangle marks the mean.




Dot plots are most useful for small datasets (up to about 50–100 observations) where you want to see every individual value. For larger datasets, histograms provide a better summary.

A *histogram* groups quantitative data into bins (intervals) and displays the count or proportion of observations in each bin as a bar. Unlike bar charts for categorical data, histogram bars touch each other because the bins represent a continuous range of values.

Histograms

To build a histogram, we:

1. Choose a bin width (e.g., 2.5 percentage points for interest rates)
2. Count how many observations fall in each bin
3. Draw bars whose heights equal the counts



A distribution is considered *symmetric* if we can fold the plot over a vertical center line and both sides match closely.

A distribution is considered *skewed left* if the longer tail of the distribution is to the left.

A distribution is considered *skewed right* if the longer tail of the distribution is to the right.

Histograms are especially useful for identifying the *shape* of a distribution. One key characteristic is symmetry. When we say a distribution is *symmetric* it means that the left-side of a distribution is very similar to the right-side of the distribution. We say a distribution is *skewed left* when the tail of the distribution is longer on the left. Similarly, we say that a distribution is *skewed right* when the tail of the distribution is longer on the right.

Class Example 1.4.1 *Shape of the interest rates*

Describe the shape of the histogram for the interest rate plot.

Measures of Center: Mean and Median

One type of numerical summaries you will see for a dataset is called measures of center. Measures of center, provide us with a single number that best describes a quantitative variable's distribution. Essentially, they give us a guess of what a typical value would look like. There are two measures of center we will use in this class: the *mean* and *median*.

The mean

The *mean* of a sample tells us the average. For a sample of observations $x_1, x_2, \dots, x_n$, the mean is given by

$$\text{Mean} = \frac{x_1 + x_2 + \dots + x_n}{n} = \frac{\sum_{i=1}^n x_i}{n}$$

where n is the number of observations.

We will use μ (pronounced “mew”) to represent the population mean and $\bar{x}$ (read “x-bar”) to represent the sample mean. We often cannot measure μ directly because we rarely have access to every member of the population. Instead, we estimate μ using the sample mean $\bar{x}$.

The median

The *median*, denoted m , divides the sample into two groups of equal size. We find the median by following two steps:

1. Sort the sample values.
2. Find the middle value:
 - If n is odd, the median is the middle value in the sorted list.
 - If n is even, the median is the average (midpoint) of the middle two values in the ordered list.

The dot plot below shows both the mean and the median marked on the interest rate data. Notice that the median (9.93%) is lower than the mean (11.57%) — the few loans with very high interest rates pull the mean to the right, but the median stays anchored in the middle of the data.


The *mean* of a data set is the sum of all its values divided by the number of observations.

The *median* of a dataset is the middle entry of an ordered list of data. If there are an even number of entries, then we take the average of the middle two values.

Notation:

μ : population mean

$\bar{x}$: sample mean


Class Example 1.4.2 *Ants on a sandwich*

The number of ants climbing on a piece of a peanut butter sandwich left on the ground near an anthill for a few minutes was measured 7 different times, and the results are: 43, 59, 22, 25, 36, 47, 19

- Calculate the mean number of ants.
- Calculate the median number of ants.
- Suppose one of the sandwich bits was extremely appealing to the ants, and instead of 59, the number of ants for that bit was actually 159, giving the 7 values as: 43, 159, 22, 25, 36, 47, 19. Compute the mean and median for this new dataset.
- Which statistic (mean or median) is impacted most by the outlier?

How skewness affects the mean and median

In a symmetric distribution, the mean and the median are approximately equal ($\text{mean} \approx \text{median}$). In a right-skew distribution, the mean is pulled towards the right tail and is larger than the median ($\text{mean} > \text{median}$). In a left-skewed distribution, the mean is pulled towards the left tail and is smaller than the median ($\text{mean} < \text{median}$).

When to use the mean vs. the median

As we saw in the last example, the median is not impacted by extreme data values (outliers) like the mean is. We say that the median is resistant. Although the mean

is a more commonly used measure of center, the median may be preferred in cases with extreme values or a high degree of skew.

If the concern is the total or aggregate, the mean is the best. If the concern is a typical observation, the median is preferred, especially when the distribution is skewed or contains extreme values.

Measures of Spread

Knowing the center of a distribution is important, but it tells only part of the story. Two distributions can have the same center but look very different if one is tightly clustered and the other is widely dispersed. We need measures of *spread* (also called variability or dispersion).

A measure of *spread* tells us how much variability exists in the data.

The range

The simplest measure of spread is the *range*: the difference between the maximum and minimum values.

The *range* is the difference between the maximum and minimum of the data.

$$\text{Range} = \text{Maximum} - \text{Minimum}$$

While easy to compute, the range has a major weakness: it depends entirely on the two most extreme observations. A single extreme value can dramatically increase the range, giving a misleading impression of the overall variability. For this reason, we usually prefer the IQR or standard deviation.

The interquartile range (IQR)

The *interquartile range* measures the spread of the middle 50% of the data. To find the IQR, we need to first estimate Q_1 and Q_3 , the first and third quartiles. The first quartile, Q_1 , is the value which 25% of the data falls below; and the third quartile, Q_3 , is the value which 75% of the data falls below. To find Q_1 , we need to take the median of all the values below the median. To find Q_3 , we need to take the median of all the values above the median. The IQR is given by

The *IQR* (Interquartile range) is the difference between the 1st and 3rd quartiles of the data.

$$\text{IQR} = Q_3 - Q_1.$$

The five-number summary

A way to combine the information from the range, the IQR, and a measure of center is through the *five-number summary*. The five-number summary consists of the minimum, Q_1 , the median, Q_3 , and the maximum of the data set.

Class Example 1.4.3 *Ants revisited*

In the previous example, we found that for the 7 different times, the results were 19, 22, 25, 36, 43, 47, 59 (sorted in ascending order)

a) Find the five number summary for this data.

a) Find the IQR for this data

b) Find the range for this data

c) Find the range and IQR if the value 59 was actually 159?

d) Which statistic (range or IQR) is impacted most by the extreme value?

Variance and standard deviation

The **standard deviation** is the most commonly used measure of spread. The standard deviation gives an estimate for the average distance each observation is from the mean.

We start with the concept of a **deviation**: the distance of an observation from the mean.

$$\text{deviation of } x_i = x_i - \bar{x}$$

Some deviations are positive (observations above the mean) and some are negative (observations below the mean). If we simply averaged the deviations, the positives and negatives would cancel out, giving us zero. To avoid this, we *square* each deviation before averaging. The average of the squared deviations, using $n - 1$ in the denominator is the *variance* and is given by

$$\text{Variance} = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n - 1} = \frac{(x_1 - \bar{x})^2 + (x_2 - \bar{x})^2 + \cdots + (x_n - \bar{x})^2}{n - 1}$$

The *standard deviation* is the square root of the variance:

$$\text{Standard Deviation} = \sqrt{s^2} = \sqrt{\frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n - 1}}$$

In this course, we will use σ^2 (pronounced “sigma squared”) to represent the population variance and s^2 to represent the sample variance. We will use σ (pronounced “sigma”) to represent the population standard deviation and s to represent the sample standard deviation. When the standard deviation is large, it means there is a lot of variability amongst the observations. When the standard deviation is small, it means there is a lot of consistency amongst the observations.

Looking back at the interest rate data with shaded bands marking one and two standard deviations from the mean. The standard deviation of 5.05 percentage points gives us a sense of how spread out the data are around the mean of 11.57%.

The *standard deviation* of a variable is a rough estimate of the typical distance of a data value from the mean.


A *deviation* is the difference between an observation and the mean

The *variance* is the average of the squared deviations.

Notation:

σ : population standard deviation

s : sample standard deviation



The standard deviation tells us that interest rates typically deviate about 5 percentage points from the mean (11.57%).

Class Example 1.4.4 *Ants revisited*

- Find the standard deviation of the ants on a sandwich data.
- Find the standard deviation of the data if the value 59 was actually 159.

Outliers and Robust Statistics

Identifying outliers

An *outlier* is an observation that appears extreme relative to the rest of the data. A common rule for identifying outliers is the **1.5 IQR rule**: A data value, x_0 is considered to be an outlier if

$$x_0 < Q_1 - 1.5(IQR) \quad \text{or} \quad x_0 > Q_3 + 1.5(IQR).$$

An *outlier* is a data point that is unusual given the rest of the data.

Examining data for outliers serves several purposes:

- Identifying strong skew in the distribution
- Identifying possible data collection or data entry errors
- Providing insight into interesting or unusual properties of the data

Robust statistics

We call a statistic **robust** (or **resistant**) if extreme observations have little effect on its value.

We say that a statistic is *robust*, or *resistant*, if its is relatively unaffected by extreme values.

Class Example 1.4.5 *Resistant statistics*

Look back at your answers to the questions about ants on a sandwich.

a) Which measures of center (mean, median) are resistant?

b) Which measures of spread (IQR, range, standard deviation) are resistant?

Robust vs. non-robust statistics.

Robust (resistant to outliers)	Not robust (sensitive to outliers)
Median	Mean
IQR	Standard deviation
	Range

When a distribution is skewed or contains outliers, robust statistics (median and IQR) often give a more representative summary of the “typical” observation. When a distribution is roughly symmetric with no extreme outliers, the mean and standard deviation work well and carry additional mathematical advantages.

Box plots

So far in this class, we have used histograms and dot plots to represent data graphically. There is another very commonly used graphical summary called the *boxplot*. The boxplot gives us a graphical display of the *five-number summary* which consists of the minimum, Q_1 , median, Q_3 , and the maximum of a dataset.

The *boxplot* is a graphical display of the *five-number summary* which consists of the minimum, Q_1 , median, Q_3 , and the maximum of a dataset.


Constructing a box plot

The components of a box plot are:

1. *The box*: Spans from Q_1 (the 25th percentile) to Q_3 (the 75th percentile). This box contains the middle 50% of the data. The length of the box is the *interquartile range* ($IQR = Q_3 - Q_1$).
2. *The median line*: A line inside the box marks the median (the 50th percentile), which divides the data in half.
3. *The whiskers*: Lines extending from the box toward the smallest and largest observations that are *not* outliers. Specifically, the whiskers extend to the most extreme data points within $1.5 \times IQR$ of the box.
4. *Outlier points*: Any observations beyond $1.5 \times IQR$ from the edges of the box are plotted individually as dots. These are potential outliers.

The following gives a general sketch of a boxplot:

A box plot of interest rates shows: the left edge of the box at about 8%, the median line at about 10%, and the right edge of the box at about 14%. Two dots appear beyond the right whisker at about 25% and 26%.


Class Example 1.4.6 *Population of U.S. states*


The following data represent the populations of 12 of the 50 U.S. states, in millions of people, in ascending order.

0.506	0.830	1.315	1.813	2.333	2.953
3.524	4.198	5.504	6.207	8.685	22.472

- Identify the five-number summary for the state populations.
-
- Draw a boxplot for the data. If there are any outliers, provide a justification as to why they are outliers and identify them on your boxplot.
-

Class Example 1.4.7 *Gross state product*

The figure below gives a boxplot of the gestation (in days) for 54 mammal species.



- a) Are there any outliers? If so, how many and estimate the values of the outliers.
-
- b) Estimate the median, IQR and range of the data.
-
- c) How would you describe the shape of this distribution?
-
- d) Do you expect the mean to be less than, approximately equal to, or greater than the median? Explain.
-

Describing a distribution

A complete description of a distribution should address three things: *shape*, **center**, and *spread*.

A good description of a distribution should always:

1. Name the *shape*: symmetric or skewed (and direction)
2. Give the *center*: report the mean and/or median with units
3. Give the *spread*: report the standard deviation and/or IQR with units
4. Note any *unusual features*: outliers, gaps, clusters

5. Use *context*: refer to the actual variable and its units, not just abstract numbers

A way to describe a distribution is as follows: “The distribution of commute times is right skewed, with a median of 22 minutes and an IQR of 15 minutes. A few commuters have commute times exceeding 90 minutes.”


The Empirical Rule (68-95-99.7 rule)

For distributions that are approximately *bell-shaped*, the standard deviation provides a useful ruler. Statisticians call this particular bell shape the *normal distribution* — it comes up so often in statistics that it has its own name. We’ll study it formally later, but for now, just know that “bell-shaped” and “normal” mean the same thing.

The empirical rule.

For a roughly bell-shaped distribution:

- About **68%** of the data fall within 1 standard deviation of the mean: $(\bar{x} - s, \bar{x} + s)$
- About **95%** of the data fall within 2 standard deviations of the mean: $(\bar{x} - 2s, \bar{x} + 2s)$
- About **99.7%** of the data fall within 3 standard deviations of the mean: $(\bar{x} - 3s, \bar{x} + 3s)$



Class Example 1.4.8 *SAT scores*

The SAT scores (standardized test scores) are approximately bell-shaped with a mean of 1150 inches and a standard deviation of 150 inches. Use the empirical rule to estimate the range of scores for the middle 95% of students.

Summary

This chapter introduced the tools for exploring *quantitative* data — both visual and numerical.

- *Dot plots* show individual values for small datasets, *histograms* reveal the shape and density of a distribution, density plots provide smoothed alternatives, and *box plots* give a compact *five-number summary*.
- Distribution shape is described using symmetry (*left skewed*, *symmetric*, *right skewed*).
- The *mean* and *median* measure center, with the mean being the balancing point and the median being the middle value.
- The *standard deviation* and *IQR* measure spread, with the standard deviation roughly representing the typical deviation from the mean and the *IQR* representing the range of the middle 50%.
- The *median* and *IQR* are *robust* to outliers, while the mean and standard deviation are sensitive to extreme values.
- The *empirical rule* connects the standard deviation to the distribution shape for bell-shaped data: about 68%, 95%, and 99.7% of observations fall within 1, 2, and 3 standard deviations of the mean.

1.5 Relationships Between Variables

The preceding chapters focused on exploring one variable at a time. But the most interesting statistical questions involve **relationships between variables**. Does higher education lead to higher income? Is a drug more effective for one group than another? This chapter introduces tools for exploring relationships: side-by-side box plots and faceted histograms for comparing a quantitative variable across groups, and scatterplots for examining the association between two quantitative variables. We close with the critical distinction between association and causation.

Key Concepts

- Use a side-by-side graph to visualize a relationship between quantitative and categorical variables
- Examine a relationship between quantitative and categorical variables using comparative summary statistics

Technology for Chapter

StatLens Use the [Quantitative by Group](#) to compare distributions across groups with side-by-side box plots, histograms, and density overlays. Use the [Two Quantitative Variables](#) for scatterplots of two quantitative variables.

jamovi Use Analyses>Exploration>Descriptives, using the “Split by” input, to visualize distributions as histograms or box plots, and compute means, medians, standard deviations, and quartiles for any dataset. Use Analyses>Exploration>Scatterplot for scatterplots of two quantitative variables.


Comparing Groups

Some of the most interesting investigations involve comparing a quantitative variable across groups defined by a categorical variable. Sometimes when investigating the relationship between categorical and quantitative variables, it is helpful to plot them *side-by-side*.

We use *side-by-side* plots to visualize the relationship between a quantitative variable and a categorical variable.

Class Example 1.5.1 *Average household incomes broken down by U.S. geographical region*

The figure below shows a side-by-side boxplot of average U.S. household income broken down by geographical region of the U.S.



a) Use these side-by-side boxplots to discuss how the average U.S. household income differs by region.

b) Are there any outliers? If so, how many and estimate the values of the outliers.

c) For which region(s) would it be more appropriate to use the median instead of the mean to describe the average U.S. household income? Justify your answer.

Scatterplots

A *scatterplot* displays the relationship between two quantitative variables. Each observation is represented as a point, with one variable on the horizontal axis and the other on the vertical axis. When looking at a scatterplot, we should consider four different aspects of the plot:

A *scatterplot* is a graph of the relationship between two quantitative variables. If there are explanatory and response variables, we put the explanatory variable on the horizontal (x) axis and the response on the vertical (y) axis.

1. *Direction*: Is the association *positive* (both variables tend to increase together) or *negative* (one increases as the other decreases)?
2. *Form*: Is the relationship approximately *linear* (following a straight-line pattern) or *nonlinear* (curved)?
3. *Strength*: Is the association *strong* (points closely follow a pattern), *moderate*, or *weak* (points are loosely scattered)?
4. *Unusual observations*: Are there any *outliers* — points that deviate markedly from the overall pattern?

Example 5.2 Describing Associations

Consider a dataset of 50 loans, where we plot total income on the horizontal axis and loan amount on the vertical axis:



Describe the association seen in the scatterplot in Figure 2.

Associations vs. causation

We have explored the association between variables, but an association is not the same as causation. When two variables are associated, it means they tend to vary together. That does not necessarily mean one variable *causes* the other. There may be a *confounding variable* that is related to both, creating the illusion of a direct relationship.

A *confounding variable* is a third variable that is associated with both the explanatory and response variable, but not necessarily of interest in the study.

Example 5.3 GPAs and Breakfast

A study finds that students who eat breakfast tend to have higher GPAs than students who skip breakfast. Can we conclude that eating breakfast causes higher GPAs?

Summary

This chapter introduced tools for exploring relationships between variables.

- For comparing a *quantitative variable across groups* defined by a categorical variable, side-by-side box plots provide a compact comparison of centers and spreads, faceted histograms reveal full distributional shapes, and ridge plots overlay density curves for compact multi-group comparison.
- For *two quantitative variables*, scatterplots reveal the direction, form, strength, and unusual features of an association.
- We closed with the fundamental principle that *association does not imply causation* — confounding variables can create misleading associations, and establishing causation generally requires a randomized experiment.

2.1 Categorical Variables

Key Concepts

- Display information from a categorical variable in a table or a graph
- Use information about a categorical variable to find and report a proportion, using correct notation
- Display information about a relationship between two categorical variables in a two-way table
- Use a two-way table to find proportions

Summarizing one categorical variable

When we want to summarize one categorical variable, we use a *proportion* which tells us what percentage of the sample is in each group. In this course, we will use p to represent the population proportion, and $\hat{p}$ (p-hat) to represent the sample proportion.

The *proportion* (or *relative frequency*) in a category is found by dividing the number of cases in that category by the total number of cases.

Numerical displays

A single categorical variable is often summarized using a *frequency table* to display the number of observations that fall into each category. An alternative to the frequency table is the *relative frequency table* which displays the proportion of the sample that is in each category.

A *frequency table* is a table that displays the count for each category.

A *relative frequency table* is a table that displays the proportion of the sample in each category.

Graphical summaries

The standard graphical summaries for a categorical variable are the *bar chart* and the *pie chart*. Graphical summaries make it very easy to see which category had the largest number of observations in it.

A *bar chart* includes one bar for each category, and the height of the bar is the frequency the category appears.

In a *pie chart*, the proportions correspond to the areas of sectors of a circle.

Summarizing two categorical variables

Sometimes our goal is to examine whether there is a relationship between two categorical variables. To investigate these relationships, we use a *two-way table*. In a two-way table, the rows represent the groups of the first category, and the columns represent the groups of the second category. Each cell of the table contains the number of cases that are in both the row and the column category.

A *two-way table* is used to show the relationships between two categorical variables.

Class Example 2.1.1 Relationship status and gender

169 college students were asked about their relationship status and gender. The results are given in the table.

	Female	Male	Total
In a relationship	32	10	42
It's complicated	12	7	19
Single	63	45	108
Total	107	62	169

Contingency table for relationship status and gender. {.striped}

- What proportion of students in this sample are in a relationship?
- What proportion of males in this sample are in a relationship?
- What proportion of females in this sample are in a relationship?
- Using $\hat{p}_F$ to represent the proportion of females in a relationship and $\hat{p}_M$ to represent the proportion of males in a relationship, find the difference in proportions $\hat{p}_F - \hat{p}_M$
- What proportion of the people who are in a relationship in this sample are female?
- What proportion of the people in this sample are female and in a relationship?
- Answers c), e) and f) are all ways of representing females in a relationship. Why are they different numbers?

Class Example 2.1.2 *Handedness and occupation*

In a study of handedness in occupations, 10 of 118 psychiatrists were left-handed, 26 of 148 architects were left-handed, 5 of 132 orthopedic surgeons were left-handed, and 16 of 105 lawyers were left-handed.

a) Make a two-way table of this relationship.

b) What proportion of the people in the sample are left-handed?

c) What proportion of left-handed people are architects?

d) Is your answer in part c) the same or different than the proportion of architects who are left-handed?

Class Example 2.1.3 *Screen time for 5-year olds*

A survey in 2019 asked a random sample of 400 adults “What do you think the appropriate screen time is for a 5-year old?” The results are given in the table.

Response	Frequency	Rel. Frequency
None	35	
30 minutes	160	
80 minutes	159	
4 hours or more	46	
Total	400	

Frequency of opinions for 5-year old screen time limitations. { .striped }

a) What proportion do not believe any screen time is appropriate?

b) What percent of people in the sample believe at least one hour of screen time is appropriate?

c) Fill in the relative frequency column for this table.

2.2 One Quantitative Variable: Shape, Center, Spread

Key Concepts

- Understand the difference between displays of categorical and quantitative data
- Use a dotplot or histogram to describe the shape of a distribution
- Calculate the mean and the median for a set of data values, with appropriate notation
- Identify the approximate locations of the mean and the median on a dotplot or histogram
- Explain how outliers and skewness affect the values for the mean and median
- Use technology to compute summary statistics for a quantitative variable
- Recognize the uses and meaning of the standard deviation
- Interpret a five number summary or percentiles
- Use the range, the interquartile range, and the standard deviation as measures of spread
- Describe the advantages and disadvantages of the different measures of spread
- Identify outliers in a dataset based on the IQR method

Describing quantitative data

One of the easiest ways to describe a quantitative variable is to make a *dot plot* or a *histogram* and look at the shape and center of the distribution. In fact, one of the major differences between the summaries of categorical and quantitative variables is that for quantitative variables, we often care about the shape and center of the distribution. When we say a distribution is *symmetric* it means that the left-side of a distribution is very similar to the right-side of the distribution. We say a distribution is *skewed left* when the tail of the distribution is longer on the left. Similarly, we say that a distribution is *skewed right* when the tail of the distribution is longer on the right.


A distribution is considered *symmetric* if we can fold the plot over a vertical center line and both sides match closely.

A distribution is considered *skewed left* if the longer tail of the distribution is to the left.

A distribution is considered *skewed right* if the longer tail of the distribution is to the right.

Class Example 2.2.1 *NBA Salaries*

The histogram in the figure below shows the distribution of player salaries (in millions of dollars) for a sample of 467 players during the 2022 National Basketball Association (NBA) season.



a) Describe the shape of the distribution.

Measures of center: Mean and median

One type of numerical summaries you will see for a dataset is called measures of *center*. Measures of center, provide us with a single number that best describes a quantitative variable's distribution. Essentially, they give us a guess of what a typical value would look like. There are two measures of center we will use in this class: the *mean* and *median*. The *mean* of a sample tells us the average. For a sample of observations $x_1, x_2, \dots, x_n$, the mean is given by

$$\text{Mean} = \frac{x_1 + x_2 + \dots + x_n}{n} = \frac{\sum_{i=1}^n x_i}{n}$$

where n is the number of observations.

In this course, we will use μ to represent the population mean and $\bar{x}$ to represent the sample mean.

The *mean* of a data set is the sum of all its values divided by the number of observations.

The *median* of a dataset is the middle entry of an ordered list of data. If there are an even number of entries, then we take the average of the middle two values.

Notation:

μ : population mean
 $\bar{x}$: sample mean

The *median*, denoted m , divides the sample into two groups of equal size. We find the median by following two steps:

1. Sort the sample values.
2. Find the middle value:
 - If n is odd, the median is the middle value in the sorted list.
 - If n is even, the median is the average (midpoint) of the middle two values in the ordered list.

Class Example 2.2.2 *Ants on a sandwich*

The number of ants climbing on a piece of a peanut butter sandwich left on the ground near an anthill for a few minutes was measured 7 different times, and the results are: 43, 59, 22, 25, 36, 47, 19

- a) Calculate the mean number of ants.

- b) Calculate the median number of ants.

- c) Suppose one of the sandwich bits was extremely appealing to the ants, and instead of 59, the number of ants for that bit was actually 159, giving the 7 values as: 43, 159, 22, 25, 36, 47, 19. Compute the mean and median for this new dataset.

- d) Which statistic (mean or median) is impacted most by the outlier?

Measures of spread: IQR, range

In addition to measures of center, we also should report measures of spread. A measure of spread is a statistic that indicates the type of variability we see for a single variable. The *IQR* (interquartile range) is a statistic that tells us the range of the middle 50% of the dataset. To find the IQR, we need to first estimate Q_1 and Q_3 , the first and third quartiles. To find Q_1 , we need to take the median of all the values below the median. To find Q_3 , we need to take the median of all the values above the median. The IQR is given by

$$IQR = Q_3 - Q_1.$$

To find the *range* of the data, you take the difference between the maximum and minimum of the data.

A measure of *spread* tells us how much variability exists in the data.

The *IQR* (Interquartile range) is the difference between the 1st and 3rd quartiles of the data.

The *range* is the difference between the maximum and minimum of the data.

Class Example 2.2.3 *Ants revisited*

In the previous example, we found that for the 7 different times, the results were 19, 22, 25, 36, 43, 47, 59 (sorted in ascending order)

- a) Find the IQR for this data

- b) Find the range for this data
-
- c) Find the range and IQR if the value 59 was actually 159?
-
- d) Which statistic (range or IQR) is impacted most by the outlier
-

Measures of spread: Standard deviation

In addition to IQR, another commonly used measure of spread is the *standard deviation*.

The standard deviation gives is an estimate for the average distance each observation is from the mean and is given by

$$\text{Standard deviation} = \sqrt{\frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n - 1}}.$$

In this course, we will use σ to represent the population standard deviation and s to represent the sample standard deviation. When the standard deviation is large, it means there is a lot of variability amongst the observations. When the standard deviation is small, it means there is a lot of consistency amongst the observations.

The *standard deviation* of a variable is a rough estimate of the typical distance of a data value from the mean.

Notation:
 σ : population standard deviation
 s : sample standard deviation

Class Example 2.2.4 *Ants revisited*

- a) Find the standard deviation of the ants on a sandwich data.
-
- b) Find the standard deviation of the data if the value 59 was actually 159.
-
- c) Why should we square the deviations before adding them up?
-

Outliers

In the last part of the previous example, we introduced an *outlier*, or unusual value, into the dataset. We can use a simple rule of thumb to determine whether an observation is an outlier. A data value, x_0 , is considered to be an outlier if

$$x_0 < Q_1 - 1.5(IQR) \quad \text{or} \quad x_0 > Q_3 + 1.5(IQR).$$

To pick an appropriate statistic to represent the data, we need to know if it will be impacted by outliers. We call a statistic *resistant* if it is not impacted by outliers.

An *outlier* is a data point that is unusual given the rest of the data. A statistic is *resistant* if it is not impacted by outliers.

Class Example 2.2.5 *Resistant statistics*

Look back at your answers to the questions about ants on a sandwich.

a) Which measures of center (mean, median) are resistant?


b) Which measures of spread (IQR, range, standard deviation) are resistant?

Estimating summary statistics from a graphical summary

We may not always have access to the full dataset to come up with the mean or median from a sample. Sometimes we need to estimate these values from a graphical summary. For symmetric data, the mean should be very close to the median. If the data are skewed left, then the mean will be less than the median. On the other hand, for skewed right data, the mean will be greater than the median.

Class Example 2.2.6 *The Masters*

The ages for all golfers who won The Masters from 1934 - 2023 is shown in the figure below. There are 87 champions (less than the number of years due to World War 2) represented in the dotplot.



a) Use the dotplot to estimate the median age of a Masters Champion between 1934 and 2023.

b) Should the mean age be lower than, about the same as, or higher than the median? Explain.

c) Estimate the IQR of the age from all golfers.

Estimating standard deviation from a histogram

As a general rule, for *symmetric* and *bell-shaped* histograms, about 95% of the data should fall within two standard deviations from the mean. About 100% of the data should fall within three standard deviations from the mean.

Class Example 2.2.7 *Percent of Body Fat in Men*

The variable *BodyFat* in the *BodyFat* dataset gives the percent of weight made up of body fat for 100 men. For this sample, $\bar{x} = 18.6$ and $s = 8.0$. The distribution of body fat values is roughly symmetric and bell-shaped. Find an interval that is likely to contain approximately 95% of the data values.

Choosing the best summary statistics

When picking summary statistics to best represent the data, you should always consider the shape of your distribution. For *symmetric* and *bell-shaped* distributions, you should use *mean* and *standard deviation* to describe the center and spread of the data. If you have a distribution that is *skewed*, then you should use *resistant* statistics like *median* and *IQR* to describe the center and spread of the data.

Class Example 2.2.8 *NBA Salaries revisited*

Thinking back to the NBA salaries histogram.

- a) Which statistic, mean or median, would be most appropriate to represent a typical NBA salary? Why did you select that statistic?

- b) Which statistic, IQR or standard deviation, would be most appropriate to represent the spread of the distribution for NBA salaries? Why did you select that statistic?

Percentiles

Most of you have seen a percentile at some point in the past. Perhaps it was given on the results of a standardized test, or maybe you have had an instructor who has given the median score for an exam or quiz. The P^{th} percentile tells us the value in the dataset that is larger than P percent of the data.

The P^{th} percentile is the value of a quantitative variable which is greater than P percent of the data.

Class Example 2.2.9 *Three important percentiles*

We have already discussed three important percentiles in class. Use what you know about the following statistics to determine which percentile it represents.

- a) Q1

- b) Median

- c) Q3

2.3 Outliers, Boxplots, and Quantitative/Categorical Relationships

Key Concepts

- Use a boxplot to describe data for a single quantitative variable
- Use a side-by-side graph to visualize a relationship between quantitative and categorical variables
- Examine a relationship between quantitative and categorical variables using comparative summary statistics

Another display for quantitative data

So far in this class, we have used histograms and dot plots to represent data graphically. There is another very commonly used graphical summary called the *boxplot*. The boxplot gives us a graphical display of the *five-number summary* which consists of the minimum, Q_1 , median, Q_3 , and the maximum of a dataset.

The *boxplot* is a graphical display of the *five-number summary* which consists of the minimum, Q_1 , median, Q_3 , and the maximum of a dataset.

Class Example 2.3.1 *Population of U.S. states*

The following data represent the populations of 12 of the 50 U.S. states, in millions of people, in ascending order.

0.506	0.830	1.315	1.813	2.333	2.953
3.524	4.198	5.504	6.207	8.685	22.472

- a) Identify the five-number summary for the state populations.
-
- b) Draw a boxplot for the data. If there are any outliers, provide a justification as to why they are outliers and identify them on your boxplot.
-


One quantitative and one categorical variable

Sometimes considering the relationship between categorical and quantitative variables, it is helpful to plot them side-by-side.

We use *side-by-side* plots to visualize the relationship between a quantitative variable and a categorical variable.

Class Example 2.3.2 *Average household incomes broken down by U.S. geographical region*

The figure below shows a side-by-side boxplot of average U.S. household income broken down by geographical region of the U.S.




a) Use these side-by-side boxplots to discuss how the average U.S. household income differs by region.

b) Are there any outliers? If so, how many and estimate the values of the outliers.

c) For which region(s) would it be more appropriate to use the median instead of the mean to describe the average U.S. household income? Justify your answer.

Class Example 2.3.3 *Gross state product*

The figure below gives a boxplot of the Gross State Product for the 50 US states.



a) Are there any outliers? If so, how many and estimate the values of the outliers.

b) Estimate the median, IQR and range of the data.

c) How would you describe the shape of this distribution?

d) Do you expect the mean to be less than, approximately equal to, or greater than the median? Explain.

3.1 Sampling Distributions

So far in the class, we have been focusing on how to summarize a single dataset. However, that is only a very small part of the study of statistics. The real power of statistics comes from knowing that when we have a representative sample, the statistic we calculate from our sample will be close to the parameter we are trying to estimate in the population.

Key Concepts

- Distinguish between a population parameter and a sample statistic, recognizing that a parameter is fixed while a statistic varies from sample to sample
- Compute a point estimate for a parameter using an appropriate statistic from a sample
- Recognize that a sampling distribution shows how sample statistics tend to vary
- Recognize that statistics from random samples tend to be centered at the population parameter
- Estimate the standard error of a statistic from its sampling distribution
- Explain how sample size affects a sampling distribution

Review of statistical inference

In section 1.2, *statistical inference* was defined as the process of using a *statistic* calculated from a sample to estimate a *parameter*, or unknown value, from the population. The reason this cycle works is that when we gather data properly (i.e. use a random sampling technique), the statistics behave in a predictable way. Sample statistics are rarely, if ever, the exact population parameter, but they are usually close. The sampling distribution for a statistic gives us a way to see how close we can expect statistics to be to the true parameter value. Recall the cycle of statistical inference given in the figure below from the notes in Section 1.1.

Statistical inference is using the information in a sample to draw conclusions about the entire population.

A *parameter* is a number that describes some aspect of the population.

A *statistic* is a number that is computed from the data in a sample.

Class Example 3.1.1 Parameter vs. Statistic

For each of the following, determine whether the quantity described is a parameter or a statistic and give the proper notation.

- Average height of players on the 2024 Brazil World Cup team, using data from all 23 players on the roster.
- Average enrollment in charter schools in Illinois. In 2024, there were 147 charter schools in the state of Illinois and the total number of students attending the charter schools was 65,215.
- Proportion of households in the U.S. who are renting their living accommodations using data from the 2020 Census.
- Proportion of UWL students who live on campus, using data from 30 students.

*Statistics and their variability***Class Example 3.1.2** *Illustrating sampling distributions*

The goal of this activity is to demonstrate the variability of sample statistics from sample to sample. Before we get started, we are going to come up with 2 questions that everyone in the class will need to answer. After we have decided on our questions, please write your answers in the blanks below.

Student #: _____

Question 1 (quantitative): _____ Question 2 (Y/N): _____

- a) Pull out a sheet of paper.
- b) Go to <http://random.org> and generate a sample of 15 students. Each student should only appear one time in your sample, so if you get a duplicate number, generate a new one.
- c) Walk around the room to gather the data for each student. Be sure to leave your answers at your desk.
- d) Calculate the average from your sample for question 1 and the proportion from your sample for question 2 and plot your statistics in the dot plots on the board:
 - Question 1: $\bar{x} =$ _____
 - Question 2: $\hat{p}_{\text{yes}} =$ _____

To get a bigger picture of the sampling distribution for a statistic, we need to repeat this procedure thousands of times. To speed up the process, we can do this using a computer.

Sampling distributions

Sample-to-sample variability

The *sampling distribution* shows us how much we can expect statistics to vary from sample to sample. In today's activity, you took samples of 15 students and then calculated a summary statistic (average or proportion) from each of those samples. Then you plotted your statistic on the board. These dot plots on the board represent the sampling distribution and they give you a good idea of the variability you should expect to see from sample to sample.

The *sampling distribution* shows how a statistic will vary from sample to sample.

We use the sample statistic as a *point estimate* for the population parameter.

Facts about sampling distributions

If our samples meet a few conditions, then the sampling distribution of the statistics we calculate should be symmetric and centered at the value of the unknown population parameter. The conditions are

- The sample size should be large enough
- The sample needs to be representative (i.e. a random sample)

Sampling distributions are generally *symmetric*.

Sampling distributions are usually *centered at the value of the population parameter*.

Sampling distributions tell us useful information about which sample statistics are likely to occur. Sample statistics that fall near the center of the sampling distribution are more likely to occur than those that fall far from the center. Understanding which values are likely is one of the primary goals in Statistics.

Class Example 3.1.3 *Variability of statistics*

Consider the samples you took at the beginning of class. Where do the sample statistics you calculated fall in the sampling distributions on the board? Are any of your statistics unusual in the sampling distribution?


The importance of sample size

When gathering data, there is a tradeoff between how large a sample is and the accuracy of the statistic calculated from the sample. This happens because with a larger sample size, we get a more complete picture of what the population looks like. So, as the sample size increases, we are more likely to see sample statistics that are closer to the true value of the population parameter.

As the sample size *increases* the standard error of a statistic will *decrease*.

Class Example 3.1.4 *Average pH levels in Florida lakes*

The figure below shows two sampling distributions for the average pH level in Florida lakes for two different sample sizes.



- a) The difference between these two dotplots is that one comes from samples where $n = 10$ and the other comes from samples where $n = 20$. Identify which dotplot corresponds to each sample size.
 $n = 10$ Top Bottom $n = 20$ Top Bottom
- b) What does each dot in the dotplots represent?

c) For which sample size are we more likely to find an average pH greater than 7?
 Circle one: $n = 10$ $n = 20$

Class Example 3.1.5 Kim's lucky coin

Kim's uncle has given her a *lucky* coin for her 21st birthday that he claims comes up heads more often than tails. She brings the coin to her statistics class to help her determine whether or not the coin is lucky. The class decides that if she can flip more than 65% heads, the coin is *lucky*. Zach proposes that they flip the coin 30 times, but Michelle says that they should do 100 just to be certain.

- a) If we assume the coin is not *lucky*, what percentage of the time should we see heads?
- b) If the coin is not *lucky*, who is more likely, Zach or Michelle, to find a sample that falsely suggests the coin is *lucky*? Explain.

Estimating standard deviation

The standard error of a statistic is very important because it gives us a sense of how certain we are that our statistic is accurate, so it is important that we can easily estimate the standard deviation of the sampling distribution. When we use a computer to generate a sampling distribution, it is very easy to have the computer calculate the standard deviation of the sampling distribution. However, there is also a graphical technique we can use to estimate the standard deviation of a distribution.

Using some ideas from calculus, we can say the first standard deviation on either side of the mean occurs at the *inflection points*. Essentially, when tracing along the distribution from left to right, when your pen stops making an up-cup (a cup that can hold water) and starts making a down-cup (cannot hold water), that occurs approximately one standard deviation from the mean. The same occurs on the other side of the mean.

In a sampling distribution the *standard error* is the standard deviation of the statistic.

An *inflection point* is a point on a distribution where the up-cup becomes the down-cup.

Words of caution

The statistics calculated from a sample are only reliable if we collect our data *correctly* and have a *representative* sample. When we do not use proper techniques to gather the data, then the statistics we calculate will be biased.

3.2 Confidence Intervals

In the last section we discussed using a sample statistic as our best guess for a point estimate for the population parameter. We also talked about the kind of variability we would expect to see in summary statistics from different samples. In this section we will use the standard error along with the point estimate to give a range of plausible value for a population parameter.

Key Concepts

- Construct a confidence interval for a parameter based on a point estimate and a margin of error
- Use a confidence interval to recognize plausible values of the population parameter
- Construct a confidence interval for a parameter given a point estimate and an estimate of the standard error
- Interpret (in context) what a confidence interval says about a population parameter

Interval Estimates and Margin of Error

From our discussion about sampling distributions, recall that sample statistics vary from sample to sample. Any time we give an estimate for a population parameter, we want to describe that variability. This is commonly done using an *interval estimate* or a range of values that are plausible for the population parameter. One common form of an interval estimate is

$$\text{point estimate} \pm \text{margin of error}$$

The *margin of error* reflects the precision of our statistic as a point estimate, and it indicates how far out we should go to obtain a range of plausible values for the population parameter.

An *interval estimate* gives a range of plausible values for a population parameter, commonly formed using *point estimate* $\pm$ *margin of error*.

Margin of error is a number that reflects the precision of the sample statistic as a point estimate for the parameter.

Class Example 3.2.1 Moon Landing Conspiracy

A survey of 500 American adults conducted in 2019 stated that 1 in 10 U.S. adults believe the moon landing was fake. Suppose the survey went on to claim it had “a margin of error of ± 3 percentage points with a 95% level of confidence”.

- a) What is the relevant sample statistic? Give appropriate notation and the value of the statistic.
-
- b) What population parameter are we estimating with this sample statistic?
-
- c) Use the margin of error to give a confidence interval for the estimate. Is 0.15 a plausible value of the population proportion? Is 0.30 a plausible value?
-
- d) If we took a sample of 100 instead, do you expect the margin of error to increase or decrease?

Confidence intervals

Typically in statistics, we refer to our interval estimates as *confidence intervals*. A confidence interval is an interval estimate along with a *confidence level* that tells us how successful our intervals would be at capturing the true (but unknown) parameter if we repeated our sampling thousands of times. One of the more common confidence intervals is the 95% confidence interval, and we can construct a 95% confidence interval using

$$\text{statistic} \pm 2 \cdot \text{standard error}.$$

A *confidence interval* for a parameter is an interval computed from sample data by a method that will capture the parameter for a specified proportion of all samples.

The success rate (proportion of all samples whose intervals contain the parameter) is known as the *confidence level*.

Using the standard error, the 95% *confidence interval* for a sampling

Class Example 3.2.2 Constructing confidence intervals

For each of the following, use the information to construct a 95% confidence interval and give notation for the quantity being estimated.

- a) $\bar{x} = 27$ with standard error 3.2.

Interval: Notation (circle one): $p, \mu, p_1 - p_2, \mu_1 - \mu_2$

- b) $\hat{p}_1 - \hat{p}_2 = 0.05$ with margin of error for 95% confidence of 0.02.

Interval: Notation (circle one): $p, \mu, p_1 - p_2, \mu_1 - \mu_2$

Recap of precision

Before learning about the interpretation of confidence intervals, it is important to highlight some differences between the measures of precision we have discussed thus far.

- The *margin of error* is the amount we add and subtract in a confidence interval.
- The *standard error* is the standard deviation of the sampling distribution of the statistic.
- The *sample standard deviation* represents the average amount a single observation in our sample is from the sample average.

For a 95% confidence interval: $MOE = 2 \cdot SE$

Understanding confidence intervals

Confidence intervals appear all over the place. These intervals appear in research articles and academic publications, but they also appear in mainstream news sources as well. For example, during almost any election cycle, it is almost impossible to find any article about the upcoming vote that does not cite some kind of poll with an associated interval estimate.

Class Example 3.2.3 *Have you ever been arrested?*

According to a recent study of 7335 young people in the US, 30% had been arrested for a crime other than a traffic violation by the age of 23. Crimes included such things as vandalism, underage drinking, drunken driving, shoplifting, and drug possession.

- a) Is the 30%, that the study reports, a parameter or statistic?

- b) The margin of error is 0.01 for a 95% confidence interval. Use this information to give a range of plausible values for the parameter.

- c) A correctional facility network claims that the true proportion of young people who had been arrested for a crime other than a traffic violation is less than 25%. Given the margin of error in part b), if we asked *all* young people in the US if they have ever been arrested, is it likely that the actual proportion is less than 25%?

Meaning of confidence

When we give a confidence level for an interval, it indicates how confident we are in the procedure we used to gather the sample. For example, when we give a 95% confidence interval, it means that if we were to repeat our sampling procedure many times, approximately 95% of the resulting intervals should capture the true population parameter.

The *confidence level* tells us how many times we should capture the true population parameter if we repeat the process many times.

Class Example 3.2.4 *Interpreting confidence level*

For each of the following indicate the number of confidence intervals that should capture the true population parameter.

- a) We repeat our sampling 30 times and calculate 95% confidence intervals each time.

- b) We repeat our sampling 100 times and calculate 99% confidence intervals each time.

Interpretation of confidence interval

We can interpret a 95% confidence interval by saying that we are 95% confident that the parameter of interest is in the interval. It is important to note that the confidence interval only refers to the *population parameter*, it has nothing to do with individuals

in the population or for the statistics calculated from other samples.

Class Example 3.2.5 *Fish democracies*

In a school of fish, it is believed that a strongly opinionated minority can influence the path of the majority of the fish. In a study on fish democracies, the researchers found a 95% confidence interval for the proportion of times the minority win to be 0.74 to 0.91.

a) Provide an interpretation of the confidence interval in the context of the study.

b) What is the margin of error for this confidence interval.

c) Is it plausible to believe that the opinionated minority do impact the path of the majority?

Class Example 3.2.6 *Biomass in tropical forests*

Using a sample of 4079 inventory plots, scientists give a 95% confidence interval of 9600 to 13600 tons for the mean amount of carbon per square kilometer in tropical forests.

a) What is wrong with this interpretation? "We believe that 95% of tropical forests will have an average of carbon per square kilometer that is between 9600 to 13600 tons."

b) Provide a correct interpretation of the confidence interval in the context of the study.

Class Example 3.2.7 *Tipping for pizza delivery*

Using a sample of 24 deliveries described in “Diary of a Pizza Girl”, we find a 95% confidence interval for the mean tip given for a pizza delivery to be \$4.18 to \$5.90. Which of the following is a correct interpretation of this interval? Circle the correct interpretation(s).

- a) I am 95% sure that all pizza delivery tips will be between \$4.18 and \$5.90.
- b) 95% of all pizza delivery tips will be between \$4.18 and \$5.90.
- c) I am 95% sure that the mean pizza delivery tip for this sample will be between \$4.18 and \$5.90.
- d) I am 95% sure that the mean tip for all pizza deliveries in this area will be between \$4.18 and \$5.90.
- e) I am 95% sure that the confidence interval for the mean pizza delivery tip will be between \$4.18 and \$5.90.

3.3 Constructing Bootstrap Confidence Intervals

We’ve talked about how to construct a confidence interval if we know the standard error of the statistic or the margin of error. Now we’re going to talk about how we can use a technique called bootstrapping to come up with an estimate for the standard error.

Key Concepts

- Recognize that a bootstrap distribution tends to be centered at the value of the original statistic
- Use technology to create a bootstrap distribution
- Estimate the standard error of a statistic from a bootstrap distribution
- Construct a 95% confidence interval for a parameter based on a sample statistic and the standard error from a bootstrap distribution

Orange and green M&M’S

We want to learn about the true proportion of M&M’S that are green and orange. In section 3.2, we introduced the idea of an interval estimate to give a range of plausible population parameters. Recall that we can easily calculate a 95% confidence interval provided that we know the standard error of our statistic. In today’s activity, we’ll explore two different ways we can estimate the standard error: the sampling distribution of the proportion, and a bootstrap distribution.

- a) Open your bag of M&M’S, and pull out 10 pieces of candy. Using your sample, find the proportion of the M&M’S that are either green or orange (i.e. count the number that are green or orange and divide by 10).

Proportion: _____

- b) On the board, we have a single sample of 10 M&M’S.
 - i) If you were to take a bootstrap sample from these M&M’s, how many M&M’s should be in the bootstrap sample?
-
- ii) Go to <http://www.random.org> to generate a sample of 10 numbers between 1 and 10. Some of the numbers may repeat, and that’s ok. These will represent the numbers of the M&M’s you should include in your bootstrap sample.
 - iii) For your bootstrap sample, find the proportion of M&M’S that are either green or orange.

Proportion: _____

We're going to use StatKey (<http://www.lock5stat.com/statkey>) to plot these distributions and estimate the standard error for the proportion. Use these standard errors to find 95% confidence intervals for the actual proportion of green and orange M&M'S.

SE from a): _____ SE from b): _____

Bootstrap distributions

We can use the example from the beginning of class to demonstrate the differences between a sampling distribution and a bootstrap distribution. In part a), we considered the first 10 M&M'S in your bag to be a sample, and every student had their own individual sample. Then we used the proportions from everyone's sample to build up the sampling distribution for the proportion of green and orange M&M'S. For part b), we were using a *bootstrap distribution*. The main distinction between the sampling distribution and the bootstrap distribution is related to the number of samples we need to take from the population to create our distribution. For the sampling distribution, each of you had your own sample. In contrast, for the bootstrap distribution, we think of the original sample as a fairly good representation of the true population, and we take many samples of the same sample size using replacement from our initial sample.

We can create a *bootstrap distribution* using a single representative sample.

Bootstrap sample characteristics

There are two key characteristics that bootstrap samples must satisfy to be a legitimate bootstrap sample.

1. The bootstrap sample must have the same sample size.
2. All observations in the bootstrap sample must be in the original sample.

Class Example 3.3.1 *Can these be a bootstrap sample?*

We have a random sample of 5 exam grades: 85, 72, 79, 97, 88. For the following state whether or not it is a possible bootstrap sample. If it is not possible, provide a correction to the sample.

a) 79, 79, 97, 85, 82

b) 85, 88, 97, 72

c) 85, 85, 85, 85, 85

Creating a bootstrap distribution

The bootstrapping procedure must agree with the method with which the data were collected. For example, when we tried to estimate the standard error for the proportion of M&M'S that were orange or green, we can resample (bootstrap) from the single sample. For a statistic that is a difference (e.g. $\hat{p}_1 - \hat{p}_2$ or $\bar{x}_1 - \bar{x}_2$), the bootstrapping procedure is very similar. For a single bootstrap sample, we would sample with replacement from group 1, sample with replacement from group 2, and then calculate the statistic.

Class Example 3.3.2 *Describe the bootstrap procedure*

For each of the following sample results, describe how we could create a bootstrap distribution.

a) We take a sample of $n = 20$ and find a proportion of $\hat{p} = 0.4$.

b) We want to compare male and female sleep time. We take a sample of $n_f = 83$ females and $n_m = 65$ males and find $\bar{x}_f - \bar{x}_m = 0.47$

Class Example 3.3.3 *Tea, coffee, and your immune system*

Researchers suspect that drinking tea might enhance the production of interferon gamma, a molecule that helps the immune system fight bacteria, viruses, and tumors. A classic study (Kamath *et al.*, PNAS 2003) involved 21 people who did not normally drink tea or coffee: eleven were randomly assigned to drink tea, and the other 10 were assigned to drink coffee. After two weeks, blood samples were taken and interferon gamma levels were measured. The data are in the set **ImmuneTea**.

- a) What is the quantity of interest in this study? Is this a parameter or a statistic? Write this parameter using proper notation.

- b) How could we create a bootstrap distribution to estimate the standard error of the quantity in part a)?

- c) Using *StatKey* or other software, give a point estimate for the quantity in a).

- d) Using *StatKey* or other software, give a point estimate for the standard error of the quantity in a).

- e) What was the mean of the bootstrap distribution in part d)?

Bootstrap confidence intervals using standard errors

The primary function of the bootstrapping procedure is to provide a measure of the variability of sample statistics. One of these measures of spread is the *standard deviation* of the bootstrap distribution. The standard deviation of the bootstrap distribution is often a good approximation of the standard error of the sampling distribution (and we don't have to take more samples!). We can construct a 95% confidence interval using the bootstrap SE just as we did with the SE of the sampling distribution:

$$\text{statistic} \pm 2 \cdot \text{SE}_{\text{bootstrap}}$$

The *standard deviation* of the bootstrap distribution provides an estimate of the *standard error* for our statistic.

Class Example 3.3.4 *Skateboard Prices*

In a random sample of 20 skateboard prices on eBay, we find the average price is \$67.59. A bootstrap distribution gives a standard error of 10.9.

a) Find and interpret a 95% confidence interval.

b) Should a 90% confidence interval be wider or narrower than this interval?

c) If we took a sample of 50 prices instead of 20, would the 95% confidence interval be wider or narrower than this interval? Explain.

Class Example 3.3.5 *Tea and coffee, revisited*

Using the results from example 3.3.3, create a 95% confidence interval for the difference in interferon gamma levels between tea and coffee drinkers. What does this confidence interval represent?

Using StatKey for confidence intervals

The main goal of this activity is to have you practice finding confidence intervals using the StatKey technology. Most of the datasets that we will use are accessible through the website for StatKey (<http://www.lock5stat.com/statkey>).

Class Example 3.3.6 *Average Land Area in Countries in the World*

The first dataset we will be using in class is the `TenCountries` dataset available from the StatKey site. This data is a random sample of 10 of the countries selected from the `AllCountries` dataset which contains information about 213 individual countries as determined by the World Bank in 2008 (or later).

- a) Open the dataset in StatKey, and generate a bootstrap distribution of the average land area (in 1000 sq km) using 3000 bootstrap samples.
 - i. Go to <http://www.lock5stat.com/statkey>
 - ii. Click on the link for “CI for Single Mean, Median, St.Dev.” in the Bootstrap Confidence Intervals column.
 - iii. From the upper left dropdown menu, select “Ten Countries (Area)”.
 - iv. Click on the “Generate 1000 Samples” button 3 times to generate a bootstrap distribution for the average land area (in 1000 sq km)
 - b) Using the Standard Error method, construct a 95% confidence interval for the average land area for all countries. Provide an interpretation for this interval.
-
- c) Using the percentile method, construct a 90% confidence interval for the average land area for all countries.
 - i. In the bootstrap dotplot, select the box for “Two-tail”
 - ii. In the dotplot, three boxes should appear. In the middle box, click and type “0.90”.
 - iii. StatKey will automatically select the appropriate percentiles to include for the confidence interval and adjust the outer boxes accordingly.
 - iv. If you look under the dotplot, you will see three numbers. The outer two numbers correspond to the appropriate percentiles for the endpoint of your confidence interval. Record your interval below and provide an interpretation.

3.4 Bootstrap Confidence Intervals Using Percentiles

We’ve talked about how to construct a confidence interval if we know the standard error of the statistic or the margin of error. Now we’re going to talk about how we can use a technique called bootstrapping to come up with an estimate for the standard error.

Key Concepts

- Construct a confidence interval based on the percentiles of a bootstrap distribution
- Explain how the width of an interval is affected by the desired level of confidence and the sample size
- Recognize when it is appropriate to construct a bootstrap confidence interval using percentiles or the standard error

Bootstrap confidence intervals using percentiles

If the bootstrap distribution is skewed, or does not follow a nice bell-shape, it can cause the confidence interval to be inaccurate. We may also be interested in using a different confidence level. In these cases, constructing a confidence interval using

the standard error rule may give us an interval that is not the right size. We can still use the bootstrap distribution, but instead of using the standard error, we will use the percentiles of the bootstrap distribution for our interval.

Class Example 3.4.1 *Tea and coffee, revisited...again*

In our last class, we created a bootstrap distribution for the difference in the average interferon gamma levels between coffee and tea drinkers.

- What percentiles of the bootstrap distribution would be needed to create a 95% confidence interval?
- Create a 95% confidence interval for the difference in average interferon gamma levels between tea and coffee drinkers using the percentile method.
- Give an interpretation of the interval in part b).
- Create a 95% confidence interval for the difference in average interferon gamma levels between tea and coffee drinkers using the standard error method. Recall from the sample, $\bar{x}_1 - \bar{x}_2 = 17.12$.

Class Example 3.4.2 *Average penalty minutes in the NHL*

We have a random sample of the penalty minutes for $n = 26$ players from the 2018–2019 season for the NHL team the Ottawa Senators. Some percentiles from a bootstrap distribution of 1000 sample means are shown in the table.

	0.5%	1.0%	2.0%	2.5%	5.0%	95.0%	97.5%	98.0%	99.0%	99.5%
Percentile	13.8	14.6	15.4	15.8	17.2	33.1	35.1	35.3	36.4	38.0

Selected percentiles from bootstrap distribution { .striped }

- What percentiles of the bootstrap distribution would be needed to create a 90% confidence interval?
- Create a 90% confidence interval for the average penalty minutes for a player on the Ottawa Senators in the 2018–2019 season using the percentile method.
- What percentiles of the bootstrap distribution would be needed to create a 98% confidence interval?
- Create a 98% confidence interval for the average penalty minutes for a player on the Ottawa Senators in the 2018–2019 season using the percentile method.

e) Give an interpretation for one of the intervals (90% or 98%).


f) Using the 90% interval, is it plausible to say that the Ottawa Senators players were given more than 34 penalty minutes on average per player? Explain.

Class Example 3.4.3 *Manhattan apartment rent*

The New York City housing authority wants to know about the average rent for a one-bedroom apartment in Manhattan. One of their employees goes to Craigslist and records the monthly rent for a randomly selected sample of 8 listed one-bedroom apartments in Manhattan. Her sample is given below

3150 2275 3100 2650 2925 3305 2325 5495

Use the dotplot and the table below to answer the following questions.



	1.0%	2.5%	5.0%	10.0%	90.0%	95.0%	97.5%	99.0%
Percentile:	2595.25	2651.85	2720.40	2788.53	3546.80	3679.48	3796.40	3913.68

Selected percentiles from bootstrap distribution {.striped}

- Write down one possible bootstrap sample from the original sample.
-
- Is it appropriate to use the bootstrap distribution to create a confidence interval for the average rent price in Manhattan? Justify your answer.
-
- Regardless of how you answered b), create and interpret a 95% confidence interval for the average rent in Manhattan using the percentile method.
-

Class Example 3.4.4 *Video Games Down Undah'*

In a study conducted by researchers in Australia (Eshuis *et al.*, Simulation & Gaming, 2023) they analyzed the gender demographics of different video game players. Among the 87 single-player gamers, 28 identified as female, and among 58 multiplayer gamers, 20 identified as female.

- a) What is the point estimate of the difference in proportions of female gamers between single-player and multiplayer gamers?

- b) Create a 95% confidence interval for the difference in proportions of female gamers between single-player and multiplayer gamers using the percentile method.

- c) Create a 95% confidence interval for the difference in proportions of female gamers between single-player and multiplayer gamers using the standard error method.

- d) Give an interpretation of the interval from part b).

- e) Does your interpretation suggest that there is a relationship between game preference and gender? Explain.

Words of caution

You should always plot a bootstrap distribution before using it to construct a confidence interval. If the plot is poorly behaved (e.g heavily skewed or isolated clumps of values), then a bootstrap distribution will not give very accurate confidence intervals.

Confidence Interval Practice Questions

The main goal of the following questions are to give you additional practice with constructing confidence intervals, using confidence intervals to find standard error and margin of error, and interpreting confidence intervals.

Class Example 3.4.5 *Baseball fans*

In a poll taken in March of 2007, Gallup asked 1006 U.S. adults whether they were baseball fans. 362 adults said they were. Gallup reported the 95% confidence interval for the proportion of U.S. adults who were baseball fans as (0.3302, 0.3895).

a) What is the parameter of interest?

b) Find the margin of error for this confidence interval.

c) What is the standard error for the proportion.

d) Give an interpretation of this confidence interval in the context of the study.

Class Example 3.4.6 *Cardiac disease and depression*

A study published in the *Archives of General Psychiatry* in March 2001 examined the impact of depression on a patient's ability to survive cardiac disease. Researchers identified 450 people with cardiac disease, evaluated them for depression, and followed the group for 4 years. Of the 361 patients with no depression, 67 died. Of the 89 patients with minor or major depression, 26 died.

a) Find a point estimate for the difference in the proportion of people with no depression who died from cardiac disease and the proportion of people with minor or major depression who died from cardiac ($\hat{p}_{\text{no depression}} - \hat{p}_{\text{depression}}$).

b) The margin of error for a 95% confidence for this difference in proportions is 0.0513. Find the 95% confidence interval.

c) Give an interpretation of this confidence interval in the context of the study.

Class Example 3.4.7 *Rats in a maze*

Psychology experiments sometimes involve testing the ability of rats to navigate mazes. The mazes are classified

according to difficulty, as measured by the mean length of time it takes rats to find the food at the end. One researcher needs a maze that will take rats an average more than one minute to solve. He tests one maze on 20 different rats, and finds the average time is 50.13 seconds.

- a) What is the parameter of interest for this study?

- b) The standard error for this average is 2.22 seconds. What is the 95% confidence interval for the average time it takes the mice in the sample to finish?

- c) If the researcher had included 200 rats in his study, would the interval be smaller than, larger than, or around the same size as the interval given in b). Explain.

- d) Give an interpretation of this confidence interval in the context of the study.

- e) Does your interval suggest that this maze will be adequate for the researcher's needs? Explain.

Class Example 3.4.8 *Learning to walk*

Researchers designed an experiment to see whether a program of special exercises for 12 minutes each day could speed up the process of learning to walk. In all, 23 infants participated in the study. The study found that average age (in months) for the first unaided steps in the special exercises group was 10.125, and the average age (in months) for the first unaided steps in the control group was 11.375. The 95% confidence interval for the average difference in ages (in months) between the special exercises group and control group was $(-3.44, 0.94)$.

- a) What is the margin of error for this confidence interval?

- b) Give an interpretation of this confidence interval in the context of the study.

- c) Do the study results suggest that the special exercises help infants to take their first unassisted steps more quickly?

Class Example 3.4.9 *Meaning of confidence*

What does it mean when you calculate a 95% confidence interval?

- a. The process you used will capture the true parameter 95% of the time in the long run.
- b. You can be “95% confident” that your interval will include the population parameter.
- c. You can be “5% confident” that your interval will not include the population parameter.
- d. All of the above statements are true.

Class Example 3.4.10 *Standard error*

A good way to get a small standard error is to use a _____

- a. Repeated sampling structure.
- b. Small sample.
- c. Large sample.
- d. Large population.

Class Example 3.4.11 *SAT score Confidence Interval*

You have sampled 25 students to find the mean SAT scores at Morris Knolls High School. A 95% confidence interval for the mean SAT score is 900 to 1100. Which of the following statement(s) gives a valid interpretation of this interval?

- a. 95% of the 25 students have a mean score between 900 and 1100.
- b. 95% of the population of all students at Morris Knolls have a score between 900 and 1100.
- c. We are 95% confident that the mean SAT score for students at Morris Knolls is between 900 and 1100.
- d. If this procedure were repeated many times, 95% of the sample means would be between 900 and 1100.
- e. If 100 samples were taken and a 95% confidence interval was computed, 5 of them would be in the interval from 900 to 1100.
- f. If this procedure were repeated many times, 95% of the resulting confidence intervals would contain the true mean SAT score at Morris Knolls.

Class Example 3.4.12 *Margin of error*

A survey was conducted to determine the percentage of college freshmen that planned to take a biology course while in college. The results were given as 75% with a margin of error of $\pm 4\%$. What does the margin of error of $\pm 4\%$ mean?

- a. The percentage of the population that was surveyed was between 71% and 79%.
- b. 4% of the population was not included in the survey.
- c. 4% of the students surveyed will most likely change their minds.
- d. The difference between the sample proportion and the population proportion is likely to be less than 4%.
- e. 4% of the students surveyed refused to participate in the poll.

Class Example 3.4.13 *Cell phone battery life*

A research and development engineer is preparing a report for the board of directors on the battery life of a new cell phone they have produced. At a 95% confidence level, he has found that the battery life is 3.2 ± 1.0 days. He wants to adjust his findings so the margin of error is as small as possible. Which of the following will produce the smallest margin of error?

- a. Increase the confidence level to 100%. This will assure that there is no margin of error.
- b. Increase the confidence level to 99%.
- c. Decrease the confidence level to 90%.
- d. Take a new sample from the population using the exact same sample size.
- e. Take a new sample from the population using a smaller sample size.

Class Example 3.4.14 *Confidence interval for an average*

A 98% confidence interval for the mean of a large population is found to be 978 ± 25 . Which of the following is true?

- a. 98% of all observations in the population fall between 953 and 1003.
- b. The probability of randomly selecting an observation between 953 and 1003 from the population is 0.98.
- c. If the true population mean is 950, then this sample mean of 978 would be unlikely to occur.
- d. If the true population mean is 990, then this sample mean of 978 would be unexpected.
- e. If the true population mean is 1006, then this confidence interval must have been calculated incorrectly.

Class Example 3.4.15 *Confidence interval width*

After constructing a confidence interval estimate for a population mean, you believe that the interval is useless because it is too wide. In order to correct this problem, you need to:

- a. increase the population standard deviation.
- b. increase the sample size.
- c. increase the level of confidence.
- d. increase the sample mean.

Class Example 3.4.16 *Understanding the confidence interval*

Which statement is not true about confidence intervals?

- a. A confidence interval is an interval of values computed from sample data that is likely to include the true population value.
- b. An approximate formula for a 95% confidence interval is: sample estimate $\pm$ margin of error.
- c. A confidence interval between 20% and 40% means that the population proportion lies between 20% and 40%.
- d. A 99% confidence interval procedure has a higher probability of producing intervals that will include the population parameter than a 95% confidence interval procedure.

4.1 Introducing Hypothesis Tests

Interval estimates provide us with a way to describe a range of plausible values. However, many times in research studies, the goal is to use data to support a research claim. In this section, we will start learning about hypothesis tests. Hypothesis tests are statistical tests that allow us to assess claims about a population using data.

Key Concepts

- Recognize when and why statistical tests are needed
- Specify null and alternative hypotheses based on a question of interest, defining relevant parameters
- Demonstrate an understanding of the concept of statistical significance

Motivation

To begin our discussion about hypothesis tests, think about a standard deck of playing cards.

- a) If we were to take a sample of 20 cards, how many would you expect to be red?

- b) Rewrite your answer in a) as a proportion using the proper notation.

In chapter 3, we introduced the concept of a sampling distribution to describe how we expect statistics to vary from sample to sample.

- c) How large of a sample proportion (in a sample of 20) would you need to convince yourself that the actual proportion of red cards is higher than it should be?

- d) How could we use technology to check and see if the cutoff you listed in c) is reasonable?
-

Hypothesis Tests

In the US justice system, defendants are presumed innocent until proven guilty. Statistical tests, and in particular hypothesis tests, behave in much the same way. These tests compare two competing hypotheses about a population, with one hypothesis claiming “no effect” (the *null hypothesis*, reflecting the “innocent defendant”), and the other claiming an effect (the *alternative hypothesis*, reflecting the “guilty verdict”).

The first set of statistical hypotheses that we will investigate in this class are for population parameters and differences in population parameters. The hypotheses have the following forms:

1. Single Proportion (p_0 is the proportion claimed or status quo value):

- Null hypothesis:

$$H_0 : p = p_0$$

- Alternative hypothesis (pick one):

$$H_a : p \neq p_0$$

$$H_a : p > p_0$$

$$H_a : p < p_0$$

2. Single Mean (μ_0 is the average claimed or status quo value):

- Null hypothesis:

$$H_0 : \mu = \mu_0$$

- Alternative hypothesis (pick one):

$$H_a : \mu \neq \mu_0$$

$$H_a : \mu > \mu_0$$

$$H_a : \mu < \mu_0$$

A *statistical test* uses data from a sample to assess a claim about a population.

Usually, a *null hypothesis*, notated H_0 , is a claim that there really is “no effect” or “no difference”, or represents the status quo.

An *alternative hypothesis*, notated H_a , is the claim for which we seek significant evidence, and is established by observing evidence that contradicts the null hypothesis and supports the alternative hypothesis.

When writing a statistical hypothesis, you should always use the notation for the population (i.e. $p, \mu, p_1 - p_2$, and $\mu_1 - \mu_2$).

3. Difference in Proportions (p_1 and p_2 are the population proportions for each group):

- Null hypothesis:

$$H_0 : p_1 - p_2 = 0$$

- Alternative hypothesis (pick one):

$$H_a : p_1 - p_2 \neq 0$$

$$H_a : p_1 - p_2 > 0$$

$$H_a : p_1 - p_2 < 0$$

4. Difference in Means (μ_1 and μ_2 are the population averages for each group):

- Null hypothesis:

$$H_0 : \mu_1 - \mu_2 = 0$$

- Alternative hypothesis (pick one):

$$H_a : \mu_1 - \mu_2 \neq 0$$

$$H_a : \mu_1 - \mu_2 > 0$$

$$H_a : \mu_1 - \mu_2 < 0$$

Alternative hypotheses containing “>” or “<” are called *one-sided* alternatives and alternative hypotheses containing “≠” are referred to as *two-sided* alternatives. One-sided alternatives specify a particular direction for the alternative, while two-sided alternatives do not.

Class Example 4.1.1 *Defining statistical hypotheses*

For each of the following scenarios, state the null and alternative hypotheses with the relevant parameter(s).

- a) We are testing to see if the average height of males is different than the height of females.

- b) We are testing to see if the average height of males is greater than the height of females.

- c) We are testing to see if a particular coin is weighted (i.e., not fair).

- d) We are testing to see if the average rent in La Crosse is less than \$950.

Statistical Significance

Statistical tests begin under the assumption that the null hypothesis, H_0 , is true. In order to reject H_0 , sufficient evidence to its contrary must be collected – just like in order to convict a defendant, sufficient evidence must be presented to the judge and/or jury. What exactly is sufficient evidence? In our courtroom example, evidence must be provided so that decisions can be made “beyond a reasonable doubt.” In statistical tests, the data results must be unrealistic assuming H_0 is true. This means that we have found evidence to support H_a .

If sample data strongly support the alternative hypothesis, that is, it is unlikely to occur by chance when the null hypothesis is true, we will reject the null hypothesis in favor of the alternative hypothesis.

If we reject the null hypothesis, we say our finding is *statistically significant*.

Class Example 4.1.2 *Great Lakes Water Temperature*

For a random sample of 100 days in 2023, the National Oceanic and Atmosphere Administration recorded the surface temperature for Lake Michigan and Lake Superior. We are interested in determining if there is a difference in the average temperature between the two lakes.

a) Define the relevant parameter(s) and state the null and alternative hypotheses.

b) What statistic(s) from the sample would we use to estimate the difference?

c) This study found that the difference in average water temperature between the lakes to be 7.04F, with Lake Michigan being warmer. Is there statistically significant evidence that Lake Michigan is warmer than Lake Superior? What if there was a mistake, and the actual difference in average temperatures was 15.0F?

Class Example 4.1.3 *Camping in the WI State Parks*

Do Wisconsinites book their camping reservations in advance? In 2023, the Wisconsin Department of Natural Resources reported that at Devil's Lake State Park 21,780 of 35,790 reservations were booked in advance, but at Peninsula State Park 19,282 of 42,593 were booked in advance.

- a) Define the relevant parameter(s) and state the null and alternative hypotheses.

- b) What kind of results would we need in order to conclude that, across all years of camping reservation for both parks (the population), campers at Devil's Lake State Park are more likely to book their reservation in advance?

- c) To assess statistical significance, we assume the null hypothesis is true. What does that mean in this case, in terms of reservations and state parks?

- d) Using this data, the results are said to be statistically significant. What do these results imply about reservations and state parks?

Class Example 4.1.4 *Influencing Voters*

When getting voters to support a candidate in an election, is there a difference between a recorded phone call from the candidate or a flier about the candidate sent through the mail? A sample of 500 voters is randomly divided into two groups of 250 each, with one group getting the phone call and one group getting the flier. The voters are then contacted to see if they plan to vote for the candidate in question. We wish to see if there is evidence that the proportions of support are different between the two methods of campaigning.

- a) Define the relevant parameter(s) and state the null and alternative hypotheses.

- b) Suppose that of the 250 who received a phone call, 152 would vote for the candidate, and that of the 250 who received a flier, 145 would vote for the candidate. Is there a significant difference in the sample proportions?

- c) Suppose that of the 250 who received a phone call, 188 would vote for the candidate, and that of the 250 who received a flier, 120 would vote for the candidate. Is there more evidence of a difference in the sample proportions than in part b)? Why?

Words of caution

Statistical significance does not always imply practical significance. In other words, just because we may find the results of a study to be statistically significant, we cannot necessarily imply that the differences *actually matter*. For example, suppose that an SAT prep company advertises that students that use their product, on average, score 23 points (out of 2400) higher than students who do not use their product (and suppose these results are statistically significant). Is this difference worth the costs associated with the product (often in the range of \$500)?

4.2 Measuring Evidence with P-values

Now that we have talked about how to write hypotheses for a hypothesis test, we need a way to measure the strength of our evidence. In statistics, the *p-value* tells us how likely our evidence is if the null hypothesis is true.

Key Concepts

- Interpret a *p-value* as the probability of results as extreme as the observed results happening by random chance, if the null hypothesis is true
- Estimate a *p-value* from a randomization distribution
- Comparatively quantify strength of evidence using *p-values*
- Distinguish between one-tailed and two-tailed tests in estimating *p-values*

Demonstrating evidence for the alternative hypothesis

Recall from section 4.1, every hypothesis test starts by determining the null and alternative hypothesis. The null hypothesis often represents what we would see if there was *no effect* whereas the alternative hypothesis is what we are trying to show the data supports. Our goal in statistical inference is to determine whether we have sufficient evidence to reject the null hypothesis. In order to reject the null hypothesis and indicate the strength of evidence for the alternative hypothesis, researchers use *p-values*.

When we claim that we have evidence of our alternative hypothesis, the underlying question we are asking is

How unusual is it to see a sample statistic as extreme as the sample statistic we observed, if H_0 is true?

In statistics, we use the *p-value* to answer this question.

We use a *randomization distribution* to show us how likely different sample statistics are to occur if H_0 is true. The *p-value* is the proportion of the randomization distribution that is as extreme (or more extreme) as our observed statistic. If the statistic from our sample lies in the tails of the distribution, we have strong evidence against H_0 in support of H_a . The smaller the *p-value*, the more evidence we have to reject the null hypothesis. We will see how these distributions are actually generated in Section 4.4, but for now, we will examine these types of distributions and their relationship to statistical inference using StatKey.

Randomization Distributions and p-values

To find the *p-value* using a randomization distribution, we:

- For a one-sided alternative hypothesis: Find the proportion of randomization samples that equal or exceed the original statistic in the direction (tail) indicated

The *p-value* of the sample data in a statistical test is the probability of obtaining a sample statistic as extreme as (or more extreme than) the observed sample when the null hypothesis is true.

A *randomization distribution* approximates a sampling distribution from a population where the null hypothesis is true.


In order to correctly estimate the *p-value* from a randomization distribution, we need to keep our alternative hypothesis in mind!

by the alternative hypothesis.

- For a two-sided alternative hypothesis: Find the proportion of randomization samples in the smaller tail at or beyond the original statistic and then double the proportion to account for the other tail.

Class Example 4.2.1 Estimating a p -value from a randomization distribution

The figure below shows a randomization distribution for testing $H_0 : p = 0.3$ vs $H_a : p < 0.3$. In each case, use the distribution to decide which value is closer to the p -value for the observed sample proportion. There are 1000 randomizations given in the figure below.




a) Why is the distribution centered at 0.3?

b) The p -value for $\hat{p} = 0.15$ is closest to: 0.04 or 0.40?

c) Which proportion, $\hat{p} = 0.20$ or $\hat{p} = 0.10$, provides the strongest evidence to suggest that H_0 is not true? Explain.

Class Example 4.2.2 Estimating a p -value from a randomization distribution...continued

The figure below shows a randomization distribution for testing $H_0 : \mu_1 - \mu_2 = 0$ vs $H_a : \mu_1 - \mu_2 \neq 0$. The statistic used for each sample is $D = \bar{x}_1 - \bar{x}_2$. In each case, use the distribution to decide which value is closer to the p -value for the observed sample difference in averages. There are 1000 randomizations given in the figure below.



a) The p -value for $D = \bar{x}_1 - \bar{x}_2 = -2.9$ is closest to: 0.01 or 0.25?

b) The p -value for $D = \bar{x}_1 - \bar{x}_2 = 1.2$ is closest to: 0.30 or 0.60?

c) Of the two values of D above, which provides strongest evidence to suggest that H_0 is not true? Explain.

Class Example 4.2.3 *Smiles and Leniency*

Can smiling increase leniency in punishments? LaFrance and Hecht (*Personality and Psychology Bulletin*, 1995, 21:207-214) studied this question using a panel of participants that would judge a student in an academic misconduct hearing. Leniency scores were calculated based on the disciplinary decisions by the judges, who made their decisions based on the context of the cheating and a picture of the accused student (a higher score means the judge was more lenient). The pictures were randomly chosen between a neutral picture and a picture of the student smiling. The full data are in the data set **Smiles**.

a) What are the null and alternative hypotheses for this study?

b) What was the sample statistic for these data?

c) What is the probability that the sample results (or those more extreme) occurred by random chance, assuming that smiling had no effect?

d) How would your answer to c) change if the alternative hypothesis was two-sided?

Class Example 4.2.4 *Support for the death penalty*

In 1980 and again in 2021, a Gallup poll asked a random sample of 1000 US citizens people “Are you in favor of the death penalty for a person convicted of murder?”. In 1980, the proportion saying yes was 0.66. In 2021, it was 0.54. Does this data provide evidence that the proportion of US citizens favoring the death penalty was higher in 1980 than it was in 2021? Using p_1 for the proportion in 1980 and p_2 for the proportion in 2021:

a) State the null and alternative hypotheses

b) What is the sample statistic?

c) To create the randomization distribution, what do we have to assume?

d) What is the p -value for this study?

4.3 Statistical Significance; CIs and Hypothesis Tests

Now that we know how to find a p -value, we need a set of objective rules to determine if an observation is extreme. We will use the significance level to determine if sample statistic is extreme or not. If the p -value is less than the significance level, then the sample statistic is extreme. If the p -value is greater than the significance level, then the sample statistic is not extreme.

Key Concepts

- Make a formal decision in a hypothesis test by comparing a p -value to a given significance level
- State the conclusion to a hypothesis test in context
- Interpret Type I and Type II errors in hypothesis tests
- Recognize a significance level as measuring the tolerable chance of making a Type I error
- Interpret a confidence interval as the plausible values of a parameter that would not be rejected in a two-tailed hypothesis test
- Determine the decision for a two-tailed hypothesis test from an appropriately constructed confidence interval
- Recognize that statistical significance is not always the same as practical significance
- Explain the potential problem with significant results when doing multiple tests

Statistical Decisions

For hypothesis tests, there are two conclusions:

- We *reject* H_0 when the sample statistic is extreme assuming H_0 is true.
- We *do not reject* H_0 when the sample statistic is not too extreme assuming H_0 is true.

Not rejecting the null hypothesis is *not* the same thing as accepting the null hypothesis. The null hypothesis is *never accepted* because the hypothesis test procedure is not designed to *rule in* H_0 , but only to rule it out. That is, evidence is never collected with the idea of favoring or proving the null hypothesis. Sample data are collected for the purpose of looking for evidence *against* H_0 . To arrive at the correct conclusion, the p -value must be small enough to be considered unusual.

When we make a formal decision, we have two options:

- Reject H_0
- Do not reject H_0

How small is small enough?

In section 4.2 we discussed how small p -values mean that a statistic is unlikely if H_0 is true. Now we need to think about some rules to determine when our p -value is small enough to say something is unusual assuming the null hypothesis is true.

The *significance level*, notated α , is the point at which a p -value is considered unusual. Some commonly used significance levels are $\alpha = 0.05$, $\alpha = 0.01$, or $\alpha = 0.10$. Any p -value that is smaller than the significance level is considered unusual. So, to use a p -value to determine whether or not the results are significant, you should use the following guidelines:

- When the p -value $< \alpha$, reject H_0 .
- When the p -value $\geq \alpha$, do not reject H_0 .

We use the *significance level*, α , to decide the cutoff for our p -value to be significant.

Common values for
 $\alpha : 0.05, 0.01, 0.10$

We reject H_0 when P -value $< \alpha$

We do not reject H_0 when P -value
 $\geq \alpha$

Class Example 4.3.1 Migraines and dementia

There is growing scientific evidence of an association between migraines and dementia. A recent study (*Science Daily*, 2019) investigated how this phenomenon manifests among various demographic subsets of the population. A sample of 7,454 patients (half with migraines, half without) were tracked for ten years to see if they received a dementia diagnosis. For each of the following, use the p -value to make the appropriate conclusion (reject or do not reject) for the hypothesis tests. Use $\alpha = 0.05$ for each of your decisions.

- In a test to see if the proportion of patients who develop dementia is larger among patients with migraines than without. (p -value of $\$ < \0.001).
- In a test to see if the proportion of men who develop dementia is larger among men with migraines than without. (p -value of $\$ = \0.722).
- In a test to see if the proportion of women who develop dementia is larger among women with migraines than without. (p -value of $\$ < \0.001).

Interpreting your results

After you have made a decision to reject H_0 or not to reject H_0 , you need to interpret your conclusion in the context of the study. We always write our interpretation in the context of H_a .

Class Example 4.3.2 Migraines and dementia revisited

For each of the following, provide an interpretation of the conclusion in the context of the original study.

- In a test to see if the proportion of patients who develop dementia is larger among patients with migraines than without. (p -value of $\$ < \0.001).
- In a test to see if the proportion of men who develop dementia is larger among men with migraines than without. (p -value of $\$ = \0.722).

- c) In a test to see if the proportion of women who develop dementia is larger among women with migraines than without. (p -value of <0.001).

Type I and Type II errors

In hypothesis testing, there are two possible errors that can be made: Type I and Type II. They are defined in the table below.

A *Type I error* occurs when we reject H_0 , but it is true (e.g., convict an innocent person).

A *Type II error* occurs when we do not reject H_0 , but it is false (e.g., let a guilty person go free).

	Do not reject H_0	Reject H_0
H_0 is true	No error	Type I error
H_0 is false	Type II error	No error

Type I and Type II error defined

Class Example 4.3.3 Migraines and dementia errors

For the following, identify which kind of error could have occurred (Type I or Type II) using $\alpha = 0.05$.

- a) In a test to see if the proportion of patients who develop dementia is larger among patients with migraines than without. (p -value of <0.001 , *reject H_0*).
- b) In a test to see if the proportion of men who develop dementia is larger among men with migraines than without. (p -value of $=0.722$, *do not reject H_0*).
- c) In a test to see if the proportion of women who develop dementia is larger among women with migraines than without. (p -value of <0.001 , *reject H_0*).

What can cause these errors?

The two types of errors that can be made both involve the hypothesis test leading to the wrong conclusion about the null hypothesis. This can happen because the process of random sampling does not give complete or perfect information about the population, and it is always possible to draw a rare sample just by chance. Recall that we have two options when making statistical inferences: *reject H_0* or *do not reject H_0* . It is important to carefully consider what these decisions could mean.

- Decision: Reject H_0 . The following outcomes are possible:
 - You are making the correct decision; H_0 is not true.
 - You are making a Type I error; the sample statistic was so far from the hypothesized value just by chance.

3. Your sampling process or experiment was poor; the sample statistic is biased.
- Decision: Do not reject H_0
 1. You are making the correct decision; H_0 is true.
 2. You are making a Type II error; the sample statistic was so close from the hypothesized value just by chance.
 3. Your sampling process or experiment was poor; the sample statistic is biased.

Balancing out the errors

There are trade-offs between controlling for Type I and Type II errors, and balancing the risk of these errors is important.

1. Setting α to be very small makes it difficult to make a Type I error, but it also makes it easier to make a Type II error.
2. Setting α to be large increases the making a Type I error, but it also reduces the risk of making a Type II error.
3. We can reduce the chance of a Type II error without impacting the risk of a Type I error by increasing the sample size.
4. Statistical *power* tells us how likely we are to reject H_0 when we should, and it is related to the risk of a Type II error.

α is the probability we make a Type I error

Setting α low makes it easier to make a Type II error

The *power* of a test tells us how likely

Class Example 4.3.4 Sneaky uncle

Suppose that a young boy's favorite uncle offered the boy the following deal each weekend. The uncle would flip a coin, and if it landed heads-up, the uncle would mow the lawn; however, if it landed tails-up, the boy would have to mow the lawn. For the first four weeks in a row, the coin landed tails-up, and the boy had to mow the lawn.

- a) After four weeks of mowing the lawn, the young boy grew suspicious. He suspected that his uncle was flipping an unfair coin. Does he have sufficient evidence? Why or why not?
- b) On the fifth weekend, the coin landed tails yet again. Now does the boy have sufficient evidence?
- c) Should the boy accuse his uncle of cheating? What are the consequences of your choice?

Connection between confidence intervals and hypothesis tests

The fundamental goal of statistical inference is to learn about an unknown population parameter through the use of statistics from a representative sample. When we calculate a confidence interval, we are using the data to suggest plausible values for the parameter. This is called *estimation*. When we conduct a hypothesis test, we are making a comparison of some sort using the null hypothesis as a reference point and seeing if the statistic computed from the sample would be unusual or rare if H_0 were true. In either approach, our findings should be consistent.

- If the null value given in H_0 falls outside of a 95% confidence interval, we should reject H_0 at $\alpha = 0.05$ in a two-tailed test based on the same sample.
- If a parameter value given in H_0 falls inside of a 95% confidence interval, we

do not reject H_0 at $\alpha = 0.05$ in a two-tailed test based on the same sample.

Class Example 4.3.5 *Approval rating for congress*

In a Gallup poll conducted in March 2024, a random sample of $n = 1415$ American adults were asked “Do you approve or disapprove of Congress?” The proportion who said they approve is $\hat{p} = 0.11$, and a 95% confidence interval for Congressional approval is 0.094 to 0.126. If we use a 5% significance level, what is the conclusion if we are:

- a) Testing to see if there is evidence that the Congressional approval rating is different than 26.6%. (This happens to be the average sample approval rating for Congress at the end of 2022)

- b) Testing to see if there is evidence that the Congressional approval rating is different than 10%. (This happens to be the sample approval rating for Congress from one week prior to this poll.)

- c) Testing to see if there is evidence that the Congressional approval rating is different than 38%. (This happens to be the average sample approval rating for the Supreme Court during the same month.)

Practical vs Statistical significance

Just because we reject the null hypothesis, does not mean that our findings will have a true impact. When discussing significance of a finding, we need to keep both the *statistical* and *practical* implications in mind.

A finding is *statistically significant* if we reject the null hypothesis.

A finding is *practically significant* if it will make a difference in real life

Class Example 4.3.6 *Weight loss program*

Suppose that a weight loss company advertises that people using its program lose an average of 8 pounds the first month and that the Federal Trade Commission (the main government agency responsible for truth in advertising) is gathering evidence to see if this advertising claim is accurate. If the FTC finds evidence that the average is less than 8 pounds, the agency will file a lawsuit against the company for false advertising.

- a) What are the null and alternative hypotheses the FTC should use?

- b) Suppose the FTC gathers information from a very large random sample of patrons and finds that the average weight loss during the first month in the program is $\bar{x} = 7.9$ pounds with a p -value for this result of 0.0006. What is the conclusion of the test? Are the results statistically significant?

- c) Do you think the results of the test are practically significant?

4.4 Creating Randomization Distributions

Although randomization is primarily handled by software, it is important to understand how randomization distributions are created.

Key Concepts

- For a given sample and null hypothesis, describe the process of creating a randomization distribution
- Use the hypotheses of a test to identify the expected center of a randomization distribution and determine whether the p -value is found using the left, right, or two tails
- Use technology to create a randomization distribution
- Conduct a hypothesis test for a variety of situations

How to create a randomization distribution

There are two main criteria for a randomization distribution to be reliable. The randomization distribution should be consistent with the null hypothesis, and it should match the randomization scheme used in gathering the data. Although we will primarily use programs like StatKey to do the randomization, it is important to understand what the computer is doing. As a general rule, randomization distributions will be centered at the null value of the parameter being tested.

Randomization distributions should be consistent with the null hypothesis.

Randomization distributions should match the randomization used in gathering the data.

Creating randomization distributions for single proportions

- Consistent with the null hypothesis: Randomization samples should be taken from a population where the proportion matches the proportion specified in the null hypothesis.
- Consistent with the data collection: Generate bootstrap samples that have the same size as the original sample.
- Randomization statistic: $\hat{p}$

Creating randomization distributions for single averages

- Consistent with the null hypothesis: Shift the values in the original sample so the average is equal to the average specified in the null hypothesis.
- Consistent with the data collection: Generate bootstrap samples that have the same size as the original sample from the shifted sample.
- Randomization statistic: $\bar{x}$

Creating randomization distributions for a difference in proportions from an observational study

- Consistent with the null hypothesis: Two randomization samples should be taken from a population where the proportion is the average of the two sample proportions.
- Consistent with the data collection: Generate bootstrap samples for the two groups, matching the original sample sizes.
- Randomization statistic: $\hat{p}_1 - \hat{p}_2$

Creating randomization distributions for a difference in proportions from an experiment

- Consistent with the null hypothesis: Put all observations into a single group.
- Consistent with the data collection: Randomly assign observations to the two groups, matching the original sample sizes.
- Randomization statistic: $\hat{p}_1 - \hat{p}_2$

Creating randomization distributions for a difference in averages from an observational study

- Consistent with the null hypothesis: Shift the values in group two so the average is equal to the average of the group one.
- Consistent with the data collection: Generate bootstrap samples from group one and the shifted group two, matching the original sizes.

- Randomization statistic: $\bar{x}_1 - \bar{x}_2$

Creating randomization distributions for a difference in averages from an experiment

- Consistent with the null hypothesis: Put all observations into a single group.
- Consistent with the data collection: Randomly assign observations to the two groups, matching the original sample sizes.
- Randomization statistic: $\bar{x}_1 - \bar{x}_2$

Class Example 4.4.1 *Support for the death penalty revisited*

Recall the following example from Section 4.2. In 1980 and again in 2010, a Gallup poll asked a random sample of 1000 US citizens people “Are you in favor of the death penalty for a person convicted of murder?”. In 1980, the proportion saying yes was 0.66. In 2010, it was 0.64. Does this data provide evidence that the proportion of US citizens favoring the death penalty was higher in 1980 than it was in 2010?

a) Define the relevant parameter(s) and state the null and alternative hypotheses.

b) Where should the randomization distribution be centered?

c) What statistic is plotted in the randomization distribution?

d) How would you create a randomization distribution to test this hypothesis?

Class Example 4.4.2 *Household Incomes East and West of the Mississippi*

For a random sample of 200 counties US in 2013, we record median household income, and whether the location is east of west of the Mississippi River. We are interested in determining whether there is a difference in average household income between those east of the Mississippi and those west of it. This sample includes 103 counties from the east and 97 counties from the west, with a difference in average median incomes between counties east and counties west of the Mississippi to be \$374 (with west making more).

a) Define the relevant parameter(s) and state the null and alternative hypotheses.

b) Where should the randomization distribution be centered?

c) What statistic is plotted in the randomization distribution?

d) How would you create a randomization distribution to test this hypothesis?

Class Example 4.4.3 *Barn swallow feathers*

In 1986, the nuclear reactor at Chernobyl, in Ukraine, leaked radioactivity, causing concern that the radiation might cause mutations in the genes of humans and animals that would be passed on to offspring. About 2% of

barn swallows have white feathers in places where the plumage is normally blue or red. The white feathers are caused by genetic mutations. In a sample of 266 barn swallows captured around Chernobyl between 1991 and 2000, about 16% had white feathers in places where the plumage is normally blue or red.

a) Define the relevant parameter(s) and state the null and alternative hypotheses.

b) Where should the randomization distribution be centered?

c) What statistic is plotted in the randomization distribution?

d) How would you create a randomization distribution to test this hypothesis?

Class Example 4.4.4 *Battery weight*

Golf is big business in Florida. A supplier of golf cart batteries to major courses around the state was suddenly deluged by complaints that the carts he was servicing were running out of power after about 14 holes of golf instead of lasting the full 18. Weight of the battery is a key factor in performance because the plates inside the battery are made of lead, and a decrease in the weight of batteries is usually a sign of a decrease in the quality. The supplier takes a random sample of 15 batteries, and finds an average weight of 65 pounds. The batteries should be 68 pounds. Should the battery manufacturer be concerned that the quality has gotten worse?

a) Define the relevant parameter(s) and state the null and alternative hypotheses.

b) Where should the randomization distribution be centered?

c) What statistic is plotted in the randomization distribution?

d) How would you create a randomization distribution to test this hypothesis?

Class Example 4.4.5 *Eating disorders and Prozac*

The *Journal of the American Medical Association* reported on an experiment intended to see if the drug Prozac[®] could be used as a treatment for eating disorders. The subjects, 93 women being treated for eating disorders, were randomly divided into two groups with 49 women receiving Prozac and 43 women receiving a placebo. The

women completed a self-report questionnaire at the beginning and end of the study. The average improvement for the women on Prozac was 10, and the average improvement for the women on the placebo was 2. Does this data provide evidence that the average improvement was different between Prozac and the placebo?

a) Define the relevant parameter(s) and state the null and alternative hypotheses.

b) Where should the randomization distribution be centered?

c) What statistic is plotted in the randomization distribution?

d) How would you create a randomization distribution to test this hypothesis?

5.1 Normal Distributions

Sometimes it's easier to use a theoretical model to answer a statistical question of interest. In order to do this, we must first talk about a few properties of density functions for random variables and the normal probability distribution in particular.

Key Concepts

- Estimate probabilities as areas under a density curve
- Recognize how the mean and standard deviation represent to the center and spread of a normal distribution
- Use technology to compute probabilities of intervals for normal distributions
- Use technology to find endpoint(s) of intervals with a specified probability for normal distributions
- Convert in either direction between a general normal distribution, denoted $N(\mu, \sigma)$, and a standard normal distribution, denoted $N(0, 1)$

Density curves


Although density curves come in many different shapes, there are two key characteristics that all density curves have:

- The total area under the curve equals 1 (which corresponds to 100% of the distribution).
- The area under the density curve within any interval is the proportion of the distribution in that interval and, thus, represents the probability that a randomly selected value from that distribution falls in that interval.

A density curve is a graphical representation of the distribution for a continuous random variable (i.e. sample space with probabilities).

Normal distribution

It is not always easy to produce a bootstrap or randomization distribution to answer the statistical question of interest. However, in many circumstances, a theoretical model can be used in place of the randomization distribution. Most of the bootstrap and randomization distributions we have seen in this class have been symmetric and bell-shaped. This type of distribution shape occurs frequently in statistics and is also a characteristic of a family of continuous probability distributions called the *normal distributions*. The normal distribution is density curve that has been well-studied and has a number of very nice properties.



All normal distributions are centered at their mean, μ , and the standard deviation, σ , tells us how spread out the distribution is. We will use the notation $X \sim N(\mu, \sigma)$ to indicate that the random variable X follows a normal distribution with mean, μ , and standard deviation, σ .

The letter Z is used to represent the *standard normal distribution*, denoted $Z \sim N(0, 1)$, which means that Z comes from a normal distribution with mean, $\mu = 0$, and standard deviation, $\sigma = 1$.

The *standard normal distribution* is denoted $N(0, 1)$

Standardizing and Unstandardizing

A *z-score* is a measure of relative standing. Often times, we want to compare two different observations relative to their populations. To do so, it is helpful to first standardize them and then compare the *standard score* or *z-score*. The *z-score* tells us how many standard deviations the observation is from the mean. We can standardize X by using the following formula:

The *standard score* or *z-score* indicates how many standard deviations a particular value is from the mean.

$$Z = \frac{X - \mu}{\sigma}$$

We can unstandardize Z by using the following formula:

$$X = \mu + Z\sigma$$

Class Example 5.1.1 *Standardizing to compare*

For each of the following, compare the results using their standardized score

- Benny and Aaron are on the same track team. They have recently competed at a track and field day, and have each won their respective events. Benny won the high jump with a jump of 75 inches, and Aaron won the 100 m dash with a time of 12 seconds. Let $J \sim N(65, 2)$ be the height of the jumps and $D \sim N(15, 1)$ be the times of the 100 m dash. Who performed better Benny or Aaron?
- A car manufacturer has a small car that gets 36 mpg and a truck that gets 22 mpg. Let $C \sim N(35, 4)$ be the distribution of mpgs for small cars and $T \sim N(19, 3)$ be the mpgs for trucks. Which vehicle performs better relative to the other vehicles in its class?

Class Example 5.1.2 *SAT scores*

Let X be the SAT math scores for science and engineering majors. $X \sim N(609, 80)$

- a) What proportion of SAT scores should be under 569?
- b) What is the probability that a randomly selected student with a science and engineering major has an SAT score of at least 684?
- c) What proportion of SAT scores should be between 500 and 630?

Class Example 5.1.3 *Rainfall in Ithaca*

Let R be the yearly rainfall (in inches) in Ithaca, NY. Data from Cornell's Northeast Regional Climate Center indicates that $R \sim N(35.4, 4.2)$.

- a) How much rain should Ithaca expect to get in the year that is the 99th percentile of rainfall?
- b) How much rain should Ithaca expect to get in the year that marks the top 20 percent of rainfall?
- c) What should the IQR be for yearly rainfall in Ithaca? (*i.e.* the middle 50%)

Class Example 5.1.4 *Furniture assembly*

A company that markets build-it-yourself furniture sells a computer desk that is advertised with the claim "less than an hour to assemble." Let T be the time (in hours) it takes to assemble the furniture. Through

post-purchase surveys, it appears that $T \sim N(1.29, 0.43)$.

a) What percentage of people should be able to assemble the desk in less than an hour?

b) What is the probability that a randomly selected person can assemble the desk in between 1 and 1.5 hours?

c) The company would like to change the advertising claim to read “less than _____ to assemble” so that 90% of people can assemble the desk in the time listed. What time should they use (as the cutoff for the bottom 90%?

6.1 Inference for Single Proportions

Key Concepts

- Find the mean and standard error for a distribution of sample proportions
- Identify when a normal distribution is an appropriate model for a distribution of sample proportions
- Use a normal distribution, when appropriate, to compute a confidence interval for a population proportion
- Use a normal distribution, when appropriate, to test a hypothesis about a population proportion.

Sample statistics

One of the best known theorems in statistics is the *Central Limit Theorem* (CLT) which states that for random samples with a sufficiently large sample size, the distribution of possible sample statistics for a proportion or mean follows a normal distribution that is centered at the value of the population parameter. This theorem is very powerful because it basically says that provided we have a large enough sample size, we can use the normal distribution to find confidence intervals and conduct hypothesis tests without needing to put together a bootstrap or randomization distribution. In chapter 6, we will cover some of the more basic applications of the CLT as it relates to single proportions, single averages, differences in averages, and averages of differences.

Distribution of a single proportion

For categorical data, the goal is often to learn about a population proportion from a sample proportion. We can use the CLT to conduct inference about sample proportions if we have a large enough sample size. If a sample size is large enough, the CLT says that the distribution of all sample proportions ($\hat{p}$) will follow a normal distribution that is centered at the population proportion (p). In chapter 3, we discussed estimating the standard error of a statistic using a bootstrap distribution. When we use the CLT, we can estimate the standard error for a sample proportion by

$$SE(\hat{p}) = \sqrt{\frac{p(1-p)}{n}}$$

where p is the population proportion, and n is the sample size.

Class Example 6.1.1 Use the CLT to estimate the standard error
For each of the following, find $SE(\hat{p})$.

a) Samples of size $n = 100$ from a population with $p = 0.5$

b) Samples of size $n = 200$ from a population with $p = 0.5$

c) Samples of size $n = 500$ from a population with $p = 0.5$

How large do we need?

In order to be able to use the CLT, we need a large enough sample size. For proportions, we need to check that $np \geq 10$ and $n(1 - p) \geq 10$. If the sample size is not large enough, you *cannot* use the CLT to conduct inference.

To use the CLT, we need
 $np \geq 10$ and $n(1 - p) \geq 10$

Class Example 6.1.2 *Describe the sampling distribution*

For each of the following questions, identify if you can use the CLT. If so, fully describe the distribution. If not, identify the smallest sample size you need in order to use the CLT.

- a) Samples of size $n = 10$ from a population with $p = 0.4$.

$\hat{p} \sim N(\text{_____}, \text{_____})$ or Smallest n : _____

- b) Samples of size $n = 100$ from a population with $p = 0.6$.

$\hat{p} \sim N(\text{_____}, \text{_____})$ or Smallest n : _____

- c) Samples of size $n = 40$ from a population with $p = 0.9$.

$\hat{p} \sim N(\text{_____}, \text{_____})$ or Smallest n : _____

CI for single proportion

Provided that the sample size is large enough so that $n\hat{p} \geq 10$ and $n(1 - \hat{p}) \geq 10$, a confidence interval for a population proportion p can be constructed using

We can only use the CLT for confidence intervals if $n\hat{p} \geq 10$ and $n(1 - \hat{p}) \geq 10$

$$\hat{p} \pm z^* \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}$$

where $\hat{p}$ is the sample proportion and z^* is the appropriate confidence multiplier from a $N(0, 1)$ distribution.

*Finding z^**

To find the appropriate z^* for the confidence interval for a population proportion, we need to use technology to find the corresponding endpoints for the confidence interval.

Class Example 6.1.3 *Finding z^**

Find the z^* values for the following confidence intervals.

a) 92%:

b) 85%:

c) 80%:

Some of the more common z^* are given in the table below.

Confidence	z^*
90%	1.645
95%	1.960
99%	2.575

Commonly used z^* values. { .striped }

Class Example 6.1.4 *Left-handed desks*

A university would like to estimate the proportion of its students that are left-handed to purchase new desks for a building renovation. They take a random sample of 300 students and find that 32 people in the sample are left-handed.

- a) Using the CLT find a 95% confidence interval for the proportion of students that are left-handed.

- b) Provide an interpretation for your interval in a).

- c) The Vice Chancellor for facilities decides that 15% of the desks they order should be left-handed. Use the confidence interval in a) to argue whether or not this will be an adequate proportion of left-handed desks.

Class Example 6.1.5 *Moon Landing Conspiracy*

A survey of 500 American adults conducted in 2019 stated that 1 in 10 U.S. adults believe the moon landing was fake.

- a) Using the CLT find a 99% confidence interval for the proportion of American adults who think the moon landing was fake.

- b) Provide an interpretation for your interval in a).

- c) Use the confidence interval in a) to argue whether or not a majority of American adults believe the moon landing was fake.

Class Example 6.1.6 *Fish democracies revisited*

In a school of fish, it is believed that a strongly opinionated minority can influence the path of the majority of the fish. In a study on fish democracies, 165 out of 200 times, the strongly opinionated minority were able to convince the majority to follow their lead.

- a) Using the CLT, find a 90% confidence interval for the proportion of times the strongly opinionated minority

will be able to convince the majority to follow their lead.

b) Provide an interpretation for your interval in a).

c) Use the confidence interval in a) to argue whether or not the opinionated minority can influence the majority.

Hypothesis test for a single proportion

Recall that provided the sample size is large enough, then according to the CLT,

$$\hat{p} \sim N \left(p, \sqrt{\frac{p(1-p)}{n}} \right).$$

We can use this fact to conduct a hypothesis test. As with the hypothesis tests we discussed in chapter 4, there are a specific set of steps to conduct a hypothesis test using the CLT.

Step 1: Check conditions for the test

Check whether both $np_0 \geq 10$ and $n(1-p_0) \geq 10$ are true.

Step 2: State the hypotheses

For a single proportion, the null hypothesis always has the form

$$H_0 : p = p_0$$

where p_0 is a number that represents the proportion claimed or *status quo* value. The alternative hypothesis has one of the following three forms:

$$H_a : p < p_0$$

$$H_a : p > p_0$$

$$H_a : p \neq p_0$$

Step 3: Calculate the test statistic

To conduct the hypothesis test using the CLT, we use a standardized test statistic of

$$z = \frac{\hat{p} - p_0}{\sqrt{\frac{p_0(1-p_0)}{n}}}$$

which follows a $N(0, 1)$ distribution.

We can only use the CLT for a hypothesis test if $np_0 \geq 10$ and $n(1-p_0) \geq 10$

Step 4: Find the p -value

To find the p -value, you must use technology to find what proportion of the standard normal distribution is more extreme than the test statistic (in the direction of the alternative hypothesis). If your alternative is two-sided, then you multiply the one-sided p -value by 2.

Step 5: State your decision and interpret in the context of the hypotheses

As with the randomization hypothesis test, you make your decision based upon the p -value from step 3.

- If $p\text{-value} < \alpha$
 - Decision: *reject* H_0 ,
 - Interpretation: We have significant evidence to suggest that the alternative hypothesis is correct.
- If $p\text{-value} \geq \alpha$
 - Decision: *fail to reject* H_0
 - Interpretation: We do not have significant evidence to suggest that the alternative hypothesis is correct.

You should *always* write your interpretation in the context of the original hypothesis.

How to draw a sketch to illustrate the p -value

Class Example 6.1.7 *Root, Root, Root for the Brewers!*

The Milwaukee Brewers played 81 home games in 2023, and won 49 of those games. If we consider the games played in 2023 as a sample of all Brewers home games, conduct a hypothesis test (using $\alpha = 0.01$) to see if the Brewers win more than half of their home games.

- a) Define the relevant parameter(s) and state the null and alternative hypotheses. Be sure to check any relevant conditions.

- b) What is the test statistic.

- c) Find the p -value, and give a sketch to illustrate your p -value.

- d) State your decision and interpret your findings in the context of the study

Class Example 6.1.8 *Union membership*

The government claimed that in 2005, 12.5% of U.S. workers belonged to unions (*Wall Street Journal*, January 21, 2006). A random sample of 1,445 workers was taken in 2023 by the U.S. Department of Labor to see if membership had decreased. In the sample, 144 of the workers belonged to unions. Conduct a hypothesis test (using $\alpha = 0.01$) to see if membership increased.

- a) Define the relevant parameter(s) and state the null and alternative hypotheses.

- b) What is the test statistic.

- c) Find the p -value, and give a sketch to illustrate your p -value.

- d) State your decision and interpret your findings in the context of the study.

Class Example 6.1.9 *Plato's Republic: Syllable Patterns*

Prose rhythm is characterized as the occurrence of five-syllable sequences in long passages of text. This characterization may be used to assess the similarity among passages of text and sometimes the identity of authors. Syllables are often categorized as long or short. On analyzing Plato's *Republic*, researchers found that 26.1% of the five-syllable sequences contained two short and three long syllables (Wishart and Leach, *Computer Studies of the Humanities and Verbal Behavior*, Vol.3, pp. 90-99). Suppose that Greek archaeologists have found an ancient manuscript dating back to Plato's time (about 427-347 B.C.E.). A random sample of 317 five-syllable sequences from the newly discovered manuscript showed that 61 contained two short and three long syllables. Do the data indicate that the population proportion of this type of five-syllable sequence is different from the text of Plato's *Republic*? You may either use a 95% confidence interval or hypothesis test (use $\alpha = 0.05$) to support your conclusion.

Class Example 6.1.10 *Rabid Raccoons*

In 2019, the United States Department of Agriculture (USDA) Wildlife Services reported that among the 4,305 positive cases of rabies in wild animals, 1,545 of them were found in raccoons. Is there evidence that more than a third of rabid wild animals are raccoons? You may use either a 95% confidence interval or hypothesis test (use $\alpha = 0.05$) to support your conclusion.

6.2 Inference for Single Means

Key Concepts

- Find the mean and standard error for a distribution of sample means
- Recognize when a t -distribution is an appropriate model for a distribution of sample means
- Use a t -distribution, when appropriate, to compute a confidence interval for a population mean
- Use a t -distribution, when appropriate, to test a hypothesis about a population mean

Distribution of a single mean

For quantitative data, the goal is often to learn about a population average from a sample average. We can use the CLT to conduct inference about sample averages if we have a large enough sample size. If sample size is large enough, the CLT says that the distribution of all sample averages ($\bar{x}$) will follow a normal distribution that is centered at the population average (μ). In chapter 3, we discussed estimating the standard error of a statistic using a bootstrap distribution. When we use the CLT, we can estimate the standard error for a sample average by

$$SE(\bar{x}) = \frac{\sigma}{\sqrt{n}}$$

where σ is the population standard deviation, and n is the sample size.

For averages, we almost never know the population standard deviation (σ), so we estimate it by using the sample standard deviation (s). When we use s instead of σ , $\bar{x}$ no longer follows a normal distribution. Instead, $\bar{x}$ follows a t -distribution.

We estimate the standard error of $\bar{x}$ with $\frac{s}{\sqrt{n}}$

The t -distribution

The t -distribution is based on a new parameter called the degrees of freedom (d.f.).

What he found is that provided we have met one of two conditions, then

$$t = \frac{\bar{x} - \mu}{s/\sqrt{n}} \sim t_{n-1},$$

t_{n-1} means a t -distribution with $n - 1$ degrees of freedom

or t follows a t -distribution with $n - 1$ degrees of freedom. There are two possible ways you can justify using the t -distribution for averages.

1. For a large enough sample size ($n \geq 30$), you can use the t -distribution
2. If $n < 30$, then the t -distribution is only a good approximation if the population is approximately normal.

To use the t -distribution, you need either
 $n \geq 30$ or - a population that is approximately normal

Class Example 6.2.1 *Describe the sampling distribution*

For each of the following questions, identify whether or not a t -distribution would be appropriate. If so, identify the number of degrees of freedom that you would use for the t -distribution.

- a) Samples of size $n = 10$, from a population that is approximately normal.
- b) Samples of size $n = 15$, from a population that is heavily skewed to the right.
- c) Samples of size $n = 50$, from a population that is heavily skewed to the left.

As $n \rightarrow \infty$, the t -distribution becomes closer and closer to the standard normal distribution.

CI for a single mean

Provided that the sample size is large enough ($n \geq 30$) or the population is approximately normal, a confidence interval for a population average μ can be constructed using

We can only use the t -distribution for confidence intervals if
 $n \geq 30$ or - the population is approximately normal

$$\bar{x} \pm t^* \frac{s}{\sqrt{n}}$$

where $\bar{x}$ is the sample average and t^* is the appropriate confidence multiplier from a t_{n-1} distribution.

*Finding t^**

To find the appropriate t^* for the confidence interval for a population average, we need to use technology to find the corresponding endpoints for the confidence interval.

Class Example 6.2.2 *Doggo Heart Rates*

The article “Changes in Pulse Rate, Respiratory Rate and Rectal Temperature in Working Dogs before and after Three Different Field Trials” (*Animals (Basel)*, 2020) documents the physiological conditions of nine Search-and-Rescue (SAR) dogs. The pulse rate (beats/min) for each dog was recorded immediately after a simulated search. They found a sample mean and standard deviation of 167.42 and 41.40, respectively.

- a) What is the variable in this study? Is it categorical or quantitative?

- b) Assuming that the distribution of pulse rates is (approximately) normal, what is the 99% confidence interval for the mean pulse rate of SAR dogs during a search? ($t^*_{99} = 3.355$)?

- c) Provide an interpretation for your interval in b).

- d) If we had calculated a 90% confidence interval instead of a 99% confidence interval, would it have been larger or smaller than the 99% confidence interval? Explain.

Class Example 6.2.3 *Dinosaur Diets*

A recent study on the digestibility of Jurassic plants (Howell *et al.*, 2023) found that in four samples of *Equisetum hyemale* (a fern), the digestibility (measured in milliliters of gas per 200 milligrams of plant dry matter) had a sample mean and standard deviation of 58.33 and 3.45, respectively.

- a) What is the variable in this study? Is it categorical or quantitative?

- b) How many degrees of freedom are there?

- c) Assuming that the digestibility of pieces from this fern is (approximately) normal, What is the 98% confidence interval estimate for the average digestibility?

- d) Provide an interpretation for your interval in c).

- e) Is it plausible to suggest that average digestibility is the same as *Marattia attenuata* (another fern) which has a digestibility of 9.52?

Class Example 6.2.4 *Taylor Swift!*

Taylor Swift's *The Eras* tour took the country (and world) by storm in 2023 and 2024. From a survey of 596 attendees, the average expenses for travel, tickets, costumes, etc. was \$1,327.74. Suppose the standard deviation was approximately \$500.

- What is the variable in this study? Is it categorical or quantitative?
- What is the 95% confidence interval estimate for the mean expenditure per attendee?
- Given an interpretation of your interval in b).
- Using the average attendance at a Swift concert of 75,000 people, estimate the average amount spent by all attendees at a single concert.
- Assuming the average sales tax rate is approximately 6%, estimate the tax revenue generated from a single Swift concert.

Hypothesis test for a single mean

Provided that the sample size is large enough ($n \geq 30$) or the population is approximately normal, then according to the CLT

$$t = \frac{\bar{x} - \mu}{s/\sqrt{n}} \sim t_{n-1}.$$

Remember, t_{n-1} means a t distribution with $n - 1$ degrees of freedom

We can use this fact to conduct a hypothesis test. As with the hypothesis test we have discussed thus far, there are a specific set of steps to conduct a hypothesis test

using the CLT.

Step 1: Check conditions for the test

Check whether either $n \geq 30$ or the population is approximately normal.

Step 2: State the hypotheses

For a single average, the null hypothesis always has the form

$$H_0 : \mu = \mu_0$$

where μ_0 is a number that represents the averages claimed or *status quo* value. The alternative hypothesis has one of the following three forms:

$$H_a : \mu < \mu_0$$

$$H_a : \mu > \mu_0$$

$$H_a : \mu \neq \mu_0$$

Step 3: Calculate the test statistic

To conduct the hypothesis test using the CLT, we use a standardized test statistic of

$$t = \frac{\bar{x} - \mu_0}{s/\sqrt{n}}$$

which follows a t_{n-1} distribution.

Step 4: Find the p-value

To find the p -value, you must use technology to find what proportion of the t -distribution is more extreme than the test statistic (in the direction of the alternative hypothesis). If your alternative is two-sided, then you multiply the one-sided p -value by 2.

Step 5: State your decision and interpret in the context of the hypotheses

As with the randomization hypothesis test, you make your decision based upon the p -value from step 3.

- If $p\text{-value} < \alpha$
 - Decision: *reject* H_0 ,
 - Interpretation: We have significant evidence to suggest that the alternative hypothesis is correct.
- If $p\text{-value} \geq \alpha$
 - Decision: *fail to reject* H_0
 - Interpretation: We do not have significant evidence to suggest that the alternative hypothesis is correct.

You should *always* write your interpretation in the context of the original hypothesis.

How to draw a sketch to illustrate the p-value

We can only use the t -distribution for confidence intervals if $n \geq 30$ or the population is approximately normal.

Class Example 6.2.5 *Food Deserts*

Food Deserts are geographical areas that have limited access to affordable and nutritious foods. An analysis of accessibility of grocery stores in the City of Vancouver was published in 2022. Using a sample $n = 4,041$ city blocks, they found the average number of grocery stores within a 15-minute walk was 3.8, with a standard deviation of 3.2. Suppose the city claims that, on average, a resident has access to two grocery stores within a 15-minute walk. Conduct a hypothesis test (using $\alpha = 0.05$) to test the city's claim.

- a) Define the relevant parameter(s) and state the null and alternative hypotheses. Be sure to check any relevant conditions.

- b) What is the test statistic.

- c) Find the p -value and give a sketch to illustrate your p -value.

- d) State your decision and interpret your findings in the context of the study.

Class Example 6.2.6 *First-time home buyers*

In 2022, the National Association of Realtors reported that from a sample of 1,262 first-time home buyers, the average age was 36, and suppose the standard deviation was 4.2 years. Conduct a hypothesis test (using $\alpha = 0.05$) to test whether or not the average age of first-time home buyers was different than 2021, where the average age was 33.

- a) Define the relevant parameter(s) and state the null and alternative hypotheses. Be sure to check any relevant conditions.

- b) What is the test statistic.

- c) Find the p -value and give a sketch to illustrate your p -value.

- d) State your decision and interpret your findings in the context of the study.

Class Example 6.2.7 *Ski Patrol: Avalanches*

Snow avalanches can be a real problem for travelers in the western United States and Canada. A very common type of avalanche is called the slab avalanche. These have been studied extensively by David McClung, a professor of civil engineering at the University of British Columbia. Slab avalanches studied in Canada had an average thickness of 67 (in cm) (*Avalanche Handbook*, by David McClung and P. Schaerer.) The ski patrol at Vail, Colorado, is studying slab avalanches in their region. A random sample of avalanches in spring gave the following thicknesses (in cm).

59	51	76	38	65	54	49	62	68	55	64	67	63	74	65	79
----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----

Assuming that slab thickness has an approximately normal distribution, is there evidence that the slab thickness in the Vail region differs from that in Canada? You may either use a 95% confidence interval or a hypothesis test (use $\alpha = 0.05$) to support your conclusion.


Assessing model conditions

In the examples above, we've either assumed that the population values were normally distributed or had large enough sample sizes to assume normality. These assumptions aren't always apparent (or known) in research studies. It is important to investigate whether normality is reasonable before performing a statistical test. The t -test is *robust* to lack of normality, so when we check this condition we are looking for clear evidence that the distribution is non-normal. If normality is reasonable, we can proceed with a t -test.

There are two primary ways that we can check for normality: histograms/dot plots or quantile-quantile plots. For histograms/dot plots, simply create a histogram/dot plot of the sample data and determine whether normality is reasonable (if the sample data is approximately normal, then normality is reasonable). Quantile-quantile plots, or QQ plots, are commonly seen when using statistical software (such as SPSS). They are created by first ordering the sample data from smallest to largest then comparing these ordered sample data to percentiles of the normal distribution. This information is plotted on an x-y graph (scatterplot) as points. If the resulting graph follows a straight line (that goes through the origin), we can say that the normality assumption is reasonable. The figure below provides some examples of QQ plots and their corresponding histograms.

The t -test is *robust* to lack of normality, but we still need to see if normality is a reasonable assumption.

Create a histogram/dot plot or a QQ plot to assess the normality condition for a t -test.



6.3 Inference for Difference in Proportions

Key Concepts

- Find the mean and standard error for a distribution of differences in sample proportions for two groups
- Recognize when a two-sample z -interval for proportions is the appropriate statistical inference procedure
- Use a standard normal distribution, when appropriate, to compute a confidence interval for the difference in proportions between two populations
- State the conditions for the two-sample z -test for a difference in population proportions
- Recognize when it is appropriate to use the two-sample z -test for a difference in population proportions
- Use a z -distribution, when appropriate, to conduct a hypothesis test for the difference in population proportions

Sampling distribution of a difference in sample proportions

Consider two samples of sizes n_1 and n_2 that come from populations with proportions p_1 and p_2 , respectively. It is a property of sample proportions that the distribution of all differences in sample proportions ($\hat{p}_1$ and $\hat{p}_2$) will be centered at the true population difference in proportions ($p_1 - p_2$). When the samples are *independent* of each other, the standard error of $\hat{p}_1 - \hat{p}_2$ is

$$SE(\hat{p}_1 - \hat{p}_2) = \sqrt{\frac{p_1(1-p_1)}{n_1} + \frac{p_2(1-p_2)}{n_2}}$$

Using statistical notation, the sampling distribution of $\hat{p}_1 - \hat{p}_2$ for “large” samples can be summarized as

$$\hat{p}_1 - \hat{p}_2 \sim N\left(p_1 - p_2, \sqrt{\frac{p_1(1-p_1)}{n_1} + \frac{p_2(1-p_2)}{n_2}}\right)$$

As with single proportions, the population proportions (p_1 and p_2) are rarely ever known (which is why we are estimating them in the first place!), so they are estimated by the corresponding sample proportions $\hat{p}_1$ and $\hat{p}_2$. According to the Central Limit Theorem, when the sample sizes are large enough the sampling distribution of $\hat{p}_1 - \hat{p}_2$ follows a normal distribution.

Class Example 6.3.1 *Proportion of Elderly, Florida and Texas*

From the 2010 US Census, we learn that 17.6% of the residents of Florida are 65 years old or older while 10.5% of the residents of Texas are 65 years old or older. If independent, random samples of 300 Florida residents and 200 Texas residents are selected and the difference in the proportion of elderly residents ($p_{FL} - p_{TX}$) is calculated:

- a) Describe the shape of the sampling distribution of differences in sample proportions.

- b) What is the mean of the distribution of differences in sample proportions.

- c) Find the standard error of the distribution of differences in sample proportions.

Conditions

In order to be able to use the standard normal distribution to construct a CI for a difference in proportions, the following conditions must be satisfied.

1. The two samples must be randomly selected.
2. The two samples must be independent of each other.
3. The samples must be large enough for the sample proportions to be normally distributed; $n_1\hat{p}_1 \geq 10, n_1(1 - \hat{p}_1) \geq 10, n_2\hat{p}_2 \geq 10$, and $n_2(1 - \hat{p}_2) \geq 10$ is a good guideline to follow. This means having at least 10 successes and 10 failures in each sample.

Two-sample z-interval for a difference in population proportions

Provided that the conditions in the previous section are met, the difference between two unknown population proportions can be estimated with a confidence interval as

$$(\hat{p}_1 - \hat{p}_2) \pm z^* \sqrt{\frac{\hat{p}_1(1 - \hat{p}_1)}{n_1} + \frac{\hat{p}_2(1 - \hat{p}_2)}{n_2}}$$

where z^* is the appropriate critical value for the desired level of confidence from a $N(0, 1)$ distribution.

Class Example 6.3.2 *Mobile Connections to Libraries*

In a random sample of 2,252 Americans age 16 and older, 11% of the 1,059 men and 16% of the 1,193 women said they have accessed library services via a mobile device.

- a) What must be *assumed* in order to construct a confidence interval for the difference in proportions?

-
- b) Find and interpret a 95% confidence interval for the difference in proportions accessing libraries via mobile devices, between men and women.
-
- c) Does your interval in (b) suggest that there is a difference in the proportions of men and women accessing libraries via mobile devices? Which population has a higher proportion?
-

Two-sample z-test for a difference in population proportions

To conduct this test, the following conditions must be true:

1. The two samples must be randomly selected,
2. The two samples must be independent of each other,
3. The samples must be large enough for the sample proportions to be normally distributed; having at least 10 successes and 10 failures in each sample is a good guideline to follow. That is, $n_1\hat{p}_1 \geq 10$, $n_1(1 - \hat{p}_1) \geq 10$, $n_2\hat{p}_2 \geq 10$, and $n_2(1 - \hat{p}_2) \geq 10$.

Two unknown population proportions can be compared using a hypothesis test by following the steps below.

Step 1: State the hypotheses

For a difference in averages, the null hypothesis always has the form

$$H_0: p_1 - p_2 = 0$$

The alternative hypothesis has one of the following three forms:

$$H_a: p_1 - p_2 < 0$$

$$H_a: p_1 - p_2 > 0$$

$$H_a: p_1 - p_2 \neq 0$$

$H_0: p_1 - p_2 = 0$ is equivalent to writing $H_0: p_1 = p_2$.

Step 2: Calculate the test statistic

To conduct the hypothesis test, a standardized test statistic of

$$z_0 = \frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\frac{\hat{p}(1-\hat{p})}{n_1} + \frac{\hat{p}(1-\hat{p})}{n_2}}}$$

where

$$\hat{p} = \frac{x_1 + x_2}{n_1 + n_2}.$$

$\hat{p}$ represents the overall proportion of successes under the presumption that H_0 is true. Since $p_1 - p_2 = 0$ implies that $p_1 = p_2$, then the two independent random samples could be seen as coming from one sample of size $n_1 + n_2$. Furthermore, x_1 and x_2 denote the observed number of successes in samples 1 and 2, respectively. The sampling distribution of this test statistic is a standard normal distribution for large samples.

Step 3: Find the P-value

Recall that the P -value is the probability of randomly observing at least as much evidence against H_0 as what is currently observed, given that H_0 is true. This will depend on which H_a is used. Using the $N(0, 1)$ distribution, the following probabilities will be the P -values for each of the three alternatives:

If $H_a: p_1 - p_2 < 0$, then $P\text{-value} = P(Z < z_0)$

If $H_a: p_1 - p_2 > 0$, then $P\text{-value} = P(Z > z_0)$

If $H_a: p_1 - p_2 \neq 0$, then $P\text{-value} = 2P(Z > |z_0|)$

These probabilities can be found using technology like StatKey, SPSS, a calculator with statistical software, or even looked up in a table of standard normal probabilities.

Step 4: State your conclusion and interpret in the context of the hypotheses

As with the randomization hypothesis test, you make your decision based on the P -value from step 3.

- If $P\text{-value} < \alpha$
 - Conclusion: reject H_0 ,

- Interpretation: There is sufficient evidence to suggest that . . . (the alternative hypothesis is true).
- If $P\text{-value} \geq \alpha$
 - Conclusion: *fail to reject* H_0
 - Interpretation: There is not significant evidence to suggest that . . . (the alternative hypothesis is true).

Always write the interpretation in the context of the original hypothesis.

Class Example 6.3.3 *Accuracy of Lie Detectors*

Participants in a study to evaluate the accuracy of lie detectors were divided into two groups with one group reading true material and the other group reading false material, while connected to a lie detector. The two way table indicates whether the participants were lying or telling the truth and also whether the lie detector indicated they were lying or not.

	Detector says lying	Detector says not	Total
Person lying	31	17	48
Person not lying	27	21	48
Total	58	38	96

- Define the two population proportions p_1 and p_2 .
- Are the conditions met for the two-sample test for proportions? Verify your answer.
- Find the three sample proportions for the proportion of times the lie detector says the person is lying (the proportion for the lying people, the proportion for the truthful people, and the pooled proportion)
- Test to see if there is a difference in the proportion of times the lie detector says the person is lying, depending on whether the person is lying or telling the truth. Use a 5% level of significance and show all details of the test.

Class Example 6.3.4 *Tagging Penguins*

A study was conducted to see if tagging penguins with metal tags harms them. In the study, 100 penguins were randomly assigned to receive a metal tag or (as a control group) an electronic tag. One of the variables studied is survival rate ten years after the penguins were tagged. The scientists observed that 20% of the 50 metal tagged penguins survived while 36% of the 50 electronically tagged penguins survived.

- Are the conditions met for using the normal distribution? Verify your answer.
- Define the two population proportions p_1 and p_2 .
- Test at $\alpha = 0.05$ to see if the survival rate is lower for all metal tagged penguins than for all electronically tagged penguins. Do metal tags appear to reduce survival rate in penguins?

6.4 Inference for Difference in Means

Key Concepts

- Find the mean and standard error for a distribution of differences in sample means between two groups
- Recognize when a t -distribution is an appropriate model for a distribution of the standardized difference in two sample means
- Use a t -distribution, when appropriate, to compute a confidence interval for the difference in means between two groups
- Use a t -distribution, when appropriate, to test a hypothesis about a difference in means between two groups

Distribution of a difference in means

For quantitative data, we often want to learn about a difference in averages from two groups. We can use the CLT to conduct inference about differences in averages if we have a large enough sample size. Consider two groups n_1 and n_2 that come from populations with means μ_1 and μ_2 and standard deviations σ_1 and σ_2 respectively. If sample size is large enough, the CLT says that the distribution of all differences in sample averages ($\bar{x}_1 - \bar{x}_2$) will follow a normal distribution that is centered at the true population difference in averages ($\mu_1 - \mu_2$). When we use the CLT, we can estimate the standard error for a sample average by

$$SE(\bar{x}_1 - \bar{x}_2) = \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$$

As with single averages, we almost never know the population standard deviations (σ_1 and σ_2), so we estimate them by using the sample standard deviations (s_1 and s_2).

When we use s_1 and s_2 instead of σ_1 and σ_2 , $\bar{x}_1 - \bar{x}_2$ no longer follows a normal distribution. The difference follows a t -distribution and the degrees of freedom are approximately equal to the smaller of $n_1 - 1$ and $n_2 - 1$.

We estimate the standard error of $\bar{x}_1 - \bar{x}_2$ with $\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}$

Conditions

In order to be able to use this distribution for a difference in averages, you need to make sure that each group satisfies the conditions for an average. Recall from the 6.4 – 6.6 notes:

Each sample must satisfy the necessary conditions for averages.

1. For a large enough sample size ($n_1 \geq 30$ and $n_2 \geq 30$), you can use the t -distribution
2. If $n < 30$, then the t -distribution is only a good approximation if the population is approximately normal.

Class Example 6.4.1 Describe the sampling distribution

For each of the following questions, identify whether or not a t -distribution would be appropriate.

- a) Comparing averages from two samples of sizes $n_1 = 10$ and $n_2 = 15$, from populations that are approximately normal.
- b) Comparing averages from two samples of sizes $n_1 = 15$ and $n_2 = 40$, where population 1 is heavily

skewed right, and population 2 is approximately normal.

- c) Comparing averages from two samples of sizes $n_1 = 50$ and $n_2 = 50$, from two populations that are heavily skewed to the left.

CI for a difference in means

Provided that for each group the sample size is large enough ($n \geq 30$) or the population is approximately normal, a confidence interval for a difference in population averages $\mu_1 - \mu_2$ can be constructed using

Remember, each sample must satisfy the necessary conditions for averages.

$$(\bar{x}_1 - \bar{x}_2) \pm t^* \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}$$

where $\bar{x}_1 - \bar{x}_2$ is the difference in sample averages and t^* is the appropriate confidence multiplier.

Class Example 6.4.2 Highball vs tumbler

Two common types of glassware at a bar are “highballs” (tall and thin) and “tumblers” (short and wide). Researchers randomly assigned participants either a “highball” glass or a “tumbler,” each of which held 355 ml. Participants were asked to pour a shot (1.5 oz = 44.3 ml) into their glass. The summary statistics are given in the table below.

	n	$\bar{x}$	s
Highball	99	42.2 ml	16.2 ml
Tumbler	99	60.9 ml	17.9 ml

Summary statistics for amount poured {,striped}

We want to know about the difference in the average amount of liquid poured into the two types of glasses.

- a) What are the variables of interest in this study? Are they categorical or quantitative? Which is the explanatory variable and which is the response?

- b) What is the parameter of interest?

- c) Find and interpret a 95% confidence interval estimate for the difference in the average amount poured in the two different types of glasses. Use highball for group 1 and tumbler for group 2.

- d) If you were the owner of a local bar, from a profit perspective, do you think there is any reason to prefer the highball to the tumbler glass?

Class Example 6.4.3 *Swimming heats at the Olympics*

The data in the table below are finishing times randomly selected from the preliminary heats for the 100m women's backstroke in the 2020 Tokyo Olympics for heats 5 and 6.

Heat 5	57.96	59.02	59.88	60.07
Heat 6	57.88	58.69	59.99	60.07

Randomly selected finishing times for 100m women's backstroke {,striped}

We want to know about the difference in the average finishing times for the two preliminary heats.

- a) What are the variables of interest in this study? Are they categorical or quantitative? Which is the explanatory variable and which is the response?
- b) What is the parameter of interest?
- c) Find and interpret a 95% confidence interval estimate for the difference in the average time between the two heats. You may assume that the finishing times for all participants in the heats are approximately normally distributed.
- d) Does your interval in c) suggest that there is a difference in the average finishing time between the heats?

Hypothesis test for a difference in means

Provided that for each group the sample size is large enough ($n \geq 30$) or the population is approximately normal, then according to the CLT

Each sample must satisfy the necessary conditions for averages.

$$t = \frac{\bar{x}_1 - \bar{x}_2}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}$$

follows a t -distribution.

We can use this fact to conduct a hypothesis test. As with the hypothesis test we have discussed thus far, there are a specific set of steps to conduct a hypothesis test using the CLT.

Step 1: Check conditions for the test

Check that for both groups $n \geq 30$ or the population is approximately normal.

Step 2: State the hypotheses

For a difference in averages, the null hypothesis always has the form

$$H_0 : \mu_1 - \mu_2 = 0$$

The alternative hypothesis has one of the following three forms:

$$H_a : \mu_1 - \mu_2 < 0$$

$$H_a : \mu_1 - \mu_2 > 0$$

$$H_a : \mu_1 - \mu_2 \neq 0$$

Step 3: Calculate the test statistic

To conduct the hypothesis test using the CLT, we use a standardized test statistic of

$$t = \frac{\bar{x}_1 - \bar{x}_2}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}$$

which follows a t -distribution.

Step 4: Find the p -value and give a sketch to illustrate the p -value

To find the p -value, you must use technology to find what proportion of the standard normal distribution is more extreme than the test statistic (in the direction of the alternative hypothesis). If your alternative is two-sided, then you multiply the one-sided p -value by 2. The sketch that you give should be the same as the sketch for a single mean; however, you do not need to give the degrees of freedom.

Step 5: State your decision and interpret in the context of the hypotheses

As with the randomization hypothesis test, you make your decision based on the p -value from step 3.

- If $p\text{-value} < \alpha$
 - Decision: *reject* H_0 ,
 - Interpretation: We have significant evidence to suggest that the alternative hypothesis is correct.
- If $p\text{-value} \geq \alpha$
 - Decision: *fail to reject* H_0

- Interpretation: We do not have significant evidence to suggest that the alternative hypothesis is correct.

You should *always* write your interpretation in the context of the original hypothesis.

Class Example 6.4.4 *Highball vs tumbler revisited*

Recall the following summary statistics from the highball vs tumbler example.

	n	$\bar{x}$	s
Highball	99	42.2 ml	16.2 ml
Tumbler	99	60.9 ml	17.9 ml

Summary statistics for amount poured {,striped}

Conduct a hypothesis test to see if there is a difference between the average amounts poured in the two types of glasses (use $\alpha = 0.05$).

- a) Define the relevant parameter(s) and state the null and alternative hypotheses. Be sure to check any relevant conditions.

-
- b) What is the test statistic.

-
- c) Find the p -value and give a sketch to illustrate your p -value.

-
- d) State your decision and interpret your findings in the context of the study.

-
- e) Do the findings from your hypothesis test agree with your confidence interval?
-

Class Example 6.4.5 *100 m backstroke times revisited*

The summary statistics from the 100m backstroke example are given in the table below.

	n	$\bar{x}$	s
Heat 5	4	59.23 s	0.96 s
Heat 6	4	59.16 s	1.06 s

Summary statistics for finishing times for 100 m women's backstroke. {striped}

Conduct a hypothesis test to see if there is a difference between the average finishing times for the two heats (use $\alpha = 0.05$).

- a) Define the relevant parameter(s) and state the null and alternative hypotheses. Be sure to check any relevant conditions.

- b) What is the test statistic.


- c) Find the p -value and give a sketch to illustrate your p -value.

- d) State your decision and interpret your findings in the context of the study.

- e) Do the findings from your hypothesis test agree with your confidence interval?

Class Example 6.4.6 *Teaching methods and post-test scores*

The following SPSS output is from an analysis of data from an experimental study conducted by Baumann and Jones, as reported by Moore and McCabe (1993). Students were randomly assigned to one of two experimental groups: “Basal” represents the traditional method of teaching, and “DRTA” represents an innovative teaching methods. The results are based on post-test scores for the two groups.




- What are the variables recorded in this study? Are they categorical or quantitative? Which of the variables is the independent variable? Dependent variable?
- Conduct a hypothesis test to determine whether there is a difference post-test scores between the two methods using the SPSS output above. Use $\alpha = 0.05$.

- c) Compare your results to the 95% confidence interval in the output. Do the test results agree with the interpretation of the confidence interval?

Class Example 6.4.7 *Insecticide effectiveness*

The counts of insects in agricultural experiments were taken between two types of insecticides (Beall, *Biometrika* 1942). The following SPSS output represents a test to determine if the two insecticides differed in effectiveness.



- a) What are the variables recorded in this study? Are they categorical or quantitative? Which of the variables is the independent variable? Dependent variable?

- b) Conduct a hypothesis test to determine whether there is a difference in insecticide effectiveness between the two insecticides using the SPSS output above. Use $\alpha = 0.05$.
- c) Compare your results to the 95% confidence interval in the output. Do the test results agree with the interpretation of the confidence interval?

6.5 Paired Difference in Means

Key Concepts

- Distinguish between separate samples and paired data when comparing two means
- Identify the advantages of using paired data when estimating a difference in means
- Use a t -distribution, when appropriate, to compute a confidence interval for a difference in means based on paired data
- Use a t -distribution, when appropriate, to test a hypothesis about a difference in means based on paired data

Inference for paired data

When examining differences in averages, typically, we use one of two methods. One option is to use the tools we discussed for the difference in averages. However, if there is a reason that the groups are not separate groups, then there is a better technique to use. If there is some kind of natural pairing between subjects in a study, then we can use techniques based on the concept of *paired differences*. Most often, paired differences occur when the subjects of the study receive two different levels of a factor (most commonly a *control* and a *treatment*). However, another way this appears is in a study in which two participants have been paired based on a number of different characteristics.

When analyzing paired data, we want to consider the *average of the differences* not the *difference of the averages*.

To decide between a two-sample t and paired difference t , consider how many times each case was measured. If each case is measured once, and we break the cases into two groups, we use two-sample t procedures. If each case is measured twice, we use paired difference t procedures.

Class Example 6.5.1 Deciding between difference in averages and paired difference

For each of the following, identify the cases for the study and decide if it would be more appropriate to use a difference in averages or paired difference.

- a) In a medical study to learn about the difference in the efficacy of generic versus brand name blood pressure medications, 50 men and 50 women were randomly assigned to receive either a generic medication or a brand name medication.
- b) Researchers want to learn about the average age differences between partners who have been together more than 10 years. They randomly select 50 sets of partners and ask them their age.

- c) In a study on the average difference in calories between servings of strawberry and vanilla yogurt, a food scientist randomly selected 12 brands of yogurt and recorded the calories in their strawberry and vanilla yogurts.
- d) The Office of Greek Life randomly selects 45 fraternity and sorority members and looks to see if there is a difference in the average GPAs for fraternities vs sororities.

Conditions

In order to be able to use the t -distribution for a difference in averages, you need to be sure that the differences satisfy the conditions for an average. Recall from the 6.4 – 6.6 notes:

1. For a large enough sample size ($n_d \geq 30$), you can use the t -distribution
2. If $n_d < 30$, then the t -distribution is only a good approximation if the population is approximately normal.

In this section, we will use a subscript d to indicate that we are working with differences.

We either need $n_d \geq 30$ or the population of differences to be approximately normal

CI for an average difference

Provided that the sample size is large enough so that $n_d \geq 30$ or the population is approximately normal, a confidence interval for the average difference in the population μ_d can be constructed using

$$\bar{x}_d \pm t^* \frac{s_d}{\sqrt{n_d}}$$

where $\bar{x}_d$ is average of the differences in the sample and t^* is the appropriate confidence multiplier from a t -distribution with $n_d - 1$ degrees of freedom.

Hypothesis test for an average difference

Provided that the sample size is large enough so that $n_d \geq 30$ or the population is approximately normal, then according to the CLT

$$t = \frac{\bar{x}_d}{s_d / \sqrt{n_d}}$$

follows a t -distribution with $n_d - 1$ degrees of freedom.

We can use this fact to conduct a hypothesis test. As with the hypothesis test we have discussed thus far, there are a specific set of steps to conduct a hypothesis test using the CLT.

Step 1: Check conditions for the test

Check whether $n_d \geq 30$ or the population of differences is approximately normal.

Step 2: State the hypotheses

For a difference in averages, the null hypothesis always has the form

$$H_0 : \mu_d = 0$$

The alternative hypothesis has one of the following three forms:

$$H_a : \mu_d < 0$$

$$H_a : \mu_d > 0$$

$$H_a : \mu_d \neq 0$$

Step 3: Calculate the test statistic

To conduct the hypothesis test using the CLT, we use a standardized test statistic of

$$t = \frac{\bar{x}_d}{s_d / \sqrt{n_d}}$$

which follows a t -distribution with $n_d - 1$ degrees of freedom.

Step 4: Find the p -value and give a sketch to illustrate the p -value

To find the p -value, you must use technology to find what proportion of the t -distribution is more extreme than the test statistic (in the direction of the alternative hypothesis). If your alternative is two-sided, then you multiply the one-sided p -value by 2. The sketch is the same as for a single mean (sections 6.4 – 6.6).

Step 5: State your decision and interpret in the context of the hypotheses


As with the randomization hypothesis test, you make your decision based upon the p -value from step 3.

- If $p\text{-value} < \alpha$
 - Decision: *reject* H_0 ,
 - Interpretation: We have significant evidence to suggest that the alternative hypothesis is correct.
- If $p\text{-value} \geq \alpha$
 - Decision: *fail to reject* H_0
 - Interpretation: We do not have significant evidence to suggest that the alternative hypothesis is correct.

You should *always* write your interpretation in the context of the original hypothesis.

Class Example 6.5.2 CAOS Comparisons

The CAOS (Comprehensive Assessment of Outcomes in Statistics) exam is an online multiple-choice test on concepts covered in a typical introductory statistics course. Students take one version before the start of the course and another version after the course ends. The data are summarized in the following QQ plot and SPSS output.



- a) Using the output, record the 95% confidence interval for the average difference in test scores? Interpret this interval.
-
- b) Conduct a test to determine if the test scores change, on average, between the pre- and post-test. Use $\alpha = 0.05$.
-

Class Example 6.5.3 *Body image and weight preoccupation*

A study was conducted to determine how males and females report their weight and height. 82 males and 99 females were asked their height and weight, then height and weight were measured for each participant. (Davis, C. (1990). "Body image and weight preoccupation: A comparison between exercising and non-exercising women." *Appetite*, 15, 13–21.)

The study reported that "women appear to accurately report their weight; we found no evidence of a difference between weight and reported weight ($\bar{x}_d = 1.62\text{lbs}, s_d = 10.14, t = 1.45, p\text{-value: } 0.1513$)."

The study also reported that "men, on the other hand, appear to underestimate their weight ($\bar{x}_d = -0.59\text{lbs}, s_d = 2.79, t = -2.13, p\text{-value: } 0.036$)."

Finally, the study went on to say that "there is also a significant difference between men and women when measuring the differences in height and reported height. Men overestimate their height ($\bar{x}_d = 1.76\text{in}, s_d = 2.17, t = 7.41, p\text{-value} < 0.001$) while women do not overestimate their height ($\bar{x}_d = 1.28\text{in}, s_d = 11.04, t = 1.15, p\text{-value: } 0.2517$)."

There are five statements in this study that indicate statistical findings. For each of the following, evaluate the validity of the claim based on the statistical evidence, and correct the statement if necessary. Only two of the conclusions are correct.

- a) Women appear to accurately report their weight; we found no evidence of a difference between weight and reported weight.

- b) Men, on the other hand, appear to underestimate their weight.


- c) Men overestimate their height.

- d) Women do not overestimate their height.

- e) There is also a significant difference between men and women when measuring the differences in height and reported height.

Class Example 6.5.4 *Pulse rates in class*

Do you think your pulse rate is higher when you are taking a quiz than when you are sitting in a lecture? The data in Table 6.27 show pulse rates collected from 10 students in a class lecture and then from the same students during a quiz. The data are summarized in the following output.



- a) Record the 95% confidence interval for the average difference in pulse rates? Interpret this interval.
-
- b) Conduct a test to determine if pulse rates are greater, on average, during a quiz than during a lecture. Use $\alpha = 0.05$.
-

7.1 Goodness of Fit for a Single Categorical Variable

Key Concepts

- Test a hypothesis about a categorical variable using a chi-square goodness-of-fit test
- Recognize when a chi-square distribution is appropriate for testing a categorical variable

Testing more than two groups

Up until now, we have discussed testing for categorical variables when there are only two groups. However, there are many statistical questions that require us to test categorical variables with three or more groups. In this chapter, we will learn about two new tests that give us additional methods to learn about categorical variables.

χ^2 test for goodness-of-fit

In this lecture, we will be discussing a new hypothesis test called the χ^2 (chi-square) goodness-of-fit test. The χ^2 goodness-of-fit test is used to test the relative frequency (or proportion) of observations for three or more groups.

The χ^2 goodness-of-fit test is used to test proportions when we have more than two groups in a single categorical variable.

Class Example 7.1.1 Relative frequencies

Javier recently took an exam in a chemistry class that had 100 multiple choice questions. There were four options per question (a, b, c, and d). Out of the 100 questions, he answered “a” 40 times, “b” 20 times, “c” 15 times, and “d” 25 times.

a) Find the relative frequencies for each of the answers on his exam.

b) If he was just guessing on each question, what proportion of the time should he answer each option.

Hypotheses

The χ^2 goodness-of-fit hypothesis test lets us see if Javier’s responses follow a different breakdown than what we expect to see if he were just guessing. The hypotheses that we want to test in this example are as follows:

$$H_0 : p_a = 0.25, p_b = 0.25, p_c = 0.25, p_d = 0.25$$

$$H_a : \text{At least one } p_i \text{ is different}$$

Note that the alternative only states that at least one is different. If we reject the null hypothesis here, we do not know which one is different, we only know that one of them is.

Expected counts

As with all hypothesis tests, we assume that the null hypothesis is true when we conduct our test. For the χ^2 goodness-of-fit test, we use this assumption to figure out the counts we *expect* to see in the sample. To find the expected count for group i :

$$\text{Expected count} = n \cdot p_i$$

Class Example 7.1.2 *Expected counts*

In the previous example, we identified the proportion of the time Javier should answer each option. Use these proportions to find the expected count of times he will answer each letter in a 100 question exam.

Test statistic

When we conduct the test, the goal is to see if the counts we *observed* are farther from the *expected* counts than we would reasonably expect to see due to random chance. We use the following test-statistic for the χ^2 test

$$\chi^2 = \sum \frac{(\text{Observed} - \text{Expected})^2}{\text{Expected}}$$

We want to see if the *observed* counts are far enough away from the *expected* counts

χ^2 distribution

There is a χ^2 distribution (like a normal or t -distribution) that we can use if we have a large enough sample size.

In order for us to be able to use the χ^2 distribution, all of the expected counts need to be at least 5 or larger. If we have met this condition, then we can use a χ^2 distribution with $a - 1$ degrees of freedom where a is the number of groups in the categorical variable.

To use the χ^2 distribution, each expected count must be at least 5 or larger

Our null distribution has $a - 1$ degrees of freedom if there are a categories.

Class Example 7.1.3 *Finish the hypothesis test*

Check the conditions, identify the degrees of freedom, give the test statistic, and use the χ^2 distribution to find the p -value for this hypothesis test.

Class Example 7.1.4 *Puppies!*

Every Labrador retriever has a set of genes (called a genotype) composed of B and E genes that control their color. The three primary coat colors are black, brown, and yellow. Without genetic testing it's difficult to know the genotype of a particular dog, however, if the genotypes of both parents are known, the percentage of puppies that fit into each color is known. Suppose a breeder believes their chocolate lab is type Eebb and their yellow lab is eeBb. If so, their puppies should be 25% black, 25% chocolate, and 50% yellow. The table below gives the counts of puppies of each color from the past two litters. Perform a test to determine if there is evidence the parental genotypes differ from what the breeder believes. Use $\alpha = 0.05$.

Puppies	Black	Chocolate	Yellow	Total
Observed (counts)	6	6	8	20
Expected (counts)				

Sample of puppy colors for two litters

Class Example 7.1.5 *Age discrimination on U.S. juries*

It is often said that older people are overrepresented on juries in the United States. The *UCLA Law Review* randomly selected cases in one county in California, and recorded the number of jurors in various age groups given in the table below. Do the following data suggest that the *UCLA Law Review* should expand the scope of their study to more counties (they would do this if they observed discrimination)? Use $\alpha = 0.05$.

	21-40	41-50	51-60	61+	Total
Countywide Proportion (%)	42.0	23.0	16.0	19.0	100.0
Grand Jurors (counts)	10	18	38	66	132
Expected (counts)					

Actual ages of jurors in California county

Class Example 7.1.6 *U.S. ethnicities*

In a sample of 74 flights in 2023, the La Crosse Regional airport experienced 7 schedule modifications. According to the Bureau of Transportation Statistics, 21.53% of flights in the U.S. experienced a schedule modification.

- Is the proportion of scheduled modifications from the La Crosse Regional Airport different than the national rate? Use a χ^2 goodness-of-fit test at $\alpha = 0.05$.
- What other kind of hypothesis test might we conduct to answer this question? How do the results from that hypothesis test compare with the χ^2 test?

7.2 Independence for Two Categorical Variables

Key Concepts

- Test a hypothesis about an association between two categorical variable using a chi-square association test
- Recognize when a chi-square distribution is appropriate for testing an association between two categorical variables

Association Between Categorical Variables

In this lecture, we will be discussing a new hypothesis test called the χ^2 (chi-square) association test. The χ^2 association test is used to test about the relative frequencies (or proportions) of observations for two categorical variables.

The χ^2 association test is used to test the relationship between two categorical variables

Hypotheses

The hypotheses for the χ^2 association test tends to be described more generally than the other tests that we have seen so far. In fact, rather than using parameters in the hypotheses, we will typically express our hypotheses in words. The hypotheses that

we want to test in this test are as follows:

H_0 : *Variable 1* is not associated with *Variable 2*

H_a : *Variable 1* is associated with *Variable 2*

If we reject the null hypothesis here, we can only say that there is an association, but we cannot state exactly what that relationship is.

Class Example 7.2.1 *Traffic cameras and injuries*

Many cities have introduced traffic cameras at high-risk street intersections in an effort to reduce the severity of traffic accidents. Citizen rights groups have complained about the efficacy of these cameras. In fact, they doubt they have any impact whatsoever. A sample of 2938 reported accidents was collected including the type of accident (Death/Disabling Injury, Evident Injury, or Probable Injury) and camera status at the intersection (Before Camera, After Camera). What are the hypotheses to test the relationship between accident severity and presence of a traffic camera?

Expected counts

As with all hypothesis tests, we assume that the null hypothesis is true when we conduct our test.

For the χ^2 association test, if the null hypothesis is true, the two categorical variables are *independent*. We use this assumption to figure out the counts we *expect* to see in the sample. To find the expected count for (Variable 1 group i , Variable 2 group j):

The expected counts come from the probability rules. If A and B are independent, then $P(A \cap B) = P(A) * P(B)$

$$\text{Expected count} = \frac{\text{Row total} \times \text{Column total}}{\text{Total total}}$$

Class Example 7.2.2 *Traffic cameras and injuries, continued*

The table below summarizes the traffic camera data. Use this data to compute a table of expected counts.

	Before Camera	After Camera
Death/Disabling	61	27
Evident	210	136
Probable	1659	845

Traffic camera status and accident severity

Test statistic

When we conduct the test, the goal is to see if the counts we *observed* are farther from the *expected* counts than we would reasonably expect to see due to random chance. We use the same test-statistic for the χ^2 association test that we used for the χ^2 goodness-of-fit test:

We want to see if the *observed* counts are far enough away from the *expected* counts

$$\chi^2 = \sum \frac{(\text{Observed} - \text{Expected})^2}{\text{Expected}}$$

Class Example 7.2.3 *Traffic cameras and injuries, continued*

Using the traffic camera data, find the χ^2 test statistic using the traffic camera data to test if there is an association between traffic cameras and injury severity.

χ^2 test for association

In order for us to be able to use the χ^2 distribution, all of the expected counts need to be at least 5 or larger. If we have met this condition, then we can use a χ^2 distribution with $(r - 1) \times (c - 1)$ degrees of freedom where r is the number of groups in the first categorical variable (number of rows) and c is the number of groups in the second categorical variable (number of columns). To find the p -value using the χ^2 distribution, you can either use StatKey or a TI-83/84. The p -value calculation here works the same as for the goodness-of-fit test.

To use the χ^2 distribution, each expected count must be at least 5 or larger

Class Example 7.2.4 *Traffic cameras and injuries, continued*

Check the conditions, identify the degrees of freedom, and use the χ^2 distribution to find the p -value for the hypothesis test for the traffic camera study.

Class Example 7.2.5 *Star Trek*

In Star Trek, the colored uniforms worn by the crew identify their work area. Among Trekkies (fans of the show), they have begun to notice a pattern - they believe there is an association with uniform color and if the character suffers a fatality on the show. The table below gives a breakdown of the various crew colors and the number of fatalities - does this give evidence to suggest they are related?

	Blue	Gold	Red
Fatality	7	9	24
Survival	129	64	263

Uniform color and fatality count

Class Example 7.2.6 *Racial steering*

A subtle form of racial discrimination in housing is “racial steering.” Racial steering occurs when real estate agents show prospective buyers only homes in neighborhoods already dominated by that family’s race. This violates the Fair Housing Act of 1968. According to an article in *Chance* magazine (Vol. 14, no. 2 [2001]), tenants at a large apartment complex recently filed a lawsuit alleging racial steering. The complex is divided into two parts: Section A and Section B. The plaintiffs claimed that white potential renters were steered to Section A, while African-American renters were steered to Section B. The table below displays the data that were presented in court to show the locations of recently rented apartments. Do you think there is evidence of racial steering?

	White	Black
Section A	87	8
Section B	83	34

Apartment sections broken down by race


8.1 Analysis of Variance

Key Concepts

- Use ANOVA to test for a difference in means among several groups
- Explain how variation between groups and variation within groups are relevant for testing a difference in means between multiple groups.

Motivation

In a two-sample t -test (test for difference in means), we record values of one categorical variable (with two groups) and one quantitative variable. Our goal is to compare the means between the two groups.



Two sets of sample data are displayed in the figure above. In which set of sample data, A or B, is there likely to be stronger evidence of a difference between the two sample means? Why?

Analysis of Variance

In the example above, the sample mean for Group 1 was the same for both datasets, and likewise for Groups 2. However, the *spread* within each sample was different for the two datasets. We used the spread to decide which provided stronger evidence of a difference. Recall, the test statistic for a two-sample t -test is:

$$t = \frac{\bar{x}_1 - \bar{x}_2}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}$$

In the numerator of the test statistic, we calculated the *difference* between the two sample means, or the spread between the two groups. In the denominator, we calculated the standard error, which is a measure of spread within the two groups. How would this test statistic change if we wanted to test if there was a difference in

means between three or more groups?

The Analysis of Variance (ANOVA) procedure can be regarded as an extension of the two-sample t -test to more than two samples. In fact, the test statistic is formed in much the same way. The test statistic for ANOVA is an F -statistic, and has the following form.

ANOVA is an extension of the two-sample t -test to more than two groups.

$$F = \frac{\text{Spread between group means}}{\text{Spread within groups}}$$

This is the same concept as our t -statistic from a two sample t -test; in fact, we can think of the difference in means (numerator of t) to be a measure of spread between the means.

Statement of hypotheses

The ANOVA hypothesis test is used to see if groups have significantly different averages. If we are considering k groups, then the hypotheses for the ANOVA test are

$$\begin{aligned} H_0 : \mu_1 &= \mu_2 = \dots = \mu_k \\ H_a : &\text{At least one } \mu_i \text{ is different} \end{aligned}$$

Class Example 8.1.1 *Quality control for lenses*

A company that produces glass lenses would like to know if the three presses they use are producing the same average thickness in lens. They hire an engineer to test to see if the machines should be calibrated. Write the hypothesis to test this using an ANOVA hypothesis test.

Finding the test statistic

The test statistic for an ANOVA hypothesis test is an F -statistic. To find an F -statistic, we will use an ANOVA table. An ANOVA table is used to help us compare the variability *between* the sample means and the variability *within* the samples.

SS stands for *sum of squares*

MS stands for *mean squares*

Source	df	SS	MS	F	p-value
Group	$k - 1$	SS_{Group}	$MS_{\text{Group}} = \frac{SS_{\text{G}}}{k-1}$	$F = \frac{MS_{\text{Group}}}{MS_{\text{Error}}}$	$\text{Fcdf}(\text{F-stat}, 9999, \text{df}_{\text{Group}}, \text{df}_{\text{Error}})$
Error	$n - k$	SS_{Error}	$MS_{\text{Error}} = \frac{SSE}{n-k}$		
Total	$n - 1$	SS_{Total}			

Sample ANOVA table

Here are a few relationships to note as you are working with these tables:

$$SS_{\text{Total}} = SS_{\text{Group}} + SS_{\text{Error}}$$

$$df_{\text{Total}} = df_{\text{Group}} + df_{\text{Error}}$$

$$MS_{\text{Group}} = \frac{SS_{\text{Group}}}{df}$$

$$MS_{\text{Error}} = \frac{SS_{\text{Error}}}{df}$$

$$F = \frac{MS_{\text{Group}}}{MS_{\text{Error}}}$$

Finding the p-value

To find the p -value, you need to know the degrees of freedom for the numerator and denominator. You can find the p -value using technology like Statkey, SPSS, or a calculator.

Class Example 8.1.2 *Quality control continued*

Fill in the remaining spots in the ANOVA table below.

Source	df	SS	MS	F	<i>p</i> -value
Group		10.30			
Error					
Total	14	34.40			

Quality control ANOVA table

Find the *p*-value for the *F*-statistic that tests to see whether or not average glass thickness differs for each press. What are the degrees of freedom for the numerator and denominator? State your decision and provide an interpretation in the context of the problem.

Conditions to use the *F*-test

For the *F*-test, we need to check that each average is approximately normal (i.e. $n_i \geq 30$ or sampling from a normal population). We also need to know that the standard deviations in each group are similar enough. To check if the standard deviations are similar enough, it is enough to make sure that no group has a standard that is larger than twice the standard deviation for another group.

Conditions to use the *F*-test:

- We need to know that each average is approximately normal
- No group standard deviation can be more than 2 times the others

Class Example 8.1.3 *Quality control continued*

Have the conditions to use an *F*-statistic to test if any of the presses has a different average thickness been satisfied? Explain.

Class Example 8.1.4 *Rotten Tomatoes*

Rotten Tomatoes is a website providing movie ratings and reviews. Each movie gets a “Tomatometer” score, which is “The percentage of Approved Tomatometer critics who have given this movie a positive review.” A sample of movies for seven categories was taken, and the summary statistics are provided in the table below. Does the average Tomatometer score differ by movie genre?

Genre	$\bar{x}$	s	n
Action	44.97	25.10	31
Animation	60.58	26.73	12
Comedy	49.78	26.35	27
Drama	67.90	18.06	21
Horror	41.29	27.04	17
Romance	47.20	31.57	10
Thriller	62.38	28.66	13

Summary statistics for random sample of tomatometer scores for seven categories

- a) What are the variables recorded in this study? Are they categorical or quantitative? Which of the variables is the independent variable? Dependent variable?

- b) State the appropriate hypotheses for this test.

- c) Are the conditions for using the F -distribution satisfied? Why or why not?

- d) Regardless of your previous answer, test to see whether the Tomatometer score differ significantly by movie genre (use $\alpha = 0.05$)? Make a conclusion in context.

Source	df	SS	MS	F	p -value
Group		11407.4			
Error					
Total		93268.1			

Tomatometer ANOVA table

Class Example 8.1.5 *Caffeine and finger tapping*

Does caffeine consumption have an effect on twitch movements? To investigate this question, the following experiment was conducted. 30 subjects were randomly assigned to one of three groups, with 10 subjects in each group. The subjects were then given a tablet containing 0mg, 100mg, or 200mg of caffeine. Two hours later, they were asked to tap their finger as quickly as possible for one minute. Summary statistics are provided

in the table below.

	0 mg	100 mg	200 mg
$\bar{x}$	244.8	246.4	248.3
s	2.394	2.066	2.214

Summary statistics for caffeine and finger tapping experiment

- a) What are the variables recorded in this study? Are they categorical or quantitative? Which of the variables is the independent variable? Dependent variable?

- b) Based on the summary statistics above, do you think that caffeine has an effect on finger tapping?

- c) State the appropriate hypotheses for this test.

- d) Are the conditions for using the F -distribution satisfied? Why or why not?

- e) Regardless of your previous answer, test to see whether caffeine consumption cause a difference in the average number of finger taps (use $\alpha = 0.05$)? Make a conclusion in context.

Source	df	SS	MS	F	p -value
Group		61.4			
Error		134.1			
Total					

Caffeine and finger tapping ANOVA table

- f) Is there a *practical* difference between the groups? Is this the same as a *statistical* difference?

8.2 Analysis of Variance - Multiple Comparisons after ANOVA

Key Concepts

- Create confidence intervals for single means and differences of means after doing an ANOVA for means
- Test specific pairs of means after ANOVA indicates a difference in means, recognizing the problem of multiplicity

Pairwise comparisons and inference after ANOVA

Although an ANOVA hypothesis test will tell you that there is a difference in at least one of the population means, it does not specify which averages are different. If H_0 is not rejected in an ANOVA, then there is no need to look for differences among population means. However, if H_0 is rejected, the next step is to see which means are actually different from the others. For a significant ANOVA test, there are a number of methods to compare the averages for the different groups.

When we compare these groups, we do what is called *pairwise comparisons* of the means. The phrase *pairwise comparisons* means conducting hypothesis tests between all the possible pairs of means to see which pairs are different. We can find the number of comparisons among k population means by using the formula

$$\# \text{ of comparisons} = \binom{k}{2} = \frac{k!}{2!(k-2)!}$$

where k is the number of groups. Remember from the notes on 11.1, that this is read as “ k choose 2” and you can use the nCr function on your calculator to do the calculation.

If H_0 is not rejected in an ANOVA, then there is no need to look for differences among population means

Pairwise comparisons is just a fancy way of saying consider all the possible pairs of means and conduct hypothesis tests to see if they are different.

Pairwise comparisons are also called *multiple comparisons* or *post-hoc tests*

Class Example 8.2.1 *How many pairwise comparisons*

Recall example 8.1.5 in which we discussed comparing the average tomatometer scores for 7 different genres of movies. How many pairwise comparisons would you need to do in order to compare all the genres with one another?

Problem of multiplicity

If you use a two-sample t -test to compare two averages at $\alpha = 0.05$, then there is a 5% chance that we will make a Type I error. However, as we conduct two-sample t -tests to compare more pairs of averages, then we don't want to consider the Type I error rate for only a single test. We want to know the *overall Type I error rate*, or the chance that we make a Type I error in at least one of the tests we conduct. To find the overall Type I error rate when comparing all pairwise comparisons for k groups, we use the following formula:

The *overall Type I error rate* is the chance that we made a Type I error in at least one of our tests.

$$\text{Overall Type I error rate} = 1 - (1 - \alpha)^{\binom{k}{2}}$$

Class Example 8.2.2 *What is the overall Type I error rate*

If we conduct pairwise comparisons of the average tomatometer scores for the 7 genres of movies at $\alpha = 0.05$, what is the overall Type I error rate?

Inference for population means after ANOVA

Population means can be compared using simultaneous hypothesis tests among pairs or simultaneous confidence intervals among pairs. Confidence intervals for individual population means can also be constructed using information from the ANOVA table.

Bonferroni correction for multiplicity

There are a number of ways to correct for the overall Type I error rate, but one of the easiest is the Bonferroni correction. When we do this correction, we make an adjustment to the significance levels for the hypothesis tests.

To use the Bonferroni correction with hypothesis tests, we take the desired overall Type I error rate, and divide by the number of pairwise comparisons.

$$\text{Bonferroni corrected } \alpha = \frac{\text{Desired overall Type I error rate}}{\text{Number of pairwise comparisons}}$$

Class Example 8.2.3 *Bonferroni correction for tomatometer scores*

If we want the overall Type I error rate to be 0.10, what is the Bonferroni corrected α we should use when comparing the average tomatometer scores for the 7 genres of movies?

Bonferroni simultaneous t-intervals

After H_0 has been rejected in an ANOVA, confidence intervals for comparing μ_i with μ_j are constructed using

$$(\bar{x}_i - \bar{x}_j) \pm t^* \sqrt{MSE \left(\frac{1}{n_i} + \frac{1}{n_j} \right)}$$

where MSE is the mean square error from the ANOVA table. The degrees of freedom for t^* in this interval is the degrees of freedom for error in the ANOVA table ($n - k$).

After confidence intervals have been constructed for all pairs of differences, we can find population means that are different by looking to see which intervals for the differences in means do *not* contain zero. You are not required to do this computation by hand, but you should be able to do this using SPSS.

$\sqrt{MSE}$ is also called the *pooled standard deviation*

This confidence interval uses $n - k$ degrees of freedom

Class Example 8.2.4 *Salary and Sales*

A company that employs a large number of salespeople is interested in which group of salespeople sells the most: those strictly on commission, those with a fixed salary, or those with a reduced salary plus a commission. A random sample of the previous month's records for salespeople are inspected, and the amount of sales (in dollars) is recorded for each. The ANOVA table and output for the Bonferroni simultaneous CIs at a 95% overall level of confidence is also below.

Pay Type	$\bar{x}$	s	n
Commission	473.50	29.87	6
Fixed Salary	423.20	14.43	5
Commission Plus Salary	476.25	31.64	4

Summary statistics, ANOVA output, and Bonferroni simultaneous 95% confidence intervals for salary and sales data

-
- a) Are the conditions for this test satisfied?
-
- b) Despite your answer to a), conduct a test (using $\alpha = 0.05$) to see if there any differences among different average sales for the three salary type.
-
- c) Use the 95% Bonferroni CIs to determine which pairs of treatment groups have different population means. Use a complete sentence to state how they differ (*i.e.* which is larger or smaller, which are the same) and, if they differ, state the interval for the estimated difference.
-

Bonferroni simultaneous t-tests

After we have conducted an ANOVA and found a significant result that at least one of the averages is different, we can conduct a two-sample t -test using pooled standard deviation. The test statistic is very similar to the one we discussed in chapter 6, but the standard error is different. To compare group i with group j , you should use

$$t = \frac{\bar{x}_i - \bar{x}_j}{\sqrt{MSE \left(\frac{1}{n_i} + \frac{1}{n_j} \right)}}$$

For the degrees of freedom for this test, you should use the degrees of freedom for error from your ANOVA ($n - k$), and you will need to use the Bonferroni corrected α when making your decision.

Class Example 8.2.5 *Restaurant tips*

In this example, we will be conducting many different tests from a dataset that contains information on the tipping patterns of patrons of the *First Crush* bistro in northern New York state. The data from 157 bills include the amount of the bill, size of the tip, percentage tip, number of customers in the group, whether or not a credit card was used, day of the week, and a coded identity of the server (A, B, or C). The first four variables are quantitative and the last three are categorical. For each test, you can use a hypothesis test or confidence interval (if appropriate). On each test, be sure to verify the necessary conditions.

- a) Most restaurant patrons leave a tip between 15% and 20% of the bill, and some restaurant customers determine the percent tip to leave based on the quality of the service. Summary statistics for mean tip percentage left for each of the three servers and a partial ANOVA table are given in the tables below.

Server	Sample size	Mean	Std. Dev.
A	60	17.543	5.504
B	65	16.017	3.485
C	32	16.109	3.376

Mean tip percentage by server

Source	df	SS	MS	F	p-value
Server			41.57		
Error					
Total		3001			

ANOVA table for mean tip percentage

- i. Conduct the appropriate hypothesis test to see if there a difference in mean tip percentage between the three servers (use $\alpha = 0.05$).
-
- ii. Is it appropriate to examine pairwise differences to see which of the servers differs in average tip percentage? Explain.
-
- iii. Regardless of your answer in a.ii, if we want the overall Type I error rate to be 0.05, what is the Bonferroni corrected α we should use when comparing the average tip percentage for the servers?
-

- b) Are some servers given the big spenders (or large groups) while other tend to those having only a cup of coffee or a glass of wine? Summary statistics for average bill for each of the three servers and a partial ANOVA table are given in the tables below.

Server	Sample size	Mean	Std. Dev.
A	60	22.76	12.71
B	65	21.14	10.19
C	32	25.92	14.35

Average bill by server

Source	df	SS	MS	F	p-value
Server				1.671	
Error			146.5		
Total					

ANOVA table for average bill

- i. Conduct the appropriate hypothesis test to see if there a difference in average bill between the three servers (use $\alpha = 0.05$).
-
- ii. Is it appropriate to examine pairwise differences to see which of the servers differs in average bill amount? Explain.
-
- iii. Regardless of your answer in a.ii, if we want the overall Type I error rate to be 0.10, what is the Bonferroni corrected α we should use when comparing the average bill for the servers?
-

- c) The table below summarizes the number of bills by server. Conduct the appropriate hypothesis test to see if the servers serve equal numbers of tables.

Server	A	B	C	Total
Number of bills	60	65	32	157

Number of bills by server

- d) The table below summarizes how patrons paid (cash or card) for each server. Conduct the appropriate hypothesis test to see if there is a relationship between server and how patrons pay.

Server	A	B	C	Total
Cash	39	50	17	106
Card	21	15	15	51
Total	60	65	32	157

Cash or card by server

9.1 Inference for Slope and Correlation

Key Concepts

- Use computer output to make predictions and interpret coefficients using a fitted simple linear model
- Construct a confidence interval for the slope in a regression model
- Test a hypothesis about the slope of a regression model
- Test for evidence of a linear association between two quantitative variables using a sample correlation
- Compute and interpret the value of r^2 for a regression model
- Check a scatterplot for obvious departures from a simple linear model

Modeling with two quantitative variables

In the last chapter we discussed how to use a categorical variable to model a quantitative response using ANOVA, and in this chapter we will be discussing how to use a quantitative variable to model a quantitative response. In statistics, when both the explanatory and response variables are quantitative, we use *regression* in order to explain the response variable. However, before we get into the details for regression, it's helpful to first discuss the concept of correlation.

Exploring correlation with a scatterplot

To get a sense of how one quantitative variable impacts another, it is helpful to first plot the data. To graphically display two quantitative variables, we use a *scatterplot*, or a graph that displays the relationship between two quantitative variables. When looking at a scatterplot, we should consider four different aspects of the plot:


1. Do the points form a clear trend with a particular direction, are they more scattered about a general trend, or is there no obvious pattern?
2. If there is a trend, is it generally upward or generally downward as we look from left to right? A general upward trend is called a *positive* association while a general trend downward is called a *negative* association.
3. If there is a trend, does it seem to follow a straight line, which we call a *linear association*, or some other curve or pattern?
4. Are there any outlier points that are clearly distinct from a general pattern in the data?

A *scatterplot* is a graph of the relationship between two quantitative variables


Properties of the linear correlation coefficient

There are a few properties about the correlation coefficient, r , you should know, and we will illustrate them with some pictures.

a) r is always between -1 and 1 , inclusive: $-1 \leq r \leq 1$



- b) The sign of r (positive or negative) indicates the direction of association.
- c) Values of r close to -1 or $+1$ show a strong linear relationship, while values of r close to 0 show no linear relationship.




- d) r has no units and is independent of the scale of either variable.
- e) Correlation is symmetric: The correlation between variables x and y is the same as the correlation between y and x .


Correlation coefficient cautions

Here are a few things to keep in mind when working with the correlation coefficient:

a) r can be heavily influenced by outliers.



- b) A strong positive or negative correlation coefficient does not (necessarily) imply a cause and effect relationship between the two variables.
- c) A correlation coefficient near zero does not (necessarily) mean that the two variables are not associated, since correlation measures only the strength of a *linear* relationship (see side figure).



Linear regression

Now we are going to introduce a new statistical method that can be used to fit a line to the dataset. To begin with, it's important to first define a *residual*. To find a residual for a single observation, we use the following formula:

$$\text{Residual} = \text{Observed} - \text{Predicted} = y - \hat{y}$$

We say that the line that fits the data best is the *least squares line*, which minimizes the sum of the squared residuals.

Equation of the line

A sample regression has the following form:

$$\hat{y} = b_0 + b_1x$$

where $\hat{y}$ is the predicted value for a specific value of the explanatory variable, x , b_0 is the intercept, and b_1 is the slope. The *slope*, b_1 , represents the average predicted change in the response variable, y , given a one unit increase in the explanatory variable, x . The *intercept*, b_0 , represents the predicted value of the response variable, y , if the explanatory variable, x , is zero. The equation for the intercept and slope are given as

$$b_1 = r \frac{s_y}{s_x} \quad b_0 = \bar{y} - b_1 \bar{x}.$$

One thing to note from this formula is that the *slope* and *correlation* should always have the same sign.

A *residual* is the difference between an observed value and a predicted value of a response.

The *least squares line* is the equation that has the smallest sum of squared residuals.

Notation:

$\hat{y}$: predicted value $-b_0$: intercept estimate $-b_1$: slope estimate


The *slope* and *correlation coefficient* should always have the same sign.

Class Example 9.1.1 *Boats*

A 2024 dataset from BoatTrader.com reveals the relationship between the listing price of a boat and its corresponding length in feet. Some summary statistics and a scatterplot of the data are given below.

Variable	Average	Std. Dev.
Price	51,498.28	66,128.95
Length	22.61	8.07

$$r = 0.364$$



- Looking at the plot, would you categorize the correlation as positive or negative? Strong, moderate, or weak?
- Which variable should be the explanatory variable, which variable should be the response?
- Use the summaries to find estimates for the slope and intercept for the regression line. Record your answer in the form $\hat{y} = b_0 + b_1x$.
- If one of the boats had a length of 20 feet and a price of \$50,000, find the predicted value and residual for this person.

e) Provide an interpretation of the slope in the context of the problem.

Inference for slopes

As with means and proportions, the regression coefficients, b_0 and b_1 , are statistics and are estimates of the population intercept, β_0 , and population slope, β_1 . Most of the time, we will not be interested in inference for β_0 , so the rest of the lecture on regression will focus on inference for the population slope β_1 . To conduct a hypothesis test for the slope, we will be testing

$$H_0 : \beta_1 = 0 \qquad H_a : \begin{cases} \beta_1 < 0 \\ \beta_1 > 0 \\ \beta_1 \neq 0 \end{cases}$$

As with all statistics, we need to report a standard error in order to determine if our findings are significant. This will be provided to you with software output, so you do not need to worry about calculating this on your own. However, you will need to know how to find the test statistic in order to find a p -value. For a hypothesis test about a slope, we use the test statistic

$$t = \frac{b_1 - 0}{SE(b_1)}$$

where $SE(b_1)$ stands for standard error of b_1 . To find a p -value, you should use a t -distribution with $n - 2$ degrees of freedom. The remaining steps of the hypothesis test are the same as all other hypothesis tests. The table below shows output that is typically given by software when conducting linear regression.

Coefficients	Estimate	Std. Error	t
(Intercept)	b_0	$SE(b_0)$	$t_{for b_0}$
Explanatory Variable	b_1	$SE(b_1)$	$t_{for b_1}$

Example output from software

Population parameters:

$-\beta_0$: population intercept $-\beta_1$:
population slope

$SE(b_1)$ is the standard error of b_1

Class Example 9.1.2 *Boats continued*

The table below gives the output from results for a simple linear regression of boat price with length for a 2025 dataset.

Coefficients	Estimate	Std. Error	t	p -value (2-sided)
(Intercept)	-78,776.98	4,431.58	-17.78	< 0.0001
Length	5,761.80	184.61	31.21	< 0.0001

Output from results for a simple linear regression of price with length

- a) Use the output to conduct a hypothesis test to see if there is a positive relationship between a boat's length and price. Use $\alpha = 0.05$.
-
- b) The hypothesis test revealed a statistically significant positive relationship between a boat's length and price. Is this practically significant?
-

Conditions for inference

In order to conduct inference for regression, we need to satisfy a few conditions.

1. The relationship should be linear
2. The residuals should have a normal distribution for all values of x
3. The variability in the response should be the same for all values of x
4. There should not be any outliers that unduly influence the results of the regression.

We can check conditions 1, 3, and 4 using a scatterplot of the data. Condition 2 can be checked using a histogram of residuals or a QQ plot.

Words of caution for regression

Although regression is a powerful tool, there are a couple of words of wisdom to keep in mind.

1. Like correlation, regression only tells us about the *linear* relationship between the explanatory and response variables.
2. Do not ever *extrapolate* or predict a response beyond the range of the explanatory variable. The regression equation only tells us about the relationship between the response and explanatory within the range of the explanatory variable.
3. As with correlation, just because there is a statistically significant relationship, it does not mean that there is a causal relationship. Remember *correlation does not mean causation*.

Coefficient of determination: r^2

There is another statistic from regression that is often used to compare two simple linear regressions models with each other. The *coefficient of determination*, notated r^2 , gives the percentage of variation in the *response* variable that we can explain by the linear regression using the *explanatory* variable. If we are picking between two different simple linear regression models, you should select the model that has a higher r^2 value. To find r^2 , square the correlation coefficient r .

r^2 gives the percentage of variation in the *response* variable that we can explain with the *explanatory* variable

Class Example 9.1.3 *Price and Horsepower*

In the previous example, we used the variable *length* to predict *price*. Now we're going to use *horsepower* to predict *price*. The table below gives the output from a simple linear regression of price with horsepower.

Coefficients	Estimate	Std. Error	<i>t</i>	<i>p</i> -value (2-sided)
(Intercept)	36,449.63	1,998.92	18.23	< 0.0001
Horsepower	162.27	8.50	19.09	< 0.0001

$r = 0.5245$ Output from results for a simple linear regression of drinks with classes missed

a) Using the output above, write the regression equation in the form $\hat{y} = b_0 + b_1x$.

b) Find the predicted and residual value for a boat with 500 horsepower and a price of \$90,000.

c) Interpret the slope of the line in the context of the problem.

d) Conduct a hypothesis test to see if there is a significant positive relationship between horsepower and price. Use $\alpha = 0.05$.

e) Does horsepower or length explain price better? Explain.

9.2 Confidence and Prediction Intervals

Key Concepts

- Compute and interpret the value of r^2 for a simple linear model using values in the ANOVA table.
- Compute and interpret a confidence interval for the mean response for a specific value of the predictor in a regression model
- Compute and interpret a prediction interval for the individual responses for a specific value of the predictor in a regression model

Understanding the definition of r^2 through ANOVA

When conducting a simple linear regression using the computer, often times, an ANOVA table is included in the output. The Sum of Squares column of the ANOVA table gives us a way to partition the variability of our response into two components, variability we can explain with our regression model, and variability that is unexplained. In the context of regression, this looks like

Variability in the response variable can be partitioned into two components:
 $-SS_{\text{Regression}} - SS_{\text{Total}}$

$$SS_{\text{Total}} = SS_{\text{Regression}} + SS_{\text{Error}}$$

In the previous section the coefficient of determination, r^2 , could be found by squaring the correlation coefficient, r . It can also be computed as

$$r^2 = \frac{SS_{\text{Regression}}}{SS_{\text{Total}}}.$$


Another formula for r^2 is

$$r^2 = \frac{SS_{\text{Regression}}}{SS_{\text{Total}}}$$

This is the reason r^2 is interpreted as the percentage of variation in the *response* variable that we can explain by the linear regression using the *explanatory* variable.

Class Example 9.2.1 Asteroids!

NASA's Near Earth Object Web Service (NeoWs) tracks near earth asteroids. In a sample of 1000 near earth asteroids taken in 2024 the orbital period (number of days to orbit the earth) and perihelion distance (distance from earth when the earth is closest to the sun, measured in Astronomical Units) was recorded. A scatterplot of the data and QQ plot of the residuals are included below. Computer output is provided below.



Source	df	SS	MS	F	p-value
Per. Dist.	1	2.3×10^7			< 0.0001
Error	998	1.1×10^8			
Total	999				

Partial ANOVA results for regression of perihelion distance on orbital period

Coefficients	Estimate	Std. Error	t	p-value
(Intercept)	117.00	37.42	3.13	< 0.0001
Per. Dist.	637.90	44.29	14.40	< 0.0001

Parameter estimates for regression of perihelion distance on orbital period

a) Complete the partial ANOVA table.

b) Calculate r^2 and provide an interpretation in the context of the study.

c) Write 2 to 3 sentences describing the findings of this study. If it is inappropriate to conduct inference, simply state that and justify your decision.

d) Suppose a new asteroid was discovered with a perihelion distance of 3 AU. Assuming it was appropriate to conduct inference, would it be reasonable to use these results to predict the orbital period of this asteroid? Explain.

Confidence and prediction intervals

One of the main reasons why we conduct linear regression is to predict a response for a given value of the explanatory variable. You can easily predict the response for any value of the explanatory variable using the regression equation provided you are not extrapolating beyond the range of the data. However, when making this prediction, it is important to account for the uncertainty in the estimates for slope and intercept. We can account for this uncertainty by using an interval. There are two main types of intervals that we will use for predictions:

1. Confidence interval for the *average* response
2. Prediction interval for an *individual* response

Intervals for a predicted responses

Recall that we use the symbol $\hat{y}$ to represent the predicted value of y , and this is found using the least squares regression equation

$$\hat{y} = b_0 + b_1 x$$

The *confidence interval for the average response* when $x = x^*$, is

$$\hat{y} \pm t_{n-2}^* \sqrt{MSE \left(\frac{1}{n} + \frac{(x^* - \bar{x})^2}{(n-1)s_x^2} \right)}.$$

The interpretation for the confidence interval for the average response when $x = x^*$ is:

The *prediction interval for an individual response* when $x = x^*$, is

$$\hat{y} \pm t_{n-2}^* \sqrt{MSE \left(1 + \frac{1}{n} + \frac{(x^* - \bar{x})^2}{(n-1)s_x^2} \right)}.$$

The interpretation for the prediction interval for an individual response when $x = x^*$ is:

We use *confidence* and *prediction* intervals to account for our uncertainty in making predictions using the regression equation.

Two facts about confidence and prediction intervals

1. As we predict further away from $\bar{x}$, the intervals get wider. For example, if $\bar{x} = 5$, a prediction interval for $x^* = 7$ will be smaller than $x^* = 10$.
2. The prediction interval for an individual response is always wider than the confidence interval for the average response.

Class Example 9.2.2 *NFL Rushing*

The tables below provide computer output for the regression line to predict number of rushing yards a player will have in a season based on their number of rushing attempts. The dataset contains all players in the NFL who attempted a rush from 2001 to 2023.

Coefficients	Estimate	Std. Error	t	p -value
(Intercept)	-4.424	0.782	-5.66	< 0.0001
Attempts	4.31	0.009	478.89	< 0.0001

Parameter estimates for regression of season rushing yards on number of rushes attempted

Source	df	SS	MS	F	p -value
Regression	1	7.14×10^8	7.14×10^8	211099	< 0.0001
Error	7514	2.54×10^7	3.38×10^3		
Total	7515	7.39×10^8			

ANOVA results for regression of season rushing yards on number of rushes attempted

- a) Is the number of rushing attempts a good predictor of the number of season rushing yards? Justify your answer using specific values in the output.
- b) Find and interpret r^2 for this regression.
- c) If a player has 300 rushing attempts in a season, how many rushing yards would you predict they would have?
- d) Suppose the 95% prediction interval for players that have 300 season rushing attempts is (1188.58, 1388.58). Provide an interpretation of this interval in the context of the question.

Class Example 9.2.3 *Predicting Mustang Prices*

The dataset MustangPrice contains information for a sample of 25 used Mustang cars offered for sale on a website. Suppose that we are interested in predicting the Price of a Mustang (in thousands of dollars) based on how old it is (Age in years). Some output for fitting this model is shown in the tables below. You may assume that any necessary conditions for inference hold.

Coefficients	Estimate	Std. Error	<i>t</i>	<i>p</i> -value
(Intercept)	30.264	3.440	8.80	< 0.0001
Age	-1.717	0.365	-4.71	< 0.0001

Parameter estimates for regression of mustang price on age in years

Source	df	SS	MS	F	<i>p</i> -value
Regression	1	1454.4	1454.4	22.15	< 0.0001
Error	23	1509.9	65.6		
Total	24	2964.3			

ANOVA results for regression of mustang price on age in years

- a) Is age of a used Mustang an effective predictor of its price? Justify your answer using the information from the output above.

- b) You are thinking about buying a 5 year-old Mustang, what would you predict the price to be?

- c) Two intervals are given below. One is the 95% confidence interval for mean price and one is the 95% prediction interval for individual prices, for Mustangs that are 5 years old. Which is which?
 (17.49, 25.86): _____ (4.40, 38.96): _____
 d) Provide an interpretation for both of the intervals given in c) above in the context of the question.

- e) The 95% prediction interval for Mustangs that are 10 years old is (9.51, 16.68). Is the average age of Mustangs in the dataset closer to 5 years or 10 years? Explain.

10.1 Probability Rules

“Probability theory is nothing but common sense reduced to calculation.” – Pierre-Simon Laplace

Key Concepts

- Compute the probability of events if outcomes are equally likely
- Identify when a probability question is asking for A and B, A or B, not A, or A if B
- Use the complement, additive, multiplicative, and conditional rules to compute probabilities of events
- Recognize when two events are disjoint

Equally Likely Outcomes

Recall that a *population* is the collection of all individuals or items under consideration. An individual could refer to a person, a playing card, or whatever object we are interested in. We typically refer to a population when referencing sampling. However, when we talk about experiments, we use the phrase *sample space*. The *sample space* is the set of all possible non-overlapping outcomes for a random experiment, and is denoted Ω . *Events* can be simple (e.g., the outcome of a single flip of a coin) or complex combinations of outcomes (e.g., the results from 10 flips of a coin). We have three important rules for probability:

P.1) Any probability must be between 0 and 1 (inclusive)

P.2) When outcomes are equally likely, the probability of the event E is given by

$$P(E) = \frac{N(E)}{N(\Omega)}$$

P.3) The sum of the probabilities for all of the experimental outcomes must equal 1

P.4) Suppose the event E is composed of several non-overlapping outcomes $E_1, \dots, E_n$, then

$$P(E) = P(E_1) + \dots + P(E_n)$$

P.5) $P(\text{not } A) = P(A^c)$ = Probability A does not occur is given by

$$P(A^c) = 1 - P(A)$$

The probability of an event is the long run frequency or proportion of times the event occurs. Probabilities are always between 0 and 1.

We use the notation $P()$ to mean *probability of*.

We use the notation $N()$ to mean *number of outcomes in* an event. When outcomes are equally likely, the probability of event is the number of outcomes in the event divided by the total number of outcomes.

Sample space, Ω , is the set of all possible outcomes.

The *complement* of a condition, written “not A” or A^c , includes events in which the condition does not happen.

Class Example 10.1.1

Suppose a fair, six-sided die is rolled twice. Determine the following.

a) $N(\Omega)$

b) $N(A)$, where A is the event that the sum of the two rolls is 5

c) $P(B)$, where B is the event that the two rolls are the same

d) $P(C)$, where C is the event that the sum of the two rolls is even

e) $P(D)$, where D is the event that the two rolls are not the same

Combinations of Events

Many of the terms we used for describing the relationship between two categorical variables is also used when describing events matching two (or more) conditions. In addition to the ideas of *joint*, *marginal*, and *conditional* proportions, we will use terms like *intersection*, and *union* of conditions.

- $P(A \text{ and } B) = P(A \cap B)$ = Probability conditions A and B both occur.
- $P(A \text{ or } B) = P(A \cup B)$ = Probability either condition A or condition B (or both) occur.

The following equations describe the probability of these complex events.

P.6) Additive Rule:

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

P.7) Multiplication Rule:

$$\begin{aligned} P(A \cap B) &= P(B|A) * P(A) \\ &= P(A|B) * P(B) \end{aligned}$$

Venn diagrams

Venn diagrams are visual displays that are useful for investigating *intersections*, *unions*, and *complements*. These diagrams are constructed by first drawing a rectangle, representing the sample space. Within this rectangle, overlapping circles are drawn for each of the conditions of interest.

The *joint* or *intersection* of two conditions, written "A and B" or $A \cap B$, includes events that satisfy *both* conditions.

The *union* of two conditions, written "A or B" or $A \cup B$, includes events that satisfy *either* or *both* conditions.

The *conditional occurrence* of condition A given the occurrence of condition B is written "A if B" or $A | B$.

Class Example 10.1.2

Recall the following example from Section 2.1. 169 college students were asked about relationship status and gender. The results are given in the table below. For each of the following, assume that a student is chosen at random from this sample.

	Female	Male	Total
In a relationship	32	10	42
It's complicated	12	7	19
Single	63	45	108
Total	107	62	169

Contingency table for relationship status and gender.

- a) What is the probability the student is in a relationship? Is this a joint, marginal, or conditional probability?
-
- b) What is the probability the student is in a relationship given the student is female? Is this a joint, marginal, or conditional probability?
-
- c) What is the probability the student is female given the student is in a relationship? Is this a joint, marginal, or conditional probability?
-
- d) What is the probability the student is a male and single? Is this a joint, marginal, or conditional probability?
-

Class Example 10.1.3

Suppose that in a particular group of people, 10% are rich, 5% are famous, and 3% are both. Use a Venn diagram, as well as the definitions of probability, to answer the following.

- a) What is the probability that a randomly selected person in this group is not rich?
-
- b) What is the probability that a randomly selected person in this group is rich but not famous?
-
- c) What is the probability that a randomly selected person in this group is neither rich nor famous?
-

Class Example 10.1.4

Police report that when stopping a driver on suspicion of driving while under the influence that 78% of drivers are given a breath test, 36% of drivers are given a blood test, and 8% of drivers are given neither. Use a Venn diagram, as well as the definitions of probability, to answer the following.

- a) What is the probability that a randomly selected person in this population is not given a breath test?
- b) What is the probability that a randomly selected person in this population is given a breath test but not a blood test?
- c) What is the probability that a randomly selected person in this population is given both tests if they are given at least one test?

Class Example 10.1.5

In 1974, the University of California, Berkeley was sued for bias against women who had applied for admission to graduate schools there. The table below displays the admissions data from the Fall 1973 semester.

	Male	Female	Total
Admitted	3546	1512	5058
Rejected	4896	2809	7705
Total	8442	4321	12763

Contingency table for UC Berkeley graduate admissions in Fall 1973.

- a) What is the probability a randomly selected applicant was female? Is this a joint, marginal, or conditional probability?
- b) What is the probability a randomly selected applicant was accepted given the applicant was female? Is this a joint, marginal, or conditional probability?
- c) What is the probability a randomly selected applicant was accepted given the applicant was male? Is this a joint, marginal, or conditional probability?

Special cases: Independent and Disjoint

There are two special cases of rules for probability *independent* and *disjoint*. If two events, A and B , are *independent* it means that the chance of A occurring is the same regardless of whether or not B happened. If two events, A and B , are *disjoint* it means that events A and B cannot occur at the same time.

P.8) If A and B are independent, then

a. $P(A | B) = P(A)$ b. $P(B | A) = P(B)$ c. $P(A \cap B) = P(A) * P(B)$

P.9) If A and B are disjoint, then $P(A \cap B) = 0$

Class Example 10.1.6

Suppose we have a fair coin.

a) Suppose we flip the coin two times, are the flips independent of one another?

b) Suppose that the first three flips were HHT. What is the probability the 4th flip is heads?

c) Suppose that the first three flips were TTT. What is the probability the 4th flip is tails?

d) What is the probability of the event the 4 flips were HHTH?

e) What is the probability of the event the 4 flips were TTTT?

f) Is getting a head on the first flip disjoint from getting a tail on the first flip?

g) Is getting a head on the third flip disjoint from getting a tail on the first flip?

Counting Rules

It is often useful to be able to count (enumerate) the number of possible outcomes of a random experiment. Understanding a couple of simple counting rules can help to determine probabilities of events.

C.1) *Basic Counting Rule*: If event A consists of ractions that are performed in a specific order, with m_1 possibilities for the first action, m_2 for the second, etc. Then the total outcomes in A is

$$N(A) = m_1 * m_2 * \cdots * m_r.$$

C.2) *Arranging items*: If event A has n items, then the number of ways we can arrange the items in A is

$$n! = n * (n - 1) * (n - 2) * \cdots * 2 * 1.$$

C.3) *Combinations*: If event A has n items, then the number of ways we can choose k items from A is

$$\binom{n}{k} = \frac{n!}{(n - k)! * k!}$$

The *basic counting rule* provides the number of possible outcomes of a random experiment in which several actions are performed.

A *combination* is any unordered collection elements chosen from a collection of elements.

Class Example 10.1.7

Suppose that a professor gives her students a 10 question True/False quiz.

a) How many ways are there to answer the quiz?

b) How many ways are there to answer exactly 5 questions correctly?

Class Example 10.1.8

Suppose that for a particular lottery ticket, you are asked to pick 6 numbers from the numbers 1-59. No matter what order with which you choose the 6 numbers, they are rearranged to be in numeric order.

a) How many different tickets can be purchased?

b) If all 6 numbers match the official values, you win the jackpot. What is the probability of winning the jackpot?

c) If 3 or more numbers match the official values, you win. What is the probability of choosing a winning ticket?

Class Example 10.1.9

Suppose that three glasses are filled with the same cola, and then labeled C, D, and P. A randomly selected person is asked to order the three glasses by preference (this person is unaware that all three glasses contain the same cola).

a) How many ways can the cola preferences be ranked?

b) What is the probability that C is ranked first?

c) What is the probability that C is ranked first and D is ranked last?

Additional Example Questions

Class Example 10.1.10

Use the definition of independent and disjoint events to determine whether the following pairs of events are independent, disjoint, both, or neither.

- a) Rolling doubles; Rolling a sum of 8

- b) Rolling a sum of 8; Getting a 2 on the first die rolled

- c) Rolling a sum of 7; Getting a 1 on the first die rolled

- d) Rolling a sum of 5; Getting doubles

Class Example 10.1.11

We have a data from a medical study with 210 people about smoker status and cancer status. The summary counts are given in the table below.

	Cancer	No Cancer	Total
Smoker	10	50	60
Non Smoker	6	144	150
Total	16	194	210

Cancer and smoker status.

- a) What is the probability of being a smoker if you have cancer? Is this a joint, marginal, or conditional probability?

- b) What is the probability of having cancer if you are a smoker? Is this a joint, marginal, or conditional probability?

- c) Is having cancer independent of being a smoker? Explain.

Class Example 10.1.12

We have information about blood pressure and cholesterol for American males. Suppose there is a 11% chance a man has both high cholesterol and high blood pressure, a 52% chance a man has neither high cholesterol or high blood pressure, and a 27% chance that a man has high blood pressure.

- a) What is the chance a man has high cholesterol?

- b) What is the chance a man has high cholesterol or high blood pressure?

- c) If a man does not have high blood pressure, what's the chance he has high cholesterol?

- d) Is having high blood pressure independent from having high cholesterol? Explain.

Class Example 10.1.13

Fifty-six percent of all American workers have an employer-sponsored retirement plan, 68% have employer-sponsored health insurance, and 49% have both benefits. We select a worker at random.

- a) What is the probability a worker has neither employer-sponsored health insurance nor a retirement plan?

- b) What is the probability a worker has employer-sponsored health insurance or an employer-sponsored retirement plan?

- c) What is the probability a worker has employer-sponsored health insurance if they have an employee-sponsored retirement plan?

- d) Are having these two benefits disjoint? Explain.

Class Example 10.1.14

Suppose that 23% of adults smoke cigarettes. It's known that 57% of smokers and 13% of nonsmokers develop a certain lung condition by age 60.

- a) What is the probability that a randomly selected 60-year-old has this lung condition?

- b) What is the probability that a randomly selected 60-year-old does not have this lung condition?

- c) What is the probability that a randomly selected 60-year-old is a smoker if he/she has this lung condition?

Class Example 10.1.15

Lie detectors are controversial instruments, barred from use as evidence in many courts. Nonetheless, many employers use lie detector screening as a part of their hiring process in the hope that they can avoid hiring people who might be dishonest. There has been some research, but no agreement, about the reliability of polygraph tests. Based on this research, suppose that a polygraph can detect 65% of lies, but incorrectly identifies 15% of true statements as lies.

A certain company believes that 95% of its job applicants are trustworthy. The company gives everyone a polygraph test, asking, "Have you ever stolen anything from your place of work?" Naturally, all the applicants answer "No," but the polygraph identifies some of those answers as lies, making the person ineligible for a job. What's the probability that a job applicant rejected under suspicion of dishonesty was actually trustworthy?

Summary of Probability and Counting Rules

Probability rules

P.1) $0 \leq P(A) \leq 1$

P.2) $P(A) = \frac{N(A)}{N(\Omega)}$

P.3) $P(\Omega) = 1$

P.4) If the event E is composed of several non-overlapping outcomes $E_1, \dots, E_n$, then $P(E) = P(E_1) + \dots + P(E_n)$

P.5) $P(A^c) = 1 - P(A)$

P.6) Addition Rule:

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

P.7) Multiplication Rule:

$$\begin{aligned} P(A \cap B) &= P(B | A) * P(A) \\ &= P(A | B) * P(B) \end{aligned}$$

P.8) If A and B are independent, then

a. $P(A | B) = P(A)$ b. $P(B | A) = P(B)$ c. $P(A \cap B) = P(A) * P(B)$

P.9) If A and B are disjoint, then $P(A \cap B) = 0$

Counting rules

C.1) Basic counting rule: If event A consists of r actions, and for each action there are $m_1, m_2, \dots, m_r$ possibilities, then the number of outcomes in A is

$$N(A) = m_1 * m_2 * \dots * m_r$$

C.2) Arranging items: If event A has n items, then the number of ways we can arrange the items in A is

$$n! = n * (n - 1) * (n - 2) * \dots * 2 * 1$$

C.3) Choosing items: If event A has n items, then the number of ways we choose k items from A is

$$\binom{n}{k} = \frac{n!}{(n - k)! * k!}$$

- $P(A)$ is the probability of event A
 - Ω is the sample space, and includes all possible outcomes - $N(A)$ is the number of outcomes in event A
 - $P(A^c) = P(\text{not } A)$
 - $P(A \cup B) = P(A \text{ or } B)$
 - $P(A \cap B) = P(A \text{ and } B)$
 - $P(A | B) = P(A \text{ if } B)$

10.2 Binomial Probability

A binomial random variable is a special kind of random variable that only has two possible outcomes for each trial, a fixed number of trials, and each trial has the same probability of success.

Key Concepts

- Identify a binomial random variable
- Compute probabilities for a binomial random variable
- Compute the mean and/or standard deviation for a binomial random variable

Binomial Random Variables

Random variables often follow simple, predictable patterns. In these cases, calculating probabilities and the mean or standard deviation are much simpler and do not require a table to be created. One type of random variable with a simple pattern is called a “Binomial Random Variable.” Suppose we call the occurrence of an event a *success*. “Success” does not necessarily indicate the outcome was positive (e.g., if we are investigating cancer in a sample of mice, success could be defined as the presence of cancer). A *binomial random variable* measures the number of successes in a fixed number of trials.

A *binomial random variable* measures the number of successes in a fixed number of trials.

In order to be considered a binomial random variable, the following properties must hold:

1. We must be measuring the number of successes in n independent trials
2. The number of trials, n , is fixed in advance
3. The probability of success, p , does not change from trial to trial
4. Each trial is independent of the others (the outcome of one trial does not affect the outcome of another trial)

Class Example 10.2.1

For each of the following, determine if the random variable is binomial. If it is, determine n and p .

- a) On average, 25% of La Crosse residents recycle. 500 residents of La Crosse were asked if they recycle. A is the number of ‘yes’ answers in the sample.
- b) A box contains 50 marbles, 35 green and 15 white. Fifteen marbles are selected without replacement. B is the number of white marbles in the sample.
- c) An American roulette wheel consists of 18 red, 18 black, and 2 green numbers. Suppose you observed red on 10 consecutive spins. J is the number of additional spins needed until you observe black.
- d) D is the number of 5s in ten rolls of a fair die.
- e) Customers arrive at Alice’s with a rate of 5 per hour. E is the number of customers that enter Alice’s between 2 A.M. and 4 A.M.

Binomial Probabilities, Mean, and Standard Deviation

When a random variable is binomially distributed, probabilities and other characteristics can be expressed as formulas. For a binomial random variable X with n trials and probability of success p ,

$$P(X = k) = \binom{n}{k} p^k (1 - p)^{n-k} \text{ for } k = 0, 1, 2, \dots, n. \quad \text{- Mean: } \mu = E[X] = np$$

$$\text{Standard Deviation: } \sigma = sd(X) = \sqrt{np(1 - p)}$$

Class Example 10.2.2

To test for ESP, we have 4 cards, each with different symbols. These cards are shuffled and one is chosen at random (symbol down). Each time, you guess the symbol on the card. Suppose that you repeat this test 8 times and you do not have ESP.

- a) Let X be the number of times you guess correctly. Is X binomial? If so, what are n and p ?

- b) What is the expected value of X ?

- c) What is the standard deviation of X ?

- d) To be certified as having ESP, you need to correctly identify at least 6 cards of the 8 drawn. What is the probability you get certified?

Class Example 10.2.3

Joan takes 10 phone calls per day from customers, and, on average, 30% of these callers are angry. If 7 or more of these callers in a given day are angry, Joan has a bad day.

- a) Let C be the number of angry callers in a day. Is C binomial? If so, what are n and p ?

- b) How many angry callers will Joan speak to on average each day? What is the standard deviation for the number of angry callers?

- c) What is the probability that Friday is a good day?

- d) What is the probability that 3 of 5 days in her workweek are good?

Class Example 10.2.4

In the 2023 NFL season, Jordan Love threw 579 passes. A fair-weather fan happened to only see a few games that season. The fan only saw Jordan Love throw 33 times, and of those 18 were completed (caught).

a) What would the fan estimate is Jordan Love's probability to complete a pass?

b) Based on the probability in part (a), how many passes would the fan expect Jordan Love to complete in the next 10 throws?

10.3 Data Sources

The datasets used in this course pack come from the following sources. We gratefully acknowledge the original authors and data collectors.

Lock5Data R Package (GPL-2)

The following datasets are from the **Lock5Data** R package, which accompanies *Statistics: Unlocking the Power of Data* by Lock, Lock, Lock, Lock, and Lock (Wiley). The Lock5Data package is distributed under the GPL-2 license.

- **CAOSExam.csv** — Pre-test and post-test scores from a Comprehensive Assessment of Outcomes in Statistics (CAOS) exam. 10 students, 3 variables.
- **FloridaLakes.csv** — Water quality measurements from 53 Florida lakes. Original source: Lange, Royals, and Connor (1993), *Journal of Environmental Quality*.
- **PulseRates.csv** — Resting pulse rates for 10 students before and after exercise.
- **SleepStudy.csv** — Sleep habits and academic performance for 253 college students. Original source: Onyper, Thacher, Gilbert, and Gradess (2012), *Teaching of Psychology*.
- **USStates.csv** — State-level data (50 US states) compiled from US government sources including the Census Bureau and Bureau of Labor Statistics.

R Packages (GPL-2+)

- **Baumann.csv** — Reading comprehension scores from a study comparing three teaching methods (Basal, DRTA, Strat). 66 observations. From the **carData** R package (GPL-2+). Original source: Baumann and Jones (1992).
- **InsectSprays.csv** — Insect counts for six different spray treatments. 72 observations. Built-in R dataset (GPL-2/3). Original source: Beall (1942).

Public Domain / Custom

- **HollywoodMovies2023.csv** — Opening weekend revenue and world gross for six 2023 Hollywood films. Created for this course pack from publicly available box office data (Box Office Mojo).
- **MastersWinners.csv** — Winning scores at the Masters Tournament (1934–2024). 87 observations. Compiled from public tournament records.
- **NBASalaries2022.csv** — NBA player salaries for the 2022–23 season. 467 observations. Synthetic data generated to match the published salary distribution.
- **Salaries.csv** — Monthly sales figures (in dollars) for 15 salespeople under three pay structures (Commission, Fixed Salary, Commission Plus Salary). Synthetic data created for this course pack to illustrate one-way ANOVA and post-hoc comparisons.