Name _____ _____ Baggett/Fall2026/207/Quiz6

first name last name

No Calculators.

**Questions 1 and 2.** Circle **True** or **False**. Then, in **one sentence**, say why — if the statement is false, the fastest way is usually to give an example that breaks it. **1 point for the circle, 1 for the reason.**

**1.** (**2** pts) In a related rates problem you differentiate the relating equation with respect to time *before* substituting the given numerical values.

**True** **False**

Why? **Substituting first freezes the variables, and a constant has derivative zero – the relationship you needed is destroyed.**

/2

**2.** (**2** pts) $\dfrac{d}{dx}(\ln 10) = \dfrac{1}{10}$

**True** **False**

Why? $\ln 10$ **is a constant, so the derivative is** $0$**. The rule** $\frac{d}{dx}\ln x = \frac{1}{x}$ **needs a variable inside.**


/2

**3.** (**6** pts) **Derive** $\dfrac{d}{dx}[\arcsin x]$ using implicit differentiation. **Show every step.**

You may use the derivative rules for sin and cos, but **not** any derivative rule for an inverse trig function. Quoting the formula earns nothing.

$$\frac{d}{dx}[\arcsin x] = \frac{1}{\sqrt{1-x^2}}$$

| | |
|---|---|
| $y = \arcsin x$ | name it |
| $\sin y = x$ | **rewrite** — no inverse left in it |
| $\cos y \dfrac{dy}{dx} = 1$ | **differentiate** both sides |
| $\dfrac{dy}{dx} = \dfrac{1}{\cos y}$ | solve — still in $y$, so not finished |
| $\cos y = \sqrt{1-x^2}$ | **convert**: triangle, or $\cos y = \sqrt{1-\sin^2 y}$ |
| $\dfrac{dy}{dx} = \dfrac{1}{\sqrt{1-x^2}}$ | substitute |




**Step 5 in a picture.** With $\sin y = x = \frac{x}{1}$, label the **opposite** side $x$ and the **hypotenuse** 1. Pythagoras gives the adjacent side $\sqrt{1-x^2}$, and $\cos y$ is adjacent over hypotenuse.

The identity $\cos y = \sqrt{1-\sin^2 y}$ reaches the same place.

/6

**4.** (**3** pts) Find $\dfrac{dy}{dx}$ by implicit differentiation:

$$x \sin y = y$$

$\dfrac{dy}{dx} = \dfrac{\sin y}{1 - x \cos y}$

| | |
|---|---|
| $\dfrac{d}{dx}[x \sin y] = \dfrac{d}{dx}[y]$ | differentiate both sides |
| $\sin y + x \cos y \dfrac{dy}{dx} = \dfrac{dy}{dx}$ | **Product Rule**; $\sin y$ needs the **Chain Rule** |
| $\sin y = \dfrac{dy}{dx} - x \cos y \dfrac{dy}{dx}$ | gather the $\frac{dy}{dx}$ terms on one side |
| $\sin y = \dfrac{dy}{dx}(1 - x \cos y)$ | factor |
| $\dfrac{dy}{dx} = \dfrac{\sin y}{1 - x \cos y}$ | divide |

/3

**5.** (**2** pts) Differentiate $y = 5^{x^2}$. You do not need to simplify.

$y' = 5^{x^2} (\ln 5)(2x)$

| | |
|---|---|
| $\dfrac{d}{dx}[a^u] = a^u (\ln a) \dfrac{du}{dx}$ | the any-base rule, with $a = 5,\ u = x^2$ |
| $y' = 5^{x^2} (\ln 5)(2x)$ | **Chain Rule**: $\frac{du}{dx} = 2x$ |

/2

**6.** (**3** pts) Use **logarithmic differentiation** to find $\dfrac{dy}{dx}$ for

$$y = x^{2x}, \qquad x > 0.$$

$\dfrac{dy}{dx} = x^{2x}(2 \ln x + 2)$

| | |
|---|---|
| $\ln y = \ln\left(x^{2x}\right)$ | take ln of both sides |
| $\ln y = 2x \ln x$ | $\ln(A^p) = p \ln A$ |
| $\dfrac{1}{y}\dfrac{dy}{dx} = 2 \ln x + 2x \cdot \dfrac{1}{x}$ | differentiate; **Product Rule** on the right |
| $\dfrac{1}{y}\dfrac{dy}{dx} = 2 \ln x + 2$ | simplify |
| $\dfrac{dy}{dx} = y(2 \ln x + 2) = x^{2x}(2 \ln x + 2)$ | multiply by $y$, then put $y$ back |

/3

**7.** (**2** pts) Each edge of a cube is growing at 2 cm/s. Let $s$ be the edge length and $V$ the volume.

Write the relating equation and **differentiate both sides with respect to $t$. Do not substitute any numbers and do not solve.**

$V = s^3$ **and** $\dfrac{dV}{dt} = 3s^2 \dfrac{ds}{dt}$

$V = s^3$ the relating equation — true at **every** instant

$\dfrac{dV}{dt} = 3s^2 \dfrac{ds}{dt}$ **Chain Rule**: $s$ moves, so it carries its own rate

/2