

first name

last name

No Calculators.

Questions 1 and 2. Circle **True** or **False**. Then, in **one sentence**, say why — if the statement is false, the fastest way is usually to give an example that breaks it. **1 point for the circle, 2 for the reason.**

1. (3 pts) $\lim_{x \rightarrow 1} \frac{x-3}{x^2+2x-4} = \frac{\lim_{x \rightarrow 1}(x-3)}{\lim_{x \rightarrow 1}(x^2+2x-4)}$

☒ True☐ False

Why? **Here the denominator limit is $-1 \neq 0$, so the Quotient Law does apply.**

/3

2. (3 pts) If $f(0.1)$, $f(0.01)$ and $f(0.001)$ all round to 2, then $\lim_{x \rightarrow 0} f(x) = 2$.

☒ True☐ False

Why? **A table samples finitely many points and says nothing about the rest. $f(x) = \sin(\frac{\pi}{x}) + 2$ hits exactly 2 at every one of those inputs while oscillating wildly in between.**

Where points were lost. The most common answer was that we are only told about x approaching 0 from the *right*. That is true, but it is not what breaks the statement — fix it by listing $f(-0.1)$, $f(-0.01)$, $f(-0.001)$ as well and the statement is *still* false. Three points are three points, and a limit is a claim about **all** x near 0. The function above passes through exactly 2 at every listed input while swinging between 1 and 3 everywhere in between. No finite table is evidence for a limit — which is also why you were asked to estimate signs rather than plug in values.

/3

3. (5 pts) Let

$$g(x) = \begin{cases} 2x+1, & x < 1 \\ 5, & x = 1 \\ 6-3x, & x > 1 \end{cases}$$

(a) $\lim_{x \rightarrow 1^-} g(x) = \underline{\quad 3 \quad}$ (b) $\lim_{x \rightarrow 1^+} g(x) = \underline{\quad 3 \quad}$

(c) $\lim_{x \rightarrow 1} g(x) = \underline{\quad 3 \quad}$ (d) $g(1) = \underline{\quad 5 \quad}$

(e) Is g continuous at $x = 1$? If not, say which requirement fails.

No. The limit exists and $g(1)$ exists, but they disagree — requirement 3.

/5

4. (4 pts) Evaluate the limit. Show the rewriting — substituting first and finding $\frac{0}{0}$ is not an answer, it is the reason a rewrite is needed.

$$\lim_{x \rightarrow 0} \frac{\sqrt{x+4} - 2}{x}$$

$\frac{1}{4}$ — **multiply by the conjugate** $\frac{\sqrt{x+4}+2}{\sqrt{x+4}+2}$:

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{\sqrt{x+4} - 2}{x} &= \lim_{x \rightarrow 0} \frac{(\sqrt{x+4} - 2)(\sqrt{x+4} + 2)}{x(\sqrt{x+4} + 2)} \\ &= \lim_{x \rightarrow 0} \frac{(x+4) - 4}{x(\sqrt{x+4} + 2)} = \lim_{x \rightarrow 0} \frac{x}{x(\sqrt{x+4} + 2)} \\ &= \lim_{x \rightarrow 0} \frac{1}{\sqrt{x+4} + 2} = \frac{1}{\sqrt{4} + 2} = \frac{1}{4} \end{aligned}$$

Where points were lost. Two things, and the first cost a point on its own. **(1) Keep lim on every line until you actually take it.** Many of you dropped it after the first step. Without it you have written that a *function* equals a *number*, which is false — and it is not bookkeeping: the cancellation $\frac{x}{x} = 1$ is only legal because $x \neq 0$, and the only thing on the page saying $x \neq 0$ is the $\lim_{x \rightarrow 0}$. Take the symbol away and you have divided by zero. That is the whole reason 2.3 kept insisting on it. **(2) Algebra.** The conjugate gives $(\sqrt{x+4})^2 - 2^2 = (x+4) - 4 = x$ on top. Common slips were leaving $x+4-2$, squaring only part of the root, or cancelling the x against the 4 inside it.

/4

5. (3 pts) Let $f(x) = \frac{x+1}{x-3}$. **No calculator.** Decide each limit by naming the sign and size of the numerator and of the denominator.

(a) $\lim_{x \rightarrow 3^-} f(x) = \underline{\quad -\infty \quad}$ (b) $\lim_{x \rightarrow 3^+} f(x) = \underline{\quad +\infty \quad}$

(c) Write the equation of the vertical asymptote.

$x = 3$

/3

Just left of 3 the numerator is near 4 (positive) and $x-3$ is small and **negative**, so the quotient is large and negative. Just right of 3 the denominator is small and positive.

6. (2 pts) Find the value of the constant c that makes

$$h(x) = \begin{cases} x^2 - 1, & x < 3 \\ cx + 2, & x \geq 3 \end{cases}$$

continuous at $x = 3$.

/2

Both pieces are polynomials, so each is continuous on its own side and the only place continuity can fail is the junction. The pieces therefore just have to meet at $x = 3$:

$$3^2 - 1 = 3c + 2 \quad \implies \quad 8 = 3c + 2 \quad \implies \quad \mathbf{c = 2}$$