

EJP 52570

NNC-112¹, NNC-687 and NNC-756¹, new selective and highly potent dopamine D₁ receptor antagonists

Peter H. Andersen^a, Frederik C. Grønvald^b, Rolf Hohlweg^b, Louis B. Hansen^b, Erling Guddal^b,
Claus Braestrup^c and Erik B. Nielsen^d

From the Department of Molecular Pharmacology, ^a Biosciences, and the ^d Departments of CNS Pharmacology
and ^b Medicinal Chemistry Laboratory I, ^c CNS Division, Novo Nordisk, DK-2880 Bagsvaerd, Denmark

Received 28 November 1991, revised MS received 29 April 1992, accepted 19 May 1992

The neurochemical properties of three novel benzazepine derivatives NNC-112, NNC-687 and NNC-756 were assessed. These compounds inhibited dopamine D₁ receptor binding in vitro with low nanomolar to picomolar dissociation constants whereas those for the D₂ receptor were in the micromolar range. Contrary to classical neuroleptics, but similar to the atypical neuroleptics, clozapine and fluperlapine, NNC-112, NNC-687 and NNC-756 were relatively more potent in inhibiting dopamine-stimulated adenylyl cyclase than [³H]SCH 23390 binding. Both NNC-112 and NNC-756 had high affinity for the 5-HT₂ receptor whereas NNC-687 had low affinity for this receptor. The affinity for other receptors or neurotransmitter transporters was very low. In vivo, the dopamine D₁ receptor selective profile of NNC-112, NNC-687 and NNC-756 was evident from the potent inhibition of D₁ receptor binding whereas no effect on D₂ receptor binding was apparent. In addition, the compounds blocked D₁ receptor-mediated rotation in unilaterally 6-hydroxydopamine-lesioned rats, but had no effect on D₂-induced rotation. Thus, NNC-112, NNC-687 and NNC-756 are potent and selective dopamine D₁ receptor antagonists that may be useful in the treatment of schizophrenia.

Dopamine D₁ receptors; NNC-112; NNC-687; NNC-756; Binding (in vivo)

1. Introduction

Until recently, dopamine receptors were classified into D₁ and D₂ types (see recent review by Andersen et al., 1990). Previously, D₁ receptors were studied primarily with respect to their function. However, following the identification of SCH 23390 as a D₁-selective ligand (Hyttel, 1983), [³H]SCH 23390 has been used to identify dopamine D₁ receptors both in vitro and in vivo (Billard et al., 1984; Andersen et al., 1985; Andersen and Grønvald, 1986). The use of this radioligand resulted in a number of interesting findings. First, in vitro studies indicated that the D₁ receptor apparently exists in two states or subtypes (Andersen and Braestrup, 1986; see also Seeman et al., 1985). Only one of these subtypes is associated with the adenylyl cyclase whereas the other may be linked to phosphoinositol turnover (Andersen et al., 1990; Monsma et al.,

1990). Second, atypical neuroleptics of the clozapine type have some selectivity for blocking the adenylyl cyclase-coupled D₁ receptor, whereas classical neuroleptics are more potent on the other state or subtype (Andersen and Braestrup, 1986). Third, in vivo studies have indicated that the dopamine D₁ receptor is an important target for atypical neuroleptic drugs of the clozapine type and, interestingly, a derivative of clozapine, fluperlapine, appears to be D₁-selective (Andersen et al., 1986, 1988). The importance of the D₁ receptor as an in vivo target for clozapine has been further corroborated by recent PET-scan studies in humans (Farde et al., 1989). Together, these data suggest that, at least in part, the atypical profile of clozapine is due to antagonism of the dopamine D₁ receptor and, perhaps, of a specific subpopulation of D₁ receptors. However, it should also be mentioned that the D₄ receptor has also been found to be an important target for clozapine (Van Tol et al., 1991).

Because of continuing interest in compounds with a clozapine-like dopaminergic profile, we synthesized a number of analogues of SCH 23390. In the present paper, we describe the neurochemical and behavioural profile of three such novel dopamine D₁ receptor

Correspondence to: P.H. Andersen, Department of Molecular Pharmacology, Biosciences, Novo Nordisk, Novo Allé, DK-2880 Bagsvaerd, Denmark. Tel. 45.44 44 88 88, ext. 3522, fax 45.42 98 50 07.

¹ Previously NO-112 and NO-756, respectively.




Fig. 1. Structure of NNC-112, NNC-756 and NNC-687.

antagonists, NNC-112, NNC-687 and NNC-756 (previous names, NO-112 and NO-756, respectively) (see fig. 1). Some of these data have been presented at the 18th Annual Neuroscience Meeting (Andersen et al., 1988; Nielsen et al., 1988).

2. Materials and methods

2.1. Animals

Male Albino Wistar rats (Møllegaard's Breeding Labs, Lille Skensved, Denmark), weighing 175 ± 25 g were used. The animals were housed in group cages (four to eight animals/cage) placed in rooms with a 12-h light cycle (06:00-18:00). The room temperature varied between 19–24°C and the relative humidity varied from 30 to 60%.

Male Albino mice (NMRI type, bred at Novo Nordisk or from Møllegaard) weighing 20 ± 2 g were also used. They were housed in group cages under the same general conditions as outlined for rats (above). All animals had access to food (standard lab chow) and tap water ad libitum.

2.2. Radioligands

[³H]SCH 23390 (73.4 Ci/mmol), [³H]2-(N,N-di-2,3(n)-propylamino)-8-hydroxy-1,2,3,4-tetrahydronaphthalene([³H]8-OH-DPAT) (176 Ci/mmol), [³H]prazosin (79 Ci/mmol), 1-[³H]quinuclidinyl-benzilate (QNB) (39 Ci/mmol), 1-[³H]noradrenaline (36.6 Ci/mmol), [³H]serotonin (5-HT) (20 Ci/mmol), [³H]dopamine (49.9 Ci/mmol), [³H]choline (75 Ci/mmol), [³H]RX 781094 (40 Ci/mmol), [³H]pyrilamine (26 Ci/mmol) and [³H]dihydroalprenolol (DHA) (73 Ci/mmol) were obtained from Amersham, U.K., and [³H]muscimol (36.9 Ci/mmol), [³H]diazepam (70 Ci/mmol), [³H]spiperone (29.6 Ci/mmol) and [³H]ketanserin (95 Ci/mmol) were obtained from NEN, Boston, MA, U.S.A.

2.3. Drugs

The following drugs were used: methylphenidate (Ciba-Geigy, Copenhagen, Denmark); pergolide (Eli

Lilly, Indianapolis, IN); cis-flupentixol, trans-flupentixol and citalopram (H. Lundbeck A/S, Copenhagen, Denmark); l-propranolol (ICI, Copenhagen, Denmark); haloperidol, spiperidol, domperidone and ketanserin (Janssen, Beerse, Belgium); apomorphine (Nordisk Droge, Copenhagen, Denmark); (+)-8-chloro-7-hydroxy-3-methyl-5-(7-benzofuranyl)-2,3,4,5-tetrahydro-1H-3-benzazepine (NNC-112) and (–)-8-chloro-7-hydroxy-3-methyl-5-(7-benzofuranyl)-2,3,4,5-tetrahydro-1H-3-benzazepine ((–)-NNC-112), (+)-8-nitro-7-hydroxy-3-methyl-5-(7-(2,3-dihydrobenzofuranyl))-2,3,4,5-tetrahydro-1H-3-benzazepine (NNC-687) and (+)-8-chloro-7-hydroxy-3-methyl-5-(7-(2,3-dihydrobenzofuranyl))-2,3,4,5-tetrahydro-1H-3-benzazepine (NNC-756) were synthesized at Novo Nordisk A/S; prazosin and thiothixene (Pfizer, Copenhagen, Denmark); R- and S-sulpiride (Ravizza Laboratories, Milan, Italy); SKF 38393 (Research Biochemicals, Natick, MA); fluperlapine and clozapine (Sandoz, Copenhagen, Denmark); (+)-8-chloro-7-hydroxy-3-methyl-5-phenyl-2,3,4,5-tetrahydro-1H-3-benzazepine (SCH 23390) (Schering Corp., Bloomfield, NJ) and RS 21361193 (Syntex Research Lab., Edinburgh, U.K.). All other drugs and chemicals used were of the best quality commercially available.

2.4. Drug dissolution

For all in vitro assays, stock solutions were prepared in water whenever possible, otherwise in methanol. Further dilution was made in water to obtain a maximal concentration of 0.1% methanol in any assay. By itself, this concentration of methanol had no effect in any of the assays.

For in vivo assays, water-soluble compounds were dissolved in 0.9% NaCl and used directly. Water-insoluble compounds were dissolved in 1 M tartaric acid + propyleneglycol (1:1) and brought up to volume with 0.9% saline.

2.5. Tissue preparation for D₁ receptor binding and adenylyl cyclase assays

Rat striatal tissue was processed as described by Andersen et al. (1985). Briefly, the tissue was homogenized gently by hand using a glass-teflon homogenizer (10 strokes) in 100 volumes of 10 mM imidazole-HCl (pH 7.4 at 25°C) containing 2 mM EDTA and centrifuged at $25\,000 \times g$ for 20 min at 4°C. The pellet was resuspended in 100 volumes of the same buffer and the homogenization/centrifugation step was repeated a total of 3 times. The final pellet was resuspended in 100 volumes of 2 mM imidazole-HCl (pH 7.4 at 25°C) containing 2 mM EDTA and used immediately for binding or adenylyl cyclase assays.

2.6. *In vitro* [^3H]SCH 23390 binding

A mixture of 100 μl of the above tissue suspension, 600 μl of 16.67 mM imidazole-HCl (pH 7.4 at 25°C) containing 16.67 mM theophylline, 1 mM EGTA and 10 mM MgSO_4 , 100 μl [^3H]SCH 23390 (final concentration 0.2 nM), and 200 μl of water/test compound/1 μM cis-flupentixol (control/compound testing/non-specific binding) were incubated at 30°C for 60 min followed by rapid filtration through Whatman GF/B filters under vacuum. The filters were immediately washed with 2×10 ml ice-cold 0.9% NaCl and transferred to scintillation vials. Four millilitres of scintillation fluid was added and the radioactivity retained on the filters was counted in a conventional scintillation counter.

2.7. Adenylyl cyclase assay

A mixture of 50 μl tissue suspension (see section 2.5.), 300 μl of 16.67 mM imidazole-HCl (pH 7.4 at 25°C) containing 16.67 mM theophylline, 1 mM EGTA, and 10 mM MgSO_4 was combined with 50 μl of agonist solution and 50 μl of GTP (final concentration 15 μM). This mixture was preincubated for 15 min at 0°C, whereupon 50 μl of ATP (final concentration 1.5 mM) was added and the samples were incubated for 5 min at 30°C. The reaction was terminated by boiling the mixture for 3 min followed by centrifugation at $2800 \times g$ for 20 min. To ensure equilibrium, the effects of certain drugs (as indicated in legends) were retested using the same procedure but with a 1-h incubation at 30°C. Identical results were obtained as shown previously (Andersen and Nielsen, 1986), indicating that equilibrium is established even with short incubation times. Cyclic AMP (cAMP) was measured in the clear supernatant as described by Geisler et al. (1977). In brief, 50 μl of the supernatant was incubated for 90 min at 4°C with 50 μl of 1.25 pmol [^3H]cAMP, 50 μl of 50 mM Tris-HCl (pH 7.4 at 4°C) containing 30 mM EDTA and

1 mM mercaptoethanol and 50 μl of purified cAMP-dependent protein kinase. One millilitre of 70% $(\text{NH}_4)_2\text{SO}_4$ was added and the incubation was continued for 15 min at 4°C. The incubation mixtures were then centrifuged at $2800 \times g$ for 30 min at 4°C and decanted. After addition of 0.1 ml H_2O and 1.2 ml scintillation fluid, radioactivity was determined by conventional scintillation counting. Standards containing 0–8 pmol cAMP/50 μl were assayed in parallel.

2.8. [^3H]Spiroperidol binding

As described previously by Andersen et al. (1985), rat striatal tissue was homogenized in 2×10 ml 50 mM Tris-HCl (pH 7.4 at 30°C) containing 120 mM NaCl and 4 mM MgCl_2 and centrifuged for 10 min at $40000 \times g$. The pellet was resuspended in 1000 volumes (original wet weight) and used immediately for D_2 receptor binding assay. The assay mixture consisted of 2.5 ml tissue suspension, 25 μl [^3H]spiroperidol (final concentration 0.05 nM) and 25 μl H_2O /test compound/100 nM domperidone (control/compound testing/non-specific binding). The incubation was carried out at 30°C for 20 min followed by 10 min on ice. The reaction was terminated by rapid filtration through Whatman GF/B filters followed by a 2×10 ml wash with ice-cold 0.9% NaCl. Radioactivity trapped on the filters was measured by conventional scintillation counting.

2.9. *In vivo* binding

Mice were injected i.v. into the tail vein with the radioligand. After 5–20 min the mice were decapitated and the brain (minus the cerebellum) of each mouse was homogenized in 14 ml of 50 mM potassium-phosphate buffer (pH 7.1 at 25°C) or in 50 mM Tris-citrate (pH 7.4 at 30°C) containing 120 mM NaCl and 4 mM MgCl_2 . For further details see table 1.

For further details concerning D_1 and D_2 binding, see Andersen (1988).

TABLE 1

Conditions for *in vivo* receptor binding.

	Receptor type		
	D_1	D_2	5-HT ₂
Ligand	[^3H]SCH 23390	[^3H]Raclopride	[^3H]Ketanserin
Ci/mouse ^a	4 μCi	4 μCi	4 μCi
Decapitation time after ligand injection	20 min	5 min	10 min
Non-specific binding defined with	20 mg/kg cis-flupentixol i.p. 2 h	20 mg/kg cis-flupentixol i.p. 2 h	100 mg/kg methysergide i.p. 1 h
Level of non-specific binding	5–15%	5–15%	25–35%

^a I.v. injection volume was 200 μl .

TABLE 2

Effects of NNC-112, NNC-687 and NNC-756 on dopamine-stimulated adenylyl cyclase and D₁, D₂ and 5-HT₂ receptors. A comparison with selected reference compounds.

The results shown are mean values from 2 to 11 separate experiments performed in duplicate, using 2–14 separate concentrations in each experiment; > means less than 10% inhibition of specific binding at the concentration indicated. K_i values calculated from IC₅₀ values (see below) and with certain compounds confirmed from Scatchard analysis ^c. K_i ^a values obtained from Schild analysis. ^c agonist; ^b identical K_i values obtained with a 1-h incubation. ^d Pharmacologically inactive isomer. IC₅₀ converted to K_i values using $K_i = IC_{50} / (1 + [L]/K_D)$, where IC₅₀ is the concentration of inhibitor causing 50% inhibition of the specific binding, [L] is the radioligand concentration used and K_D the dissociation constant of the radioligand. Some of these data have been published in Andersen et al. (1988); Andersen and Nielsen (1986); Andersen et al. (1985).

Compound	K _i (nM)			
	Adenylyl cyclase ^a	D ₁ ([³ H]SCH 23390)	D ₂ ([³ H]spiperone)	5-HT ₂ ([³ H]ketanserin)
Benzazepines				
NNC-112	2.8 ^b	0.18 ^c	898 ^c	18 ^c
(–)NNC-112 ^d	NT	170	988	NT
NNC-687	9.1	5.8 ^c	> 10 000	355
NNC-756	2.2	0.17 ^c	942 ^c	4.5 ^c
SCH 23390	39.9 ^b	0.14 ^c	895 ^c	37 ^c
SCH 23388 ^d	NT	173	900	> 10 000
SKF 38393 ^c	415 (EC ₅₀)	18	9 300	> 10 000

2.10. Other *in vitro* assays

Other neurotransmitter receptor binding assays or neurotransmitter uptake assays used in the present study have been described and characterized in detail in Andersen (1989).

2.11. Inhibition of rotational behaviour in 6-hydroxy-dopamine (6-OHDA)-lesioned rats

Rats were stereotactically injected with 9.7 µg of 6-OHDA into the right substantia nigra, using conven-

tional procedures. Upon recovery from surgery, the animals were injected with 0.25 mg/kg s.c. of apomorphine. Only animals exhibiting strong contralateral turning were used in further experiments. In these experiments, rotation dependent upon D₁ receptor stimulation was elicited by 2.5 mg/kg s.c. of SKF 38393. D₂-dependent rotations were produced by s.c. administration of pergolide (0.05 mg/kg).

Test drugs were injected at the indicated times (below) before the agonists. After agonist administration, the animals were attached to a swivel via a Velcro band strapped around the waist and placed in plexi-

TABLE 3

Results from *in vitro* receptor profiling of NNC-112, NNC-687, NNC-756 and selected reference compounds.

Results are means of two to six separate experiments performed in duplicate with 2–14 different concentrations in each assay. > indicates less than 10% inhibition of binding at the concentration indicated.

Receptor/ligand	IC ₅₀ (nM)					
	NNC-112	NNC-687	NNC-756	SCH 23390	cis-Flupentixol	Haloperidol
Dopamine receptors						
D ₁ [³ H]SCH 23390	0.4	12.4	0.4	0.4	0.72	150
D ₂ [³ H]spiperone	1857	> 15 000	1 938	1 800	0.75	5
Adrenoceptors						
α ₁ [³ H]prazosin	2 300	> 10 000	1 560	10 000	5.2	50
α ₂ [³ H]RX 781094	369	> 1 000	98	891	695	> 10 000
β [³ H]DHA	> 10 000	> 100 000	> 10 000	> 10 000	> 10 000	> 10 000
Serotonin receptors						
5-HT _{1A} [³ H]8-OH-DPAT	1 500	> 10 000	2 600	768	345	1 318
5-HT ₂ [³ H]ketanserin	37	663	7	63	7	175
GABA–benzodiazepine complex						
GABA _A [³ H]muscimol	> 10 000	> 10 000	> 10 000	> 10 000	> 10 000	> 10 000
Benzodiazepine [³ H]diazepam	> 10 000	> 10 000	> 10 000	> 10 000	> 10 000	> 10 000
Histamine receptors						
H ₁ [³ H]pyrilamine	1 800	1 800	1 400	1 400	963	1 400

TABLE 4

Effects of NNC-112, NNC-687, NNC-756 and selected reference compounds on neurotransmitter uptake.

Results are means of two to four separate experiments performed in duplicate with two to eight different concentrations of each compound in each assay. > indicates less than 10% inhibition of uptake at the concentration indicated.

Uptake	IC ₅₀ (nM)					
	NNC-112	NNC-687	NNC-756	SCH 23390	cis-Flupentixol	Haloperidol
Dopamine	4300	> 10000	> 10000	9600	2200	5900
Noradrenaline	790	2300	> 10000	780	4100	4600
Serotonin	835	> 1000	1500	124	> 1000	> 1000
GABA	> 10000	> 10000	> 10000	> 10000	7500	> 10000

glass bowls (diameter 49 cm). A computer detected rotation-operated photocell interruptions. Three to six animals were tested at each dose of antagonist. The total number of counts were measured for 2 h after agonist treatment.

2.12. Statistical analysis

Computer programmed log-probit methods were used to generate ED₅₀ values for rotation data. Regression analysis of the role of cyclase inhibition vs. inhibition of D₁ binding for antagonism of D₁-dependent rotational behaviour was performed with the SAS REG program (SAS Institute, NC, USA).

3. Results

3.1. In vitro profile and mechanism of action

Similar to SCH 23390, NNC-112, NNC-687 and NNC-756 were selective dopamine D₁ receptor antagonists. Thus, the K_i values for inhibition of [³H]SCH 23390 binding to the D₁ receptor were more than a thousand fold lower than the K_i value for displacement of [³H]spiroperidol from the D₂ receptor (see table 2). Other compounds with neuroleptic activity primarily inhibited D₂ receptor binding or were equally

potent on both D₁ and D₂ receptor subtypes (e.g. clozapine, fluperlapine and cis-flupentixol). Besides inhibiting D₁ receptor binding, NNC-112 and NNC-756 had fairly high affinity for the 5-HT₂ receptor whereas NNC-687 had low affinity for this receptor. The affinity of these novel benzazepine derivatives for other neurotransmitter receptors was marginal (table 3). Similarly, NNC-112, NNC-687 and NNC-756 had no effect on neurotransmitter transporters. However, SCH 23390 inhibited 5-HT uptake with an IC₅₀ of 120 nM (table 4).

The effect of NNC-112, NNC-687 and NNC-756 on the D₁ receptor was investigated in further detail. As observed with SCH 23390, all three compounds inhibited [³H]SCH 23390 binding with Hill slopes close to unity (not shown) (Andersen and Braestrup, 1986). Further, Scatchard analysis indicated a competitive mechanism of action since inclusion of NNC-112, NNC-687 or NNC-756 in the assay affected only the slope of the Scatchard transformation (not shown).

The dopamine D₁ receptor is coupled to adenylyl cyclase in a stimulatory fashion. Inhibition of dopamine-stimulated adenylyl cyclase activity by NNC-112, NNC-687 and NNC-756 occurred in a dose-dependent manner (not shown). Schild transformation indicated a slope close to unity, suggesting that the mechanism behind this inhibition was competitive in nature.

TABLE 5

Effects of NNC-112, NNC-687, NNC-756 and selected reference compounds on D₁, D₂ and 5-HT₂ receptors in vivo.

Results are means from two to three separate experiments with two to six different doses of each compound in each experiment with three mice per dose, ^c means from three mice tested in a single experiment. Drugs were administered ^b s.c. 0.5 h, ^c s.c. 1 h or ^d i.p. 2 h before the animals were killed. > means less than 10% inhibition of the specific binding at the dose indicated. ^a ED₅₀ values were always corrected for changes in the cerebral blood flow. Some of these data have been published in Andersen and Grønvald (1986); Andersen and Nielsen (1986); Andersen et al. (1988).

Receptor/ligand	ED ₅₀ (mg/kg) ^a						
	NNC-112 ^b	NNC-687 ^b	NNC-756 ^b	SCH 23390 ^b	Clozapine ^c	cis-Flupentixol ^d	Haloperidol ^d
<i>Dopamine</i>							
D ₁ [³ H]SCH 23390	0.015	0.35	0.02	0.017	10	0.4	72.5
D ₂ [³ H]spiperone	> 10 ^c	> 10	> 10 ^c	13.5	7.5	0.24	0.06
<i>Serotonin</i>							
5-HT ₂ [³ H]ketanserin	0.9	> 10	0.2	1.5	0.7	0.4	0.72

3.2. *In vivo* profile

The effect of NNC-112, NNC-687 and NNC-756 on the primary *in vitro* targets was analysed *in vivo*. As expected, all three NNC compounds were potent and selective inhibitors of D_1 receptor binding *in vivo*. The effect of NNC-112 and NNC-756 on the 5-HT₂ receptor was seen at substantially higher doses whereas, again as expected, NNC-687 was inactive. NNC-112, NNC-687 and NNC-756 were all inactive in inhibiting [³H]raclopride binding to D_2 receptors (table 5).

The dopamine D_1 receptor selective profile of the NNC compounds was confirmed by the data from the experiments on rotational behaviour in unilaterally 6-OHDA-lesioned rats. Here, the D_1 -mediated rotation induced by SKF 38393 was potently blocked by NNC-112, NNC-687, NNC-756 as well as by SCH 23390 and the non-selective dopamine receptor antagonist cis-flupentixol. On the other hand, haloperidol, a D_2 -selective antagonist, did not affect D_1 receptor-mediated rotation. Rotation induced by administration of the D_2 -selective agonist pergolide was neither affected by NNC-112, NNC-687 or NNC-756 nor by the classical dopamine D_1 receptor antagonist SCH 23390 but was potently antagonized by cis-flupentixol and haloperidol (table 6).

Finally, the role of potency to inhibit cyclase vs. D_1 binding in the ability of the dopamine D_1 receptor antagonists to block D_1 -dependent rotational behaviour was examined by regression analysis. The D_1 binding affinity correlated significantly with the blockade of SKF 38393-induced rotations ($r = 0.75$, $P = 0.036$), but not with the blockade of inhibition of adenylyl cyclase ($r = 0.03$, $P < 0.15$).

4. Discussion

Although neuroleptic drugs share the ability to antagonize brain dopamine receptors, they also bind to many other neurotransmitter receptors (table 3 and

TABLE 6

Effect of NNC-112, NNC-687, NNC-756 and selected reference compounds on D_1 or D_2 receptor-mediated rotation.

Rotation induced by D_1 -selective SKF 38393^a (2.5 mg/kg) or D_2 -selective pergolide^b (0.05 mg/kg), both administered s.c.

Drug	ED ₅₀ (mg/kg)	
	D_1 ^a	D_2 ^b
NNC-112 s.c. 0.5 h	0.08	> 2
NNC-687 i.p. 2 h	0.48	> 50
NNC-756 s.c. 0.5 h	0.03	> 1
SCH 23390 s.c. 0.5 h	0.14	> 2
cis-Flupentixol i.p. 2 h	0.28	0.05
Haloperidol i.p. 2 h	> 5	0.1




Fig. 2. Correlation between K_i values obtained in [³H]SCH 23390 binding and adenylyl cyclase assays. For details see the legend of table 2. Numbers on graph refer to the following compounds: 1, NNC-112; 2, NNC-756; 3, NNC-687; 4, clozapine; 5, fluperlapine; 6, SCH 23390; 7, cis-flupentixol; 8, (+)-butaclamol; 9, fluphenazine; 10, chlorpromazine; 11, trans-flupentixol; 12, spiroperidol.

Peroutka and Snyder, 1980). However, NNC-112, NNC-687 and NNC-756 showed a high degree of selectivity for the dopamine D_1 receptor as their affinity constant for this receptor was in the low nanomolar to picomolar range (table 2). In contrast, their affinity for other secondary targets, such as 5-HT₂ and α_2 -adrenoceptors, was 25- to 1000-fold lower. The benzazepine SCH 23390 was a relatively potent 5-HT uptake inhibitor. Although the NNC dopamine D_1 receptor antagonists are structurally related to SCH 23390, they failed to block 5-HT uptake.

NNC-112, NNC-687 and NNC-756 dose dependently inhibited dopamine-stimulated adenylyl cyclase. The Schild analysis suggested a competitive mechanism of inhibition as previously observed with a number of neuroleptics and SCH 23390 (Andersen and Braestrup, 1986). Further, both compounds were relatively more potent in inhibiting dopamine-stimulated adenylyl cyclase activity than [³H]SCH 23390 binding. These characteristics have previously been observed with atypical neuroleptics of the clozapine type (Andersen and Braestrup, 1986). A graphic presentation of the data (from table 2) clearly indicates that, like the clozapine-type compounds the correlation between pK_i values obtained from [³H]SCH 23390 binding and inhibition of dopamine-stimulated adenylyl cyclase for NNC-112, NNC-687 and NNC-756 was shifted to the left (fig. 2).

The selectivity for the dopamine D_1 receptor was further corroborated by the results of *in vivo* binding experiments. Thus, NNC-112, NNC-687 and NNC-756 inhibited [³H]SCH 23390 binding to the D_1 receptor in

vivo with ED_{50} values in the $\mu\text{g/kg}$ range whereas all three compounds were inactive in inhibiting [^3H]raclopride binding to the D_2 receptor. As observed in vitro, NNC-687 was inactive whereas both NNC-112 and NNC-756 inhibited [^3H]ketanserin binding to the 5-HT $_2$ receptor. However, the potencies were 60 and 10 times lower, respectively, than for the D_1 receptor.

The D_1 selectivity of the NNC benzazepines was confirmed in the behavioural experiments. Thus, in unilaterally 6-OHDA-lesioned rats, NNC-112, NNC-687, NNC-756 and SCH 23390 antagonized only D_1 receptor-induced rotational behaviour. The compounds were without effect on D_2 receptor-mediated rotation, which was blocked by the D_2 -selective dopamine receptor antagonist haloperidol. The non-selective dopamine D_1/D_2 receptor antagonist cis-flupentixol antagonized both the D_1 - and D_2 -mediated rotation.

Interestingly, the inhibitory potency of the dopamine D_1 receptor antagonists at the D_1 receptor binding site, but not on the inhibition of cyclase correlated with their antagonistic potency in blocking D_1 -dependent rotations. The significance of the differential ability of some compounds (e.g. clozapine) to block the dopamine-sensitive adenylate cyclase is presently not known. However, enhanced dopamine-dependent cyclase sensitivity is observed in post-mortem striatal tissue from schizophrenics, suggesting that inhibition of cyclase may be important in the antipsychotic effect of clozapine (e.g. Memo et al., 1983).

Except for clozapine, antipsychotic drugs induce extrapyramidal side-effects and increase plasma prolactin levels (e.g. Andersen and Nielsen, 1991). Can these side-effects be expected with NNC-112, NNC-687 and NNC-756 which, unlike classical neuroleptics, are highly selective dopamine D_1 receptor antagonists? First, conflicting data have been reported concerning the ability of SCH 23390 to induce catalepsy in rodents (a putative animal counterpart of human extrapyramidal side-effects) (Iorio et al., 1983; Christensen et al., 1984; Morelli and DiChiara, 1985; Barnett, 1986). In our laboratory, dopamine D_1 receptor antagonists readily induce a cataleptic syndrome which is indistinguishable from that induced by either D_2 selective or non-selective dopamine D_1 , D_2 receptor antagonists (see Nielsen et al., 1992). However, the propensity of dopamine D_1 receptor antagonists to induce extrapyramidal side-effects in patients may be lower than that observed with dopamine D_2 receptor antagonists. Thus, recent studies indicate that, in contrast to dopamine D_2 receptor antagonists, dopamine D_1 receptor antagonists are incapable of inducing classical extrapyramidal side-effects in *drug-naïve* primates (Coffin et al., 1989). This has been confirmed in primates with NNC-756 in doses up to 1 mg/kg s.c. (Hansen and Gerlach, 1991). It should be noted that the D_1 antagonist SCH 23390 readily induces extrapyramidal side-effects in primates

that have *previously* received D_2 haloperidol (Kistrup and Gerlach, 1987). However, i.m. injected (but not p.o. injected) SCH 23390 was found by Lawrence et al. (1991) to produce signs of 'parkinsonism' in drug-naïve primates (mainly increased sedation, decreased blink rate and, at high doses, tremor). Thus, the route of administration and species may play a role in the expression of motor side-effects by SCH 23390. Further, there may be a difference in the symptomatology of 'Parkinsonism' (e.g. as defined by Lawrence et al.) and extrapyramidal side-effects (as defined by Coffin et al., 1989). Second, since the dopamine D_1 receptor antagonists are devoid of D_2 affinity, they should be without effect in tissues containing only this receptor subtype, e.g. the mammothrophic cells of the pituitary. Consistent with this, dopamine D_1 receptor antagonists have little effect on plasma prolactin levels in animal experiments (Iorio et al., 1983; Rovescalli et al., 1987).

NNC-112, NNC-687 and NNC-756 are new dopamine D_1 receptor-selective antagonists with a neurochemical profile suggestive of antipsychotic potential. Since D_1 -selective antagonists may have a lower propensity to induce extrapyramidal symptoms, such compounds hold great clinical promise.

Acknowledgements

The expert technical assistance of Ms. A. Hansen, H.F. Beck, D.P.D. Andersen and typing assistance of Ms. H.F. Estrup is greatly appreciated. Further, the generous gift of drugs from various drug companies is acknowledged.

References

- Andersen, P.H., 1988, Comparison of the pharmacological characteristics of [^3H]raclopride and [^3H]SCH 23390 binding to dopamine receptors in vivo in mouse brain, *Eur. J. Pharmacol.* 146, 113.
- Andersen, P.H., 1989, The dopamine uptake inhibitor GBR 12909: selectivity and molecular mechanism of action, *Eur. J. Pharmacol.* 166, 493.
- Andersen, P.H. and C. Braestrup, 1986, Evidence for different states of the dopamine D_1 receptor: Clozapine and fluperlapine may preferentially label an adenylate cyclase-coupled state of the D_1 receptor, *J. Neurochem.* 47, 1822.
- Andersen, P.H. and F.C. Grønvald, 1986, Specific binding of [^3H]SCH 23390 to dopamine D_1 receptors in vivo, *Life Sci.* 38, 1507.
- Andersen, P.H. and E.B. Nielsen, 1986, The dopamine D_1 receptor, biochemical and behavioral aspects, in: *Neurobiology of Central D_1 Dopamine Receptors*, Vol. 204, eds. G.B. Breese and I. Creese (Plenum Press, New York) p. 73.
- Andersen, P.H. and E.B. Nielsen, 1991, Novel approaches to development of antipsychotics, *Drug News Perspect.* 4, 150.
- Andersen, P.H., F.C. Grønvald and J.A. Jansen, 1985, A comparison between dopamine-stimulated adenylate cyclase and [^3H]SCH 23390 binding in rat striatum, *Life Sci.* 37, 1971.
- Andersen, P.H., E.B. Nielsen, F.C. Grønvald and C. Braestrup, 1986, Some atypical neuroleptics inhibit [^3H]SCH 23390 binding in vivo, *Eur. J. Pharmacol.* 120, 143.

- Andersen, P.H., F.C. Grønvald, R. Hohlweg, L.B. Hansen, E. Guddal and C. Braestrup, 1988, NO-112, NO-756 – New dopamine D1 selective antagonists, Soc. Neuroscience, Toronto, Canada, Nov. 13–18, Abstract No. 378.13.
- Andersen, P.H., J.A. Gringrich, M.D. Bates, A. Dearry, P. Falardeau, S.E. Senogles and M.G. Caron, 1990, Dopamine receptor subtypes: beyond the D1/D2 classification, Trends Pharmacol. Sci. 11, 231.
- Barnett, A., 1986, Review on dopamine receptors, Drugs Future 11, 49.
- Billard, W., V. Ruperto, G. Crosby, L.C. Iorio and A. Barnett, 1984, Characterization of the binding of [³H]SCH 23390, a selective D-1 receptor antagonist ligand, in rat striatum, Life Sci. 35, 1885.
- Christensen, A.V., J. Arnt, J. Hyttel, J.J. Larsen and O. Svendsen, 1984, Pharmacological effects of a specific dopamine D1 antagonist SCH 23390 in comparison with neuroleptics, Life Sci. 34, 1529.
- Coffin, V.L., M.B. Latranyi and R.E. Chipkin, 1989, Acute extrapyramidal syndrome in Cebus monkeys: development mediated by dopamine D2 but not D1 receptors, J. Pharmacol. Exp. Ther. 249, 1.
- Farde, L., F.A. Wiesel, A.-L. Nordström and G. Sedvall, 1989, D1- and D2-dopamine receptor occupancy during treatment with conventional and atypical neuroleptics, Psychopharmacology 99, S28.
- Geisler, A., R. Klysner, P. Thams and S. Christensen, 1977, A simple and inexpensive protein binding assay for cyclic AMP in biological materials, Acta Pharmacol. Toxicol. 40, 356.
- Hansen, L. and J. Gerlach, 1991, Dopamine 1 and 2 receptor interaction in the development of dystonia and hypersensitivity in Cebus monkeys, Biol. Psychiat. 29, 275S.
- Hyttel, J., 1983, SCH 23390 – the first selective dopamine D1 antagonist, Eur. J. Pharmacol. 91, 153.
- Iorio, L.C., A. Barnett, F.H. Leitz, V.P. Houser and C.A. Korduba, 1983, SCH 23390, a potential benzazepine antipsychotic with unique interactions on dopaminergic systems, J. Pharmacol. Exp. Ther. 226, 462.
- Kistrup, K. and J. Gerlach, 1987, Selective D1 and D2 receptor manipulation in Cebus monkeys: relevance for dystonia and dyskinesia in humans, Pharmacol. Toxicol. 61, 157.
- Lawrence, M.S., D.E. Redmond, Jr., J.D. Elsworth, J.R. Taylor and R.H. Roth, 1991, The D1 receptor antagonist, SCH 23390, induces signs of parkinsonism in African green monkeys, Life Sci. 49, PL229.
- Memo, M., J.E. Kleinman and I. Hanbauer, 1983, Coupling of dopamine D-1 recognition sites with adenylate cyclase in nuclei accumbens and caudatus of schizophrenics, Science 221, 1304.
- Monsma, F.J., Jr., L.C. Mahan, L.D. McVittie, C.R. Gerfen and D.R. Sibley, 1990, Molecular cloning and expression of a D1 dopamine receptor linked to adenylyl cyclase activation, Proc. Natl. Acad. Sci. U.S.A. 87, 6723.
- Morelli, M. and G. DiChiara, 1985, Catalepsy induced by SCH 23390 in rats, Eur. J. Pharmacol. 117, 179.
- Nielsen, E.B., L.B. Hansen, F.C. Grønvald, R. Hohlweg, E. Guddal and P.H. Andersen, 1988, NO-112 and NO-756, new potent benzazepine D-1 antagonists: pharmacological characterization, Soc. Neuroscience, Toronto, Canada, Nov. 13–18, Abstract No. 378.14.
- Nielsen, E.B. and P.H. Andersen, 1992, Dopamine receptor occupancy in vivo: behavioral correlates using NNC-112, NNC-687 and NNC-756, new selective dopamine D-1 receptor antagonists, Eur. J. Pharmacol. 219, 35.
- Peroutka, S.J. and S.H. Snyder, 1980, Relationship of neuroleptic drug effects at brain dopamine, serotonin, alpha-adrenergic, and histamine receptors to clinical potency, Am. J. Psychiat. 137, 1518.
- Rovescalli, A.C., N. Brunello, A. Monopoli, E. Ongini and G. Racagni, 1987, Absence of [³H]SCH 23390 specific binding sites in anterior pituitary. Dissociation from effects on prolactin synthesis, Eur. J. Pharmacol. 135, 129.
- Seeman, P., C. Ulpian, D. Grigoriadis, I. Pri-Bar and O. Buchman, 1985, Conversion of dopamine D1 receptors from high to low affinity for dopamine, Biochem. Pharmacol. 34, 151.
- Van Tol, H.H.M., J.R. Bunzow, H.-C. Guan, R.K. Sunahara, P. Seeman, H.B. Niznik and O. Civelli, 1991, Cloning of the gene for a human dopamine D₄ receptor with high affinity for the antipsychotic clozapine, Nature 350, 610.