

Planarity

October 29, 2024

Motivation :

- 1** Planar graphs are a major link between graph theory and geometry/-topology.
- 2 There are three easily identifiable milestones in planar graph theory.
- 3 A formula of Euler that $V - E + F = 2$ for any convex polyhedron with V vertices/corners, E edges and F faces.
- 4 A deep characterization of planar graphs due to Kuratowski.
- 5 The 4-color-theorem of Appel, Haken and Koch.

Motivation :

- 1** Planar graphs are a major link between graph theory and geometry/-topology.
- 2** There are three easily identifiable milestones in planar graph theory.
- 3 A formula of Euler that $V - E + F = 2$ for any convex polyhedron with V vertices/corners, E edges and F faces.
- 4 A deep characterization of planar graphs due to Kuratowski.
- 5 The 4-color-theorem of Appel, Haken and Koch.

Motivation :

- 1 Planar graphs are a major link between graph theory and geometry/-topology.
- 2 There are three easily identifiable milestones in planar graph theory.
- 3 A formula of Euler that $V - E + F = 2$ for any convex polyhedron with V vertices/corners, E edges and F faces.
- 4 A deep characterization of planar graphs due to Kuratowski.
- 5 The 4-color-theorem of Appel, Haken and Koch.

Motivation :

- 1 Planar graphs are a major link between graph theory and geometry/-topology.
- 2 There are three easily identifiable milestones in planar graph theory.
- 3 A formula of Euler that $V - E + F = 2$ for any convex polyhedron with V vertices/corners, E edges and F faces.
- 4 A deep characterization of planar graphs due to Kuratowski.
- 5 The 4-color-theorem of Appel, Haken and Koch.

Motivation :

- 1 Planar graphs are a major link between graph theory and geometry/-topology.
- 2 There are three easily identifiable milestones in planar graph theory.
- 3 A formula of Euler that $V - E + F = 2$ for any convex polyhedron with V vertices/corners, E edges and F faces.
- 4 A deep characterization of planar graphs due to Kuratowski.
- 5 The 4-color-theorem of Appel, Haken and Koch.

Motivation :

- 1 Colorings of planar graphs made their first appearance in a problem of map coloring.
- 2 Recent applications of planar graphs in the design of chips and VLSI have further boosted the current research on planar graphs.

Motivation :

- 1 Colorings of planar graphs made their first appearance in a problem of map coloring.
- 2 Recent applications of planar graphs in the design of chips and VLSI have further boosted the current research on planar graphs.

Definition

- 1 A graph G is said to be **planar** or embeddable in the plane if it can be drawn in the plane so that no two edges intersect except (possibly) at their end vertices; otherwise it is said to be a **nonplanar** graph.
- 2 A planar graph embedded in the plane is called a **plane graph**.

Definition

- 1 A graph G is said to be **planar** or embeddable in the plane if it can be drawn in the plane so that no two edges intersect except (possibly) at their end vertices; otherwise it is said to be a **nonplanar** graph.
- 2 A planar graph embedded in the plane is called a **plane graph**.




Figure 10.1: Planar graph, plane graphs and nonplanar graphs.

- 1** Find necessary and sufficient conditions for a graph to be planar.
- 2 How to test a given graph for planarity?
- 3 Design a (polynomial time) algorithm to draw a given planar graph as a plane graph.

- 1 Find necessary and sufficient conditions for a graph to be planar.
- 2 How to test a given graph for planarity?
- 3 Design a (polynomial time) algorithm to draw a given planar graph as a plane graph.

- 1 Find necessary and sufficient conditions for a graph to be planar.
- 2 How to test a given graph for planarity?
- 3 Design a (polynomial time) algorithm to draw a given planar graph as a plane graph.

- 1 The first problem was solved by Kuratowski in 1930.
- 2 His characterization uses the hereditary nature of planar graphs.
- 3 A graph theoretic property P is said to be hereditary if a graph has property P then all its subgraphs too have property P .
- 4 Clearly, acyclicity, bipartiteness and planarity are hereditary properties.
- 5 Kuratowski's characterization has lead to the design of many "good" (= polynomial time) algorithms to check whether a given graph is planar, and if it is planar to draw it as a plane graph.

- 1 The first problem was solved by Kuratowski in 1930.
- 2 His characterization uses the hereditary nature of planar graphs.
- 3 A graph theoretic property P is said to be hereditary if a graph has property P then all its subgraphs too have property P .
- 4 Clearly, acyclicity, bipartiteness and planarity are hereditary properties.
- 5 Kuratowski's characterization has lead to the design of many "good" (= polynomial time) algorithms to check whether a given graph is planar, and if it is planar to draw it as a plane graph.

- 1 The first problem was solved by Kuratowski in 1930.
- 2 His characterization uses the hereditary nature of planar graphs.
- 3 A graph theoretic property P is said to be hereditary if a graph has property P then all its subgraphs too have property P .
- 4 Clearly, acyclicity, bipartiteness and planarity are hereditary properties.
- 5 Kuratowski's characterization has lead to the design of many "good" (= polynomial time) algorithms to check whether a given graph is planar, and if it is planar to draw it as a plane graph.

Definition

- 1** Given any two points a and b in the plane, any non-self-intersecting continuous curve from a to b is called a Jordan curve and it is denoted by $J[a, b]$.
- 2** If $a = b$, then J is called a closed Jordan curve.

Definition

- 1 Given any two points a and b in the plane, any non-self-intersecting continuous curve from a to b is called a Jordan curve and it is denoted by $J[a, b]$.
- 2 If $a = b$, then J is called a closed Jordan curve.

Theorem

- 1** Any closed Jordan curve J partitions the plane into 3 parts namely, interior of J ($\text{int } J$), exterior of J ($\text{ext } J$) and J .
- 2** If J is a closed Jordan curve, $s \in \text{int } J$ and $t \in \text{ext } J$, then any Jordan curve $J'[s, t]$ contains a point of J (that is, J' intersects J).

Theorem

- 1** Any closed Jordan curve J partitions the plane into 3 parts namely, interior of J ($\text{int } J$), exterior of J ($\text{ext } J$) and J .
- 2** If J is a closed Jordan curve, $s \in \text{int } J$ and $t \in \text{ext } J$, then any Jordan curve $J'[s, t]$ contains a point of J (that is, J' intersects J).

- 1** If G is a plane graph, then any path in G is identified with a Jordan curve.
- 2 Similarly, any cycle is identified with a closed Jordan curve.
- 3 In particular, an edge $e(u, v)$ of G is a Jordan curve from u to v .

- 1** If G is a plane graph, then any path in G is identified with a Jordan curve.
- 2** Similarly, any cycle is identified with a closed Jordan curve.
- 3** In particular, an edge $e(u, v)$ of G is a Jordan curve from u to v .

- 1 If G is a plane graph, then any path in G is identified with a Jordan curve.
- 2 Similarly, any cycle is identified with a closed Jordan curve.
- 3 In particular, an edge $e(u, v)$ of G is a Jordan curve from u to v .

Definition

Let G be a plane graph.

- 1** G partitions the plane into several regions. These regions are called the **faces** of G . The set of all faces of G is denoted by $F(G)$.
- 2** Except one face, every other face is a bounded region. The exceptional face is called the **exterior face** and other faces are called **interior faces** of G .
- 3** The exterior face is unbounded and interior faces are bounded ($:=$ area is finite).

Definition

Let G be a plane graph.

- 1** G partitions the plane into several regions. These regions are called the **faces** of G . The set of all faces of G is denoted by $F(G)$.
- 2** Except one face, every other face is a bounded region. The exceptional face is called the **exterior face** and other faces are called **interior faces** of G .
- 3** The exterior face is unbounded and interior faces are bounded ($:=$ area is finite).

Definition

Let G be a plane graph.

- 1** G partitions the plane into several regions. These regions are called the **faces** of G . The set of all faces of G is denoted by $F(G)$.
- 2** Except one face, every other face is a bounded region. The exceptional face is called the **exterior face** and other faces are called **interior faces** of G .
- 3** The exterior face is unbounded and interior faces are bounded ($:=$ area is finite).

Faces, Boundary and Degree

Definition

Let G be a plane graph.

- 1 The **boundary of a face** f is the set of all edges of G which are incident with f . It is denoted by $b_G(f)$ or $b(f)$.
- 2 The **degree of a face** f in a plane graph G is the number of edges in the boundary of f with cut-edges counted twice.
- 3 The degree of f is denoted by $\deg_G(f)$ or $\deg(f)$ or $d(f)$.

Definition

Let G be a plane graph.


- 1** The **boundary of a face** f is the set of all edges of G which are incident with f . It is denoted by $b_G(f)$ or $b(f)$.
- 2** The **degree of a face** f in a plane graph G is the number of edges in the boundary of f with cut-edges counted twice.
- 3** The degree of f is denoted by $\deg_G(f)$ or $\deg(f)$ or $d(f)$.

Definition

Let G be a plane graph.

- 1** The **boundary of a face** f is the set of all edges of G which are incident with f . It is denoted by $b_G(f)$ or $b(f)$.
- 2** The **degree of a face** f in a plane graph G is the number of edges in the boundary of f with cut-edges counted twice.
- 3** The degree of f is denoted by $\deg_G(f)$ or $\deg(f)$ or $d(f)$.

Example



f_1, f_2, f_3, f_4 - Faces of G

$b(f_1) = \{12, 23, 13\}$, $d(f_1) = 3$, $b(f_1)$ is a cycle.

$b(f_2) = \{12, 23, 13, 34, 45, 55, 56, 67, 74\}$, $d(f_2) = 10$.

Note that 34 is counted twice and that $b(f_2)$ does not form a cycle.

Consequences of Jordan Curve Theorem

- 1 A cyclic edge belongs to two faces.
- 2 A cut-edge belongs to only one face.
- 3 A plane graph G is acyclic if and only if $|F(G)| = 1$

Consequences of Jordan Curve Theorem

- 1 A cyclic edge belongs to two faces.
- 2 A cut-edge belongs to only one face.
- 3 A plane graph G is acyclic if and only if $|F(G)| = 1$

Consequences of Jordan Curve Theorem

- 1 A cyclic edge belongs to two faces.
- 2 A cut-edge belongs to only one face.
- 3 A plane graph G is acyclic if and only if $|F(G)| = 1$

Euler's Formula

Theorem

For a connected plane graph G ,

$$V - E + F = 2$$

Proof.

We discuss two possible cases:

- 1 G is acyclic.
- 2 G is cyclic.



Euler's Formula

Theorem

For a connected plane graph G ,

$$V - E + F = 2$$

Proof.

We discuss two possible cases:

- 1 G is acyclic.
- 2 G is cyclic.



Euler's Formula

Theorem

For a connected plane graph G ,

$$V - E + F = 2$$

Proof.

We discuss two possible cases:

1 G is acyclic.

2 G is cyclic.



Euler's Formula

Theorem

For a connected plane graph G ,

$$V - E + F = 2$$

Proof.

We discuss two possible cases:

- 1 G is acyclic.
- 2 G is cyclic.



Proof.

Case 1 : G is acyclic.

- 1 G is connected, acyclic and has n vertices.
- 2 G is a tree and has $n - 1$ edges.
- 3 G has only one face (unbounded one).
- 4 $V - E + F = n - (n - 1) + 1 = 2$.



Proof.

Case 1 : G is acyclic.

1 G is connected, acyclic and has n vertices.

2 G is a tree and has $n - 1$ edges.

3 G has only one face (unbounded one).

4 $V - E + F = n - (n - 1) + 1 = 2.$



Proof.

Case 1 : G is acyclic.

1 G is connected, acyclic and has n vertices.

2 G is a tree and has $n - 1$ edges.

3 G has only one face (unbounded one).

4 $V - E + F = n - (n - 1) + 1 = 2.$



Proof.

Case 1 : G is acyclic.

1 G is connected, acyclic and has n vertices.

2 G is a tree and has $n - 1$ edges.

3 G has only one face (unbounded one).

4 $V - E + F = n - (n - 1) + 1 = 2.$



Proof.

Case 1 : G is acyclic.

- 1** G is connected, acyclic and has n vertices.
- 2** G is a tree and has $n - 1$ edges.
- 3** G has only one face (unbounded one).
- 4** $V - E + F = n - (n - 1) + 1 = 2.$



Proof.

Case 2 : G is cyclic.

- 1 Proof by induction on the number of edges in the graph.
- 2 **Base Case :** $E = 3$ (K_3).
- 3 Then, $v = 3$ and $F = 2$.
- 4 Hence, $V - E + F = 2$.
- 5 **Induction Hypothesis :** Assume true for all connected graphs with m edges.
- 6 Now consider a graph G with $m + 1$ edges.



Proof.

Case 2 : G is cyclic.

- 1 Proof by induction on the number of edges in the graph.
- 2 **Base Case :** $E = 3$ (K_3).
- 3 Then, $v = 3$ and $F = 2$.
- 4 Hence, $V - E + F = 2$.
- 5 **Induction Hypothesis :** Assume true for all connected graphs with m edges.
- 6 Now consider a graph G with $m + 1$ edges.



Proof.

Case 2 : G is cyclic.

- 1 Proof by induction on the number of edges in the graph.
- 2 **Base Case :** $E = 3$ (K_3).
- 3 Then, $v = 3$ and $F = 2$.
- 4 Hence, $V - E + F = 2$.
- 5 **Induction Hypothesis :** Assume true for all connected graphs with m edges.
- 6 Now consider a graph G with $m + 1$ edges.



Proof.

Case 2 : G is cyclic.

- 1 Proof by induction on the number of edges in the graph.
- 2 **Base Case :** $E = 3$ (K_3).
- 3 Then, $v = 3$ and $F = 2$.
- 4 Hence, $V - E + F = 2$.
- 5 **Induction Hypothesis :** Assume true for all connected graphs with m edges.
- 6 Now consider a graph G with $m + 1$ edges.



Proof.

Case 2 : G is cyclic.

- 1 Proof by induction on the number of edges in the graph.
- 2 **Base Case :** $E = 3$ (K_3).
- 3 Then, $v = 3$ and $F = 2$.
- 4 Hence, $V - E + F = 2$.
- 5 **Induction Hypothesis :** Assume true for all connected graphs with m edges.
- 6 Now consider a graph G with $m + 1$ edges.



Proof.

Case 2 : G is cyclic.

- 1 Proof by induction on the number of edges in the graph.
- 2 **Base Case :** $E = 3$ (K_3).
- 3 Then, $v = 3$ and $F = 2$.
- 4 Hence, $V - E + F = 2$.
- 5 **Induction Hypothesis :** Assume true for all connected graphs with m edges.
- 6 Now consider a graph G with $m + 1$ edges.



Proof.

Case 2 : G is cyclic.

- 1 Proof by induction on the number of edges in the graph.
- 2 **Base Case :** $E = 3$ (K_3).
- 3 Then, $v = 3$ and $F = 2$.
- 4 Hence, $V - E + F = 2$.
- 5 **Induction Hypothesis :** Assume true for all connected graphs with m edges.
- 6 Now consider a graph G with $m + 1$ edges.



Proof.

Case 2 : G is cyclic.

- 1 Let G be a connected cyclic graph with $m + 1$ edges.
- 2 Let C be a cycle of G .
- 3 Let $e \in E(C)$.
- 4 Then, $G - e$ is connected.
- 5 $V(G - e) = V(G)$, $E(G - e) = E - 1$ and $F(G - e) = F - 1$.
- 6 Note that $V(G - e) - E(G - e) + F(G - e) = 2$ (By induction if $G - e$ is cyclic, and by Case 1 if $G - e$ is a tree).
- 7 $V(G) - E(G) + F(G) =$
 $V(G - e) - E(G - e) - 1 + F(G - e) + 1 = 2.$

Proof.

Case 2 : G is cyclic.

- 1 Let G be a connected cyclic graph with $m + 1$ edges.
- 2 Let C be a cycle of G .
- 3 Let $e \in E(C)$.
- 4 Then, $G - e$ is connected.
- 5 $V(G - e) = V(G)$, $E(G - e) = E - 1$ and $F(G - e) = F - 1$.
- 6 Note that $V(G - e) - E(G - e) + F(G - e) = 2$ (By induction if $G - e$ is cyclic, and by Case 1 if $G - e$ is a tree).
- 7 $V(G) - E(G) + F(G) =$
 $V(G - e) - E(G - e) - 1 + F(G - e) + 1 = 2.$

Proof.

Case 2 : G is cyclic.

- 1 Let G be a connected cyclic graph with $m + 1$ edges.
- 2 Let C be a cycle of G .
- 3 Let $e \in E(C)$.
- 4 Then, $G - e$ is connected.
- 5 $V(G - e) = V(G)$, $E(G - e) = E - 1$ and $F(G - e) = F - 1$.
- 6 Note that $V(G - e) - E(G - e) + F(G - e) = 2$ (By induction if $G - e$ is cyclic, and by Case 1 if $G - e$ is a tree).
- 7
$$V(G) - E(G) + F(G) = V(G - e) - E(G - e) - 1 + F(G - e) + 1 = 2.$$

Proof.

Case 2 : G is cyclic.

- 1 Let G be a connected cyclic graph with $m + 1$ edges.
- 2 Let C be a cycle of G .
- 3 Let $e \in E(C)$.
- 4 Then, $G - e$ is connected.
- 5 $V(G - e) = V(G)$, $E(G - e) = E - 1$ and $F(G - e) = F - 1$.
- 6 Note that $V(G - e) - E(G - e) + F(G - e) = 2$ (By induction if $G - e$ is cyclic, and by Case 1 if $G - e$ is a tree).
- 7 $V(G) - E(G) + F(G) =$
 $V(G - e) - E(G - e) - 1 + F(G - e) + 1 = 2.$

Proof.

Case 2 : G is cyclic.

- 1 Let G be a connected cyclic graph with $m + 1$ edges.
- 2 Let C be a cycle of G .
- 3 Let $e \in E(C)$.
- 4 Then, $G - e$ is connected.
- 5 $V(G - e) = V(G)$, $E(G - e) = E - 1$ and $F(G - e) = F - 1$.
- 6 Note that $V(G - e) - E(G - e) + F(G - e) = 2$ (By induction if $G - e$ is cyclic, and by Case 1 if $G - e$ is a tree).
- 7 $V(G) - E(G) + F(G) =$
 $V(G - e) - E(G - e) - 1 + F(G - e) + 1 = 2.$

Proof.

Case 2 : G is cyclic.

- 1 Let G be a connected cyclic graph with $m + 1$ edges.
- 2 Let C be a cycle of G .
- 3 Let $e \in E(C)$.
- 4 Then, $G - e$ is connected.
- 5 $V(G - e) = V(G)$, $E(G - e) = E - 1$ and $F(G - e) = F - 1$.
- 6 Note that $V(G - e) - E(G - e) + F(G - e) = 2$ (By induction if $G - e$ is cyclic, and by Case 1 if $G - e$ is a tree).
- 7 $V(G) - E(G) + F(G) =$
 $V(G - e) - E(G - e) - 1 + F(G - e) + 1 = 2.$

Proof.

Case 2 : G is cyclic.

- 1 Let G be a connected cyclic graph with $m + 1$ edges.
- 2 Let C be a cycle of G .
- 3 Let $e \in E(C)$.
- 4 Then, $G - e$ is connected.
- 5 $V(G - e) = V(G)$, $E(G - e) = E - 1$ and $F(G - e) = F - 1$.
- 6 Note that $V(G - e) - E(G - e) + F(G - e) = 2$ (By induction if $G - e$ is cyclic, and by Case 1 if $G - e$ is a tree).
- 7 $V(G) - E(G) + F(G) =$
 $V(G - e) - E(G - e) - 1 + F(G - e) + 1 = 2.$

Proof.

Case 2 : G is cyclic.

- 1 Let G be a connected cyclic graph with $m + 1$ edges.
- 2 Let C be a cycle of G .
- 3 Let $e \in E(C)$.
- 4 Then, $G - e$ is connected.
- 5 $V(G - e) = V(G)$, $E(G - e) = E - 1$ and $F(G - e) = F - 1$.
- 6 Note that $V(G - e) - E(G - e) + F(G - e) = 2$ (By induction if $G - e$ is cyclic, and by Case 1 if $G - e$ is a tree).
- 7 $V(G) - E(G) + F(G) =$
 $V(G - e) - E(G - e) - 1 + F(G - e) + 1 = 2.$



Consequences of Euler's formula

Theorem

If G is a simple connected planar graph with $|V| \geq 3$, then $\frac{3}{2}F \leq E \leq 3V - 6$.

Proof.

We split into two cases.

- 1 G has no bounded face.
- 2 G has a bounded face.



Consequences of Euler's formula

Theorem

If G is a simple connected planar graph with $|V| \geq 3$, then
$$\frac{3}{2}F \leq E \leq 3V - 6.$$

Proof.

We split into two cases.

- 1 G has no bounded face.
- 2 G has a bounded face.



Consequences of Euler's formula

Theorem

If G is a simple connected planar graph with $|V| \geq 3$, then $\frac{3}{2}F \leq E \leq 3V - 6$.

Proof.

We split into two cases.

- 1** G has no bounded face.
- 2** G has a bounded face.



Consequences of Euler's formula

Theorem

If G is a simple connected planar graph with $|V| \geq 3$, then $\frac{3}{2}F \leq E \leq 3V - 6$.

Proof.

We split into two cases.

- 1** G has no bounded face.
- 2** G has a bounded face.



Proof Contd. Case 1

Proof.

Case 1 : G has no bounded face.

- 1 Since G is connected, G is a tree.
- 2 Then, $|E| = |V| - 1$
- 3 Since $|V| \geq 3$, $|E| = |V| - 1 \leq 2|V| - 4 \leq 3|V| - 6$.
- 4 Also, $|F| = 1$, $|E| \geq 2$.
- 5 Hence, $\frac{3}{2}F \leq E \leq 3V - 6$.



Proof.

Case 1 : G has no bounded face.

1 Since G is connected, G is a tree.

2 Then, $|E| = |V| - 1$

3 Since $|V| \geq 3$, $|E| = |V| - 1 \leq 2|V| - 4 \leq 3|V| - 6$.

4 Also, $|F| = 1$, $|E| \geq 2$.

5 Hence, $\frac{3}{2}F \leq E \leq 3V - 6$.



Proof Contd. Case 1

Proof.

Case 1 : G has no bounded face.

1 Since G is connected, G is a tree.

2 Then, $|E| = |V| - 1$

3 Since $|V| \geq 3$, $|E| = |V| - 1 \leq 2|V| - 4 \leq 3|V| - 6$.

4 Also, $|F| = 1$, $|E| \geq 2$.

5 Hence, $\frac{3}{2}F \leq E \leq 3V - 6$.



Proof Contd. Case 1

Proof.

Case 1 : G has no bounded face.

- 1** Since G is connected, G is a tree.
- 2** Then, $|E| = |V| - 1$
- 3** Since $|V| \geq 3$, $|E| = |V| - 1 \leq 2|V| - 4 \leq 3|V| - 6$.
- 4** Also, $|F| = 1$, $|E| \geq 2$.
- 5** Hence, $\frac{3}{2}F \leq E \leq 3V - 6$.



Proof Contd. Case 1

Proof.

Case 1 : G has no bounded face.

- 1** Since G is connected, G is a tree.
- 2** Then, $|E| = |V| - 1$
- 3** Since $|V| \geq 3$, $|E| = |V| - 1 \leq 2|V| - 4 \leq 3|V| - 6$.
- 4** Also, $|F| = 1$, $|E| \geq 2$.
- 5** Hence, $\frac{3}{2}F \leq E \leq 3V - 6$.



Proof.

Case 1 : G has no bounded face.

- 1** Since G is connected, G is a tree.
- 2** Then, $|E| = |V| - 1$
- 3** Since $|V| \geq 3$, $|E| = |V| - 1 \leq 2|V| - 4 \leq 3|V| - 6$.
- 4** Also, $|F| = 1$, $|E| \geq 2$.
- 5** Hence, $\frac{3}{2}F \leq E \leq 3V - 6$.



Proof Contd. Case 2

Proof.

Case 2 : G has a bounded face.

- 1 G contains a cycle.
- 2 Then, $\deg(f) \geq 3$ (Number of edges in the boundary of a face)
- 3 Each edge is in the boundary of two faces.
- 4 $\sum \deg(f) = 2|E|$
- 5 Combining the above two, we have, Since
 $3|F| \leq 2|E| \Rightarrow \frac{3}{2}F \leq E.$
- 6 By Euler's formula,
 $F = 2 - V + E \Rightarrow 3F = 6 - 3V + 3E \leq 2E.$
- 7 Hence, $\frac{3}{2}F \leq E \leq 3V - 6.$



Proof Contd. Case 2

Proof.

Case 2 : G has a bounded face.

1 G contains a cycle.

2 Then, $\deg(f) \geq 3$ (Number of edges in the boundary of a face)

3 Each edge is in the boundary of two faces.

4 $\sum \deg(f) = 2|E|$

5 Combining the above two, we have, Since
 $3|F| \leq 2|E| \Rightarrow \frac{3}{2}F \leq E.$

6 By Euler's formula,
 $F = 2 - V + E \Rightarrow 3F = 6 - 3V + 3E \leq 2E.$

7 Hence, $\frac{3}{2}F \leq E \leq 3V - 6.$



Proof Contd. Case 2

Proof.

Case 2 : G has a bounded face.

- 1 G contains a cycle.
- 2 Then, $\deg(f) \geq 3$ (Number of edges in the boundary of a face)
- 3 Each edge is in the boundary of two faces.
- 4 $\sum \deg(f) = 2|E|$
- 5 Combining the above two, we have, Since
 $3|F| \leq 2|E| \Rightarrow \frac{3}{2}F \leq E.$
- 6 By Euler's formula,
 $F = 2 - V + E \Rightarrow 3F = 6 - 3V + 3E \leq 2E.$
- 7 Hence, $\frac{3}{2}F \leq E \leq 3V - 6.$



Proof Contd. Case 2

Proof.

Case 2 : G has a bounded face.

- 1** G contains a cycle.
- 2** Then, $\deg(f) \geq 3$ (Number of edges in the boundary of a face)
- 3** Each edge is in the boundary of two faces.
- 4** $\sum \deg(f) = 2|E|$
- 5** Combining the above two, we have, Since
 $3|F| \leq 2|E| \Rightarrow \frac{3}{2}F \leq E.$
- 6** By Euler's formula,
 $F = 2 - V + E \Rightarrow 3F = 6 - 3V + 3E \leq 2E.$
- 7** Hence, $\frac{3}{2}F \leq E \leq 3V - 6.$



Proof Contd. Case 2

Proof.

Case 2 : G has a bounded face.

- 1** G contains a cycle.
- 2** Then, $\deg(f) \geq 3$ (Number of edges in the boundary of a face)
- 3** Each edge is in the boundary of two faces.
- 4** $\sum \deg(f) = 2|E|$
- 5** Combining the above two, we have, Since
 $3|F| \leq 2|E| \Rightarrow \frac{3}{2}F \leq E.$
- 6** By Euler's formula,
 $F = 2 - V + E \Rightarrow 3F = 6 - 3V + 3E \leq 2E.$
- 7** Hence, $\frac{3}{2}F \leq E \leq 3V - 6.$



Proof Contd. Case 2

Proof.

Case 2 : G has a bounded face.

- 1** G contains a cycle.
- 2** Then, $\deg(f) \geq 3$ (Number of edges in the boundary of a face)
- 3** Each edge is in the boundary of two faces.
- 4** $\sum \deg(f) = 2|E|$
- 5** Combining the above two, we have, Since
 $3|F| \leq 2|E| \Rightarrow \frac{3}{2}F \leq E.$
- 6** By Euler's formula,
 $F = 2 - V + E \Rightarrow 3F = 6 - 3V + 3E \leq 2E.$
- 7** Hence, $\frac{3}{2}F \leq E \leq 3V - 6.$



Proof Contd. Case 2

Proof.

Case 2 : G has a bounded face.

- 1** G contains a cycle.
- 2** Then, $\deg(f) \geq 3$ (Number of edges in the boundary of a face)
- 3** Each edge is in the boundary of two faces.
- 4** $\sum \deg(f) = 2|E|$
- 5** Combining the above two, we have, Since
 $3|F| \leq 2|E| \Rightarrow \frac{3}{2}F \leq E.$
- 6** By Euler's formula,
 $F = 2 - V + E \Rightarrow 3F = 6 - 3V + 3E \leq 2E.$
- 7** Hence, $\frac{3}{2}F \leq E \leq 3V - 6.$



Proof Contd. Case 2

Proof.

Case 2 : G has a bounded face.

- 1** G contains a cycle.
- 2** Then, $\deg(f) \geq 3$ (Number of edges in the boundary of a face)
- 3** Each edge is in the boundary of two faces.
- 4** $\sum \deg(f) = 2|E|$
- 5** Combining the above two, we have, Since
 $3|F| \leq 2|E| \Rightarrow \frac{3}{2}F \leq E.$
- 6** By Euler's formula,
 $F = 2 - V + E \Rightarrow 3F = 6 - 3V + 3E \leq 2E.$
- 7** Hence, $\frac{3}{2}F \leq E \leq 3V - 6.$



Corollary 2

Corollary

If G is a connected planar graph, G has a vertex of degree less than six.

Proof.

- 1 This is clearly true if $|V(G)| = 1$ or 2.
- 2 Let $|V| \geq 3$.
- 3 By contradiction, suppose $\deg(v) \geq 6$ for all $v \in V(G)$.
- 4 $6|V| \leq \sum \deg(v) = 2|E| \leq 2(3V - 6) = 6V - 12$.
- 5 A contradiction.
- 6 Hence there is a vertex with degree less than six.



Corollary 2

Corollary

If G is a connected planar graph, G has a vertex of degree less than six.

Proof.

- 1** This is clearly true if $|V(G)| = 1$ or 2.
- 2** Let $|V| \geq 3$.
- 3** By contradiction, suppose $\deg(v) \geq 6$ for all $v \in V(G)$.
- 4** $6|V| \leq \sum \deg(v) = 2|E| \leq 2(3|V| - 6) = 6|V| - 12$.
- 5** A contradiction.
- 6** Hence there is a vertex with degree less than six.



Corollary 2

Corollary

If G is a connected planar graph, G has a vertex of degree less than six.

Proof.

- 1** This is clearly true if $|V(G)| = 1$ or 2.
- 2** Let $|V| \geq 3$.
- 3** By contradiction, suppose $\deg(v) \geq 6$ for all $v \in V(G)$.
- 4** $6|V| \leq \sum \deg(v) = 2|E| \leq 2(3|V| - 6) = 6|V| - 12$.
- 5** A contradiction.
- 6** Hence there is a vertex with degree less than six.



Corollary 2

Corollary

If G is a connected planar graph, G has a vertex of degree less than six.

Proof.

- 1** This is clearly true if $|V(G)| = 1$ or 2 .
- 2** Let $|V| \geq 3$.
- 3** By contradiction, suppose $\deg(v) \geq 6$ for all $v \in V(G)$.
- 4** $6|V| \leq \sum \deg(v) = 2|E| \leq 2(3|V| - 6) = 6|V| - 12$.
- 5** A contradiction.
- 6** Hence there is a vertex with degree less than six.



Corollary 2

Corollary

If G is a connected planar graph, G has a vertex of degree less than six.

Proof.

- 1** This is clearly true if $|V(G)| = 1$ or 2.
- 2** Let $|V| \geq 3$.
- 3** By contradiction, suppose $\deg(v) \geq 6$ for all $v \in V(G)$.
- 4** $6|V| \leq \sum \deg(v) = 2|E| \leq 2(3v - 6) = 6V - 12$.
- 5** A contradiction.
- 6** Hence there is a vertex with degree less than six.



Corollary 2

Corollary

If G is a connected planar graph, G has a vertex of degree less than six.

Proof.

- 1** This is clearly true if $|V(G)| = 1$ or 2 .
- 2** Let $|V| \geq 3$.
- 3** By contradiction, suppose $\deg(v) \geq 6$ for all $v \in V(G)$.
- 4** $6|V| \leq \sum \deg(v) = 2|E| \leq 2(3|V| - 6) = 6|V| - 12$.
- 5** A contradiction.
- 6** Hence there is a vertex with degree less than six.



Corollary 2

Corollary

If G is a connected planar graph, G has a vertex of degree less than six.

Proof.

- 1** This is clearly true if $|V(G)| = 1$ or 2 .
- 2** Let $|V| \geq 3$.
- 3** By contradiction, suppose $\deg(v) \geq 6$ for all $v \in V(G)$.
- 4** $6|V| \leq \sum \deg(v) = 2|E| \leq 2(3V - 6) = 6V - 12$.
- 5** A contradiction.
- 6** Hence there is a vertex with degree less than six.



Corollary 3

Corollary

Let G be a connected planar triangle-free graph with $|V| \geq 3$, then $|E| \leq 2|V| - 4$.

Proof.

- 1 We have already proved the case when G has no bounded face earlier.
- 2 Let G be triangle-free and has at least one bounded face.
- 3 $\Rightarrow \deg(f) \geq 4$.
- 4 $\sum \deg(f) = 2|E|$
- 5 $2|E| = \sum \deg(f) \geq 4|F| \Rightarrow 2|F| \leq |E|$.
- 6 $|V| - |E| + |F| = 2$ (Euler's formula) \Rightarrow
 $2|E| - 2|V| + 4 = 2|F| \leq |E|$.
- 7 Hence, $|E| \leq 2|V| - 4$

Corollary 3

Corollary

Let G be a connected planar triangle-free graph with $|V| \geq 3$, then $|E| \leq 2|V| - 4$.

Proof.

- 1 We have already proved the case when G has no bounded face earlier.
- 2 Let G be triangle-free and has at least one bounded face.
- 3 $\Rightarrow \deg(f) \geq 4$.
- 4 $\sum \deg(f) = 2|E|$
- 5 $2|E| = \sum \deg(f) \geq 4|F| \Rightarrow 2|F| \leq |E|$.
- 6 $|V| - |E| + |F| = 2$ (Euler's formula) \Rightarrow
 $2|E| - 2|V| + 4 = 2|F| \leq |E|$.
- 7 Hence, $|E| \leq 2|V| - 4$

Corollary 3

Corollary

Let G be a connected planar triangle-free graph with $|V| \geq 3$, then $|E| \leq 2|V| - 4$.

Proof.

- 1 We have already proved the case when G has no bounded face earlier.
- 2 Let G be triangle-free and has at least one bounded face.
- 3 $\Rightarrow \deg(f) \geq 4$.
- 4 $\sum \deg(f) = 2|E|$
- 5 $2|E| = \sum \deg(f) \geq 4|F| \Rightarrow 2|F| \leq |E|$.
- 6 $|V| - |E| + |F| = 2$ (Euler's formula) \Rightarrow
 $2|E| - 2|V| + 4 = 2|F| \leq |E|$.
- 7 Hence, $|E| \leq 2|V| - 4$

Corollary 3

Corollary

Let G be a connected planar triangle-free graph with $|V| \geq 3$, then $|E| \leq 2|V| - 4$.

Proof.

- 1 We have already proved the case when G has no bounded face earlier.
- 2 Let G be triangle-free and has at least one bounded face.
- 3 $\Rightarrow \deg(f) \geq 4$.
- 4 $\sum \deg(f) = 2|E|$
- 5 $2|E| = \sum \deg(f) \geq 4|F| \Rightarrow 2|F| \leq |E|$.
- 6 $|V| - |E| + |F| = 2$ (Euler's formula) \Rightarrow
 $2|E| - 2|V| + 4 = 2|F| \leq |E|$.
- 7 Hence, $|E| \leq 2|V| - 4$

Corollary 3

Corollary

Let G be a connected planar triangle-free graph with $|V| \geq 3$, then $|E| \leq 2|V| - 4$.

Proof.

- 1 We have already proved the case when G has no bounded face earlier.
- 2 Let G be triangle-free and has at least one bounded face.
- 3 $\Rightarrow \deg(f) \geq 4$.
- 4 $\sum \deg(f) = 2|E|$
- 5 $2|E| = \sum \deg(f) \geq 4|F| \Rightarrow 2|F| \leq |E|$.
- 6 $|V| - |E| + |F| = 2$ (Euler's formula) \Rightarrow
 $2|E| - 2|V| + 4 = 2|F| \leq |E|$.
- 7 Hence, $|E| \leq 2|V| - 4$

Corollary 3

Corollary

Let G be a connected planar triangle-free graph with $|V| \geq 3$, then $|E| \leq 2|V| - 4$.

Proof.

- 1 We have already proved the case when G has no bounded face earlier.
- 2 Let G be triangle-free and has at least one bounded face.
- 3 $\Rightarrow \deg(f) \geq 4$.
- 4 $\sum \deg(f) = 2|E|$
- 5 $2|E| = \sum \deg(f) \geq 4|F| \Rightarrow 2|F| \leq |E|$.
- 6 $|V| - |E| + |F| = 2$ (Euler's formula) \Rightarrow
 $2|E| - 2|V| + 4 = 2|F| \leq |E|$.
- 7 Hence, $|E| \leq 2|V| - 4$

Corollary 3

Corollary

Let G be a connected planar triangle-free graph with $|V| \geq 3$, then $|E| \leq 2|V| - 4$.

Proof.

- 1 We have already proved the case when G has no bounded face earlier.
- 2 Let G be triangle-free and has at least one bounded face.
- 3 $\Rightarrow \deg(f) \geq 4$.
- 4 $\sum \deg(f) = 2|E|$
- 5 $2|E| = \sum \deg(f) \geq 4|F| \Rightarrow 2|F| \leq |E|$.
- 6 $|V| - |E| + |F| = 2$ (Euler's formula) \Rightarrow
 $2|E| - 2|V| + 4 = 2|F| \leq |E|$.
- 7 Hence, $|E| \leq 2|V| - 4$

Example




Figure: Kuratowski's graph

Kuratowski's Theorem

Theorem

K_5 and $K_{3,3}$ are non-planar.

Proof.

- 1 K_5 has 5 vertices and 10 edges.
- 2 $3|V| - 6 = 15 - 6 = 9 < 10 = |E|$.
- 3 Hence, K_5 is non-planar.
- 4 $K_{3,3}$ has 6 vertices, 9 edges and is triangle-free.
- 5 But, $|E| = 9 \not\leq 2|V| - 4 = 12 - 4 = 8$.
- 6 Hence, $K_{3,3}$ is non-planar.



Kuratowski's Theorem

Theorem

K_5 and $K_{3,3}$ are non-planar.

Proof.

- 1 K_5 has 5 vertices and 10 edges.
- 2 $3|V| - 6 = 15 - 6 = 9 < 10 = |E|$.
- 3 Hence, K_5 is non-planar.
- 4 $K_{3,3}$ has 6 vertices, 9 edges and is triangle-free.
- 5 But, $|E| = 9 \not\leq 2|V| - 4 = 12 - 4 = 8$.
- 6 Hence, $K_{3,3}$ is non-planar.



Kuratowski's Theorem

Theorem

K_5 and $K_{3,3}$ are non-planar.

Proof.

1 K_5 has 5 vertices and 10 edges.

2 $3|V| - 6 = 15 - 6 = 9 < 10 = |E|$.

3 Hence, K_5 is non-planar.

4 $K_{3,3}$ has 6 vertices, 9 edges and is triangle-free.

5 But, $|E| = 9 \not\leq 2|V| - 4 = 12 - 4 = 8$.

6 Hence, $K_{3,3}$ is non-planar.



Kuratowski's Theorem

Theorem

K_5 and $K_{3,3}$ are non-planar.

Proof.

- 1 K_5 has 5 vertices and 10 edges.
- 2 $3|V| - 6 = 15 - 6 = 9 < 10 = |E|$.
- 3 Hence, K_5 is non-planar.
- 4 $K_{3,3}$ has 6 vertices, 9 edges and triangle-free.
- 5 But, $|E| = 9 \not\leq 2|V| - 4 = 12 - 4 = 8$.
- 6 Hence, $K_{3,3}$ is non-planar.



Kuratowski's Theorem

Theorem

K_5 and $K_{3,3}$ are non-planar.

Proof.

- 1 K_5 has 5 vertices and 10 edges.
- 2 $3|V| - 6 = 15 - 6 = 9 < 10 = |E|$.
- 3 Hence, K_5 is non-planar.
- 4 $K_{3,3}$ has 6 vertices, 9 edges and triangle-free.
- 5 But, $|E| = 9 \not\leq 2|V| - 4 = 12 - 4 = 8$.
- 6 Hence, $K_{3,3}$ is non-planar.



Kuratowski's Theorem

Theorem

K_5 and $K_{3,3}$ are non-planar.

Proof.

- 1 K_5 has 5 vertices and 10 edges.
- 2 $3|V| - 6 = 15 - 6 = 9 < 10 = |E|$.
- 3 Hence, K_5 is non-planar.
- 4 $K_{3,3}$ has 6 vertices, 9 edges and is triangle-free.
- 5 But, $|E| = 9 \not\leq 2|V| - 4 = 12 - 4 = 8$.
- 6 Hence, $K_{3,3}$ is non-planar.



Kuratowski's Theorem

Theorem

K_5 and $K_{3,3}$ are non-planar.

Proof.

- 1 K_5 has 5 vertices and 10 edges.
- 2 $3|V| - 6 = 15 - 6 = 9 < 10 = |E|$.
- 3 Hence, K_5 is non-planar.
- 4 $K_{3,3}$ has 6 vertices, 9 edges and is triangle-free.
- 5 But, $|E| = 9 \not\leq 2|V| - 4 = 12 - 4 = 8$.
- 6 Hence, $K_{3,3}$ is non-planar.



- 1 Both are regular graphs.
- 2 Both are nonplanar.
- 3 Removal of one edge or a vertex makes each a planar graph.
- 4 Kuratowski's first graph is the nonplanar graph with the smallest number of vertices.
- 5 Kuratowski's second graph is the nonplanar graph with the smallest number of edges.
- 6 Thus both are the simplest nonplanar graphs.
- 7 The letter K being for Kuratowski.

- 1 Both are regular graphs.
- 2 Both are nonplanar.
- 3 Removal of one edge or a vertex makes each a planar graph.
- 4 Kuratowski's first graph is the nonplanar graph with the smallest number of vertices.
- 5 Kuratowski's second graph is the nonplanar graph with the smallest number of edges.
- 6 Thus both are the simplest nonplanar graphs.
- 7 The letter K being for Kuratowski.

- 1 Both are regular graphs.
- 2 Both are nonplanar.
- 3 Removal of one edge or a vertex makes each a planar graph.
- 4 Kuratowski's first graph is the nonplanar graph with the smallest number of vertices.
- 5 Kuratowski's second graph is the nonplanar graph with the smallest number of edges.
- 6 Thus both are the simplest nonplanar graphs.
- 7 The letter K being for Kuratowski.

- 1 Both are regular graphs.
- 2 Both are nonplanar.
- 3 Removal of one edge or a vertex makes each a planar graph.
- 4 Kuratowski's first graph is the nonplanar graph with the smallest number of vertices.
- 5 Kuratowski's second graph is the nonplanar graph with the smallest number of edges.
- 6 Thus both are the simplest nonplanar graphs.
- 7 The letter K being for Kuratowski.

- 1 Both are regular graphs.
- 2 Both are nonplanar.
- 3 Removal of one edge or a vertex makes each a planar graph.
- 4 Kuratowski's first graph is the nonplanar graph with the smallest number of vertices.
- 5 Kuratowski's second graph is the nonplanar graph with the smallest number of edges.
- 6 Thus both are the simplest nonplanar graphs.
- 7 The letter K being for Kuratowski.


- 1 Both are regular graphs.
- 2 Both are nonplanar.
- 3 Removal of one edge or a vertex makes each a planar graph.
- 4 Kuratowski's first graph is the nonplanar graph with the smallest number of vertices.
- 5 Kuratowski's second graph is the nonplanar graph with the smallest number of edges.
- 6 Thus both are the simplest nonplanar graphs.
- 7 The letter K being for Kuratowski.

- 1 Both are regular graphs.
- 2 Both are nonplanar.
- 3 Removal of one edge or a vertex makes each a planar graph.
- 4 Kuratowski's first graph is the nonplanar graph with the smallest number of vertices.
- 5 Kuratowski's second graph is the nonplanar graph with the smallest number of edges.
- 6 Thus both are the simplest nonplanar graphs.
- 7 The letter K being for Kuratowski.


Subdivision

Definition

- 1** The subdivision of an edge $e(u, v) \in E(G)$ is the operation of replacing e by a path (u, w, v) , where w is a new vertex.
- 2** So, to get a subdivision of e introduce a new vertex w on e .
- 3** A graph H is said to be a subdivision of G if H can be obtained from G by a sequence of edge subdivisions. (By definition, G is a subdivision of G .)




(a) Original Graph G




(b) Subdivision $S(G)$

Definition

- 1** The subdivision of an edge $e(u, v) \in E(G)$ is the operation of replacing e by a path (u, w, v) , where w is a new vertex.
- 2** So, to get a subdivision of e introduce a new vertex w on e .
- 3** A graph H is said to be a subdivision of G if H can be obtained from G by a sequence of edge subdivisions. (By definition, G is a subdivision of G .)




(a) Original Graph G




(b) Subdivision $S(G)$

Definition

- 1** The subdivision of an edge $e(u, v) \in E(G)$ is the operation of replacing e by a path (u, w, v) , where w is a new vertex.
- 2** So, to get a subdivision of e introduce a new vertex w on e .
- 3** A graph H is said to be a subdivision of G if H can be obtained from G by a sequence of edge subdivisions. (By definition, G is a subdivision of G .)




(a) Original Graph G




(b) Subdivision $S(G)$

Definition

- 1** The subdivision of an edge $e(u, v) \in E(G)$ is the operation of replacing e by a path (u, w, v) , where w is a new vertex.
- 2** So, to get a subdivision of e introduce a new vertex w on e .
- 3** A graph H is said to be a subdivision of G if H can be obtained from G by a sequence of edge subdivisions. (By definition, G is a subdivision of G .)



(a) Original Graph G



(b) Subdivision $S(G)$

- 1** Any subdivision of G is denoted by $S(G)$.
- 2 Note that $S(G)$ is not unique.
- 3 In fact, there are infinite number of subdivisions of any graph with at least one edge.

- 1 Any subdivision of G is denoted by $S(G)$.
- 2 Note that $S(G)$ is not unique.
- 3 In fact, there are infinite number of subdivisions of any graph with at least one edge.

- 1 Any subdivision of G is denoted by $S(G)$.
- 2 Note that $S(G)$ is not unique.
- 3 In fact, there are infinite number of subdivisions of any graph with at least one edge.

- 1 $H \subseteq G$; G is planar $\implies H$ is planar.
- 2 $H \subseteq G$; H is nonplanar $\implies G$ is nonplanar.
- 3 G is planar $\implies S(G)$ is planar.
- 4 G is nonplanar $\implies S(G)$ is nonplanar.
- 5 $S(K_5)$ and $S(K_{3,3})$ are nonplanar.
- 6 G is planar $\implies S(K_5), S(K_{3,3}) \not\subseteq G$.

- 1 $H \subseteq G$; G is planar $\implies H$ is planar.
- 2 $H \subseteq G$; H is nonplanar $\implies G$ is nonplanar.
- 3 G is planar $\implies S(G)$ is planar.
- 4 G is nonplanar $\implies S(G)$ is nonplanar.
- 5 $S(K_5)$ and $S(K_{3,3})$ are nonplanar.
- 6 G is planar $\implies S(K_5), S(K_{3,3}) \not\subseteq G$.

- 1 $H \subseteq G$; G is planar $\implies H$ is planar.
- 2 $H \subseteq G$; H is nonplanar $\implies G$ is nonplanar.
- 3 G is planar $\implies S(G)$ is planar.
- 4 G is nonplanar $\implies S(G)$ is nonplanar.
- 5 $S(K_5)$ and $S(K_{3,3})$ are nonplanar.
- 6 G is planar $\implies S(K_5), S(K_{3,3}) \not\subseteq G$.

- 1 $H \subseteq G$; G is planar $\implies H$ is planar.
- 2 $H \subseteq G$; H is nonplanar $\implies G$ is nonplanar.
- 3 G is planar $\implies S(G)$ is planar.
- 4 G is nonplanar $\implies S(G)$ is nonplanar.
- 5 $S(K_5)$ and $S(K_{3,3})$ are nonplanar.
- 6 G is planar $\implies S(K_5), S(K_{3,3}) \not\subseteq G$.

- 1 $H \subseteq G$; G is planar $\implies H$ is planar.
- 2 $H \subseteq G$; H is nonplanar $\implies G$ is nonplanar.
- 3 G is planar $\implies S(G)$ is planar.
- 4 G is nonplanar $\implies S(G)$ is nonplanar.
- 5 $S(K_5)$ and $S(K_{3,3})$ are nonplanar.
- 6 G is planar $\implies S(K_5), S(K_{3,3}) \not\subseteq G$.

- 1 $H \subseteq G$; G is planar $\implies H$ is planar.
- 2 $H \subseteq G$; H is nonplanar $\implies G$ is nonplanar.
- 3 G is planar $\implies S(G)$ is planar.
- 4 G is nonplanar $\implies S(G)$ is nonplanar.
- 5 $S(K_5)$ and $S(K_{3,3})$ are nonplanar.
- 6 G is planar $\implies S(K_5), S(K_{3,3}) \not\subseteq G$.

Kuratowski's Theorem (1930)

- 1 The two example non-planar graphs $K_{3,3}$ and K_5 weren't picked randomly.
- 2 Kuratowski proved that any non-planar graph must contain a subgraph closely related to one of these two graphs.

Theorem

Kuratowski's Theorem: A graph is nonplanar if and only if it contains a subgraph that is a subdivision of $K_{3,3}$ or K_5 .

Petersen graph non-planar.

Find a subdivision of either K_5 or $K_{3,3}$

Kuratowski's Theorem (1930)

- 1 The two example non-planar graphs $K_{3,3}$ and K_5 weren't picked randomly.
- 2 Kuratowski proved that any non-planar graph must contain a subgraph closely related to one of these two graphs.

Theorem

Kuratowski's Theorem: A graph is nonplanar if and only if it contains a subgraph that is a subdivision of $K_{3,3}$ or K_5 .

Petersen graph non-planar.

Find a subdivision of either K_5 or $K_{3,3}$

Kuratowski's Theorem (1930)

- 1 The two example non-planar graphs $K_{3,3}$ and K_5 weren't picked randomly.
- 2 Kuratowski proved that any non-planar graph must contain a subgraph closely related to one of these two graphs.

Theorem

Kuratowski's Theorem: A graph is nonplanar if and only if it contains a subgraph that is a subdivision of $K_{3,3}$ or K_5 .

Petersen graph non-planar.

Find a subdivision of either K_5 or $K_{3,3}$

Kuratowski's Theorem (1930)

- 1 The two example non-planar graphs $K_{3,3}$ and K_5 weren't picked randomly.
- 2 Kuratowski proved that any non-planar graph must contain a subgraph closely related to one of these two graphs.

Theorem

Kuratowski's Theorem: A graph is nonplanar if and only if it contains a subgraph that is a subdivision of $K_{3,3}$ or K_5 .

Petersen graph non-planar.

Find a subdivision of either K_5 or $K_{3,3}$

Kuratowski's Theorem (1930)

- 1 The two example non-planar graphs $K_{3,3}$ and K_5 weren't picked randomly.
- 2 Kuratowski proved that any non-planar graph must contain a subgraph closely related to one of these two graphs.

Theorem

Kuratowski's Theorem: A graph is nonplanar if and only if it contains a subgraph that is a subdivision of $K_{3,3}$ or K_5 .

Petersen graph non-planar.

Find a subdivision of either K_5 or $K_{3,3}$

4-color theorem

Planar maps and Planar graphs first appeared in a problem called the four color conjecture (1850).

Theorem

Four-Color-Conjecture : Any map of a country can be colored with at most four colors so that no two adjacent states receive the same color.

- 1 solved by K. Appel, W. Haken and J. Koch (1977).
- 2 Their proof techniques involved making of a large number of cases by a computer. (700 pages)
- 3 A shorter, independent proof was constructed by Robertson et al. (1996) and Thomas (1998).(100 pages)

4-color theorem

Planar maps and Planar graphs first appeared in a problem called the four color conjecture (1850).

Theorem

***Four-Color-Conjecture :** Any map of a country can be colored with at most four colors so that no two adjacent states receive the same color.*

- 1 solved by K. Appel, W. Haken and J. Koch (1977).
- 2 Their proof techniques involved making of a large number of cases by a computer. (700 pages)
- 3 A shorter, independent proof was constructed by Robertson et al. (1996) and Thomas (1998).(100 pages)

4-color theorem

Planar maps and Planar graphs first appeared in a problem called the four color conjecture (1850).

Theorem

***Four-Color-Conjecture :** Any map of a country can be colored with at most four colors so that no two adjacent states receive the same color.*

- 1** solved by K. Appel, W. Haken and J. Koch (1977).
- 2 Their proof techniques involved making of a large number of cases by a computer. (700 pages)
- 3 A shorter, independent proof was constructed by Robertson et al. (1996) and Thomas (1998).(100 pages)

4-color theorem

Planar maps and Planar graphs first appeared in a problem called the four color conjecture (1850).

Theorem

***Four-Color-Conjecture** : Any map of a country can be colored with at most four colors so that no two adjacent states receive the same color.*

- 1** solved by K. Appel, W. Haken and J. Koch (1977).
- 2** Their proof techniques involved making of a large number of cases by a computer. (700 pages)
- 3** A shorter, independent proof was constructed by Robertson et al. (1996) and Thomas (1998). (100 pages)

4-color theorem

Planar maps and Planar graphs first appeared in a problem called the four color conjecture (1850).

Theorem

***Four-Color-Conjecture** : Any map of a country can be colored with at most four colors so that no two adjacent states receive the same color.*

- 1** solved by K. Appel, W. Haken and J. Koch (1977).
- 2** Their proof techniques involved making of a large number of cases by a computer. (700 pages)
- 3** A shorter, independent proof was constructed by Robertson et al. (1996) and Thomas (1998).(100 pages)