

Independent Sets and Cliques

October 8, 2024

Definition

- 1 A subset S of V is called an **independent set** of G if no two vertices of S are adjacent in G .
- 2 An independent set is maximum if G has no independent set S' with $|S'| > |S|$.
- 3 Recall that a subset K of V such that every edge of G has at least one end in K is called a covering of G .


Definition

- 1 A subset S of V is called an **independent set** of G if no two vertices of S are adjacent in G .
- 2 An independent set is maximum if G has no independent set S' with $|S'| > |S|$.
- 3 Recall that a subset K of V such that every edge of G has at least one end in K is called a covering of G .

Definition

- 1 A subset S of V is called an **independent set** of G if no two vertices of S are adjacent in G .
- 2 An independent set is maximum if G has no independent set S' with $|S'| > |S|$.
- 3 Recall that a subset K of V such that every edge of G has at least one end in K is called a covering of G .

Example



- 1 Independent set $S_1 = \{a, e\}$, and Covering $S_1^c = \{c, b, d\}$
- 2 Independent set $S_2 = \{b, c\}$, and Covering $S_2^c = \{a, d, e\}$.

Theorem

Theorem

A set $S \subseteq V$ is an independent set of G if and only if $V \setminus S = S^c$ is a covering of G .

Proof.

- 1 By definition, S is an independent set of G
- 2 if and only if no edge of G has both ends in S
- 3 if and only if each edge has at least one end in $V \setminus S = S^c$.
- 4 if and only if $V \setminus S = S^c$ is a covering of G .



Theorem

Theorem

A set $S \subseteq V$ is an independent set of G if and only if $V \setminus S = S^c$ is a covering of G .

Proof.

- 1** By definition, S is an independent set of G
- 2 if and only if no edge of G has both ends in S
- 3 if and only if each edge has at least one end in $V \setminus S = S^c$.
- 4 if and only if $V \setminus S = S^c$ is a covering of G .



Theorem

Theorem

A set $S \subseteq V$ is an independent set of G if and only if $V \setminus S = S^c$ is a covering of G .

Proof.

- 1** By definition, S is an independent set of G
- 2** if and only if no edge of G has both ends in S
- 3** if and only if each edge has at least one end in $V \setminus S = S^c$.
- 4** if and only if $V \setminus S = S^c$ is a covering of G .



Theorem

Theorem

A set $S \subseteq V$ is an independent set of G if and only if $V \setminus S = S^c$ is a covering of G .

Proof.

- 1** By definition, S is an independent set of G
- 2** if and only if no edge of G has both ends in S
- 3** if and only if each edge has at least one end in $V \setminus S = S^c$.
- 4** if and only if $V \setminus S = S^c$ is a covering of G .



Theorem

Theorem

A set $S \subseteq V$ is an independent set of G if and only if $V \setminus S = S^c$ is a covering of G .

Proof.


- 1** By definition, S is an independent set of G
- 2** if and only if no edge of G has both ends in S
- 3** if and only if each edge has at least one end in $V \setminus S = S^c$.
- 4** if and only if $V \setminus S = S^c$ is a covering of G .



Independence and Covering Number

Definition

- 1 The number of vertices in a maximum independent set of G is called the **independence number** of G and is denoted by $\alpha(G)$.
- 2 The number of vertices in a minimum covering of G is the **covering number** of G and is denoted by $\beta(G)$.




$$\alpha(G) = 2 \text{ and } \beta(G) = 3$$

Independence and Covering Number

Definition

- 1 The number of vertices in a maximum independent set of G is called the **independence number** of G and is denoted by $\alpha(G)$.
- 2 The number of vertices in a minimum covering of G is the **covering number** of G and is denoted by $\beta(G)$.



$$\alpha(G) = 2 \text{ and } \beta(G) = 3$$

Corollary

Corollary

$$\alpha + \beta = |V(G)| = n.$$

Proof.

- 1 Let S be a maximum independent set of G .
- 2 Let K be a minimum covering of G .
- 3 By Previous theorem, $V \setminus K$ is an independent set and $V \setminus S$ is a covering of G .
- 4 $\beta \leq |V \setminus S| = n - \alpha, \Rightarrow n \geq \alpha + \beta.$
- 5 $\alpha \geq |V \setminus K| = n - \beta, \Rightarrow n \leq \alpha + \beta.$
- 6 $\Rightarrow n = \alpha + \beta$



Corollary

Corollary

$$\alpha + \beta = |V(G)| = n.$$

Proof.

- 1** Let S be a maximum independent set of G .
- 2** Let K be a minimum covering of G .
- 3** By Previous theorem, $V \setminus K$ is an independent set and $V \setminus S$ is a covering of G .
- 4** $\beta \leq |V \setminus S| = n - \alpha, \Rightarrow n \geq \alpha + \beta.$
- 5** $\alpha \geq |V \setminus K| = n - \beta, \Rightarrow n \leq \alpha + \beta.$
- 6** $\Rightarrow n = \alpha + \beta$



Corollary

Corollary

$$\alpha + \beta = |V(G)| = n.$$

Proof.

- 1** Let S be a maximum independent set of G .
- 2** Let K be a minimum covering of G .
- 3** By Previous theorem, $V \setminus K$ is an independent set and $V \setminus S$ is a covering of G .
- 4** $\beta \leq |V \setminus S| = n - \alpha =, \Rightarrow n \geq \alpha + \beta.$
- 5** $\alpha \geq |V \setminus K| = n - \beta, \Rightarrow n \leq \alpha + \beta.$
- 6** $\Rightarrow n = \alpha + \beta$



Corollary

Corollary

$$\alpha + \beta = |V(G)| = n.$$

Proof.

- 1** Let S be a maximum independent set of G .
- 2** Let K be a minimum covering of G .
- 3** By Previous theorem, $V \setminus K$ is an independent set and $V \setminus S$ is a covering of G .
- 4** $\beta \leq |V \setminus S| = n - \alpha =, \Rightarrow n \geq \alpha + \beta.$
- 5** $\alpha \geq |V \setminus K| = n - \beta, \Rightarrow n \leq \alpha + \beta.$
- 6** $\Rightarrow n = \alpha + \beta$



Corollary

Corollary

$$\alpha + \beta = |V(G)| = n.$$

Proof.

- 1** Let S be a maximum independent set of G .
- 2** Let K be a minimum covering of G .
- 3** By Previous theorem, $V \setminus K$ is an independent set and $V \setminus S$ is a covering of G .
- 4** $\beta \leq |V \setminus S| = n - \alpha, \Rightarrow n \geq \alpha + \beta.$
- 5** $\alpha \geq |V \setminus K| = n - \beta, \Rightarrow n \leq \alpha + \beta.$
- 6** $\Rightarrow n = \alpha + \beta$



Corollary

Corollary

$$\alpha + \beta = |V(G)| = n.$$

Proof.

- 1** Let S be a maximum independent set of G .
- 2** Let K be a minimum covering of G .
- 3** By Previous theorem, $V \setminus K$ is an independent set and $V \setminus S$ is a covering of G .
- 4** $\beta \leq |V \setminus S| = n - \alpha, \Rightarrow n \geq \alpha + \beta.$
- 5** $\alpha \geq |V \setminus K| = n - \beta, \Rightarrow n \leq \alpha + \beta.$
- 6** $\Rightarrow n = \alpha + \beta$



Corollary

Corollary

$$\alpha + \beta = |V(G)| = n.$$

Proof.


- 1** Let S be a maximum independent set of G .
- 2** Let K be a minimum covering of G .
- 3** By Previous theorem, $V \setminus K$ is an independent set and $V \setminus S$ is a covering of G .
- 4** $\beta \leq |V \setminus S| = n - \alpha, \Rightarrow n \geq \alpha + \beta.$
- 5** $\alpha \geq |V \setminus K| = n - \beta, \Rightarrow n \leq \alpha + \beta.$
- 6** $\Rightarrow n = \alpha + \beta$



Edge Analogues

Definition

- 1** The edge analogue of an independent set is a set of links no two of which are adjacent, that is, a **matching**.
- 2** The edge analogue of a covering is called an **edge covering**. An **edge covering** of G is a subset L of E such that each vertex of G is an end of some edge in L .
- 3** An edge-cover F is called a **minimal edge-cover** if there is no edge-cover which is properly contained in F .




Edge Cover : $\{ab, cd, be\}$

Edge Analogues

Definition

- 1** The edge analogue of an independent set is a set of links no two of which are adjacent, that is, a **matching**.
- 2** The edge analogue of a covering is called an **edge covering**. An **edge covering** of G is a subset L of E such that each vertex of G is an end of some edge in L .
- 3** An edge-cover F is called a **minimal edge-cover** if there is no edge-cover which is properly contained in F .




Edge Cover : $\{ab, cd, be\}$

Edge Analogues

Definition

- 1** The edge analogue of an independent set is a set of links no two of which are adjacent, that is, a **matching**.
- 2** The edge analogue of a covering is called an **edge covering**. An **edge covering** of G is a subset L of E such that each vertex of G is an end of some edge in L .
- 3** An edge-cover F is called a **minimal edge-cover** if there is no edge-cover which is properly contained in F .



Edge Cover : $\{ab, cd, be\}$

- 1 Note that edge coverings do not always exist; a graph G has an edge covering if and only if $\delta(G) > 0$.
- 2 $\alpha'(G)$ = number of edges in a maximum matching of G
- 3 $\beta'(G)$ = number of edges in a minimum edge covering of G
- 4 $\alpha'(G)$ = edge independence number
- 5 $\beta'(G)$ = edge covering number of G .

- 1 Note that edge coverings do not always exist; a graph G has an edge covering if and only if $\delta(G) > 0$.
- 2 $\alpha'(G)$ = number of edges in a maximum matching of G
- 3 $\beta'(G)$ = number of edges in a minimum edge covering of G
- 4 $\alpha'(G)$ = edge independence number
- 5 $\beta'(G)$ = edge covering number of G .

- 1 Note that edge coverings do not always exist; a graph G has an edge covering if and only if $\delta(G) > 0$.
- 2 $\alpha'(G)$ = number of edges in a maximum matching of G
- 3 $\beta'(G)$ = number of edges in a minimum edge covering of G
- 4 $\alpha'(G)$ = edge independence number
- 5 $\beta'(G)$ = edge covering number of G .

- 1 Note that edge coverings do not always exist; a graph G has an edge covering if and only if $\delta(G) > 0$.
- 2 $\alpha'(G)$ = number of edges in a maximum matching of G
- 3 $\beta'(G)$ = number of edges in a minimum edge covering of G
- 4 $\alpha'(G)$ = edge independence number
- 5 $\beta'(G)$ = edge covering number of G .

- 1 Note that edge coverings do not always exist; a graph G has an edge covering if and only if $\delta(G) > 0$.
- 2 $\alpha'(G)$ = number of edges in a maximum matching of G
- 3 $\beta'(G)$ = number of edges in a minimum edge covering of G
- 4 $\alpha'(G)$ = edge independence number
- 5 $\beta'(G)$ = edge covering number of G .

- 1 Observe that a graph G may contain many maximal and maximum matchings.
- 2 It also may contain many minimal and minimum edge covers.
- 3 But $\alpha'(G)$ and $\beta'(G)$ are unique.

- 1 Observe that a graph G may contain many maximal and maximum matchings.
- 2 It also may contain many minimal and minimum edge covers.
- 3 But $\alpha'(G)$ and $\beta'(G)$ are unique.

- 1 Observe that a graph G may contain many maximal and maximum matchings.
- 2 It also may contain many minimal and minimum edge covers.
- 3 But $\alpha'(G)$ and $\beta'(G)$ are unique.


- 1 Matchings and edge coverings are not related to one another as simply as are independent sets and coverings
- 2 The complement of a matching need not be an edge cover.
- 3 The complement of an edge covering is not necessarily a matching.
- 4 However, it so happens that the parameters $\alpha'(G)$ and $\beta'(G)$ are related in precisely the same manner as are α and β .

- 1 Matchings and edge coverings are not related to one another as simply as are independent sets and coverings
- 2 The complement of a matching need not be an edge cover.
- 3 The complement of an edge covering is not necessarily a matching.
- 4 However, it so happens that the parameters $\alpha'(G)$ and $\beta'(G)$ are related in precisely the same manner as are α and β .

- 1 Matchings and edge coverings are not related to one another as simply as are independent sets and coverings
- 2 The complement of a matching need not be an edge cover.
- 3 The complement of an edge covering is not necessarily a matching.
- 4 However, it so happens that the parameters $\alpha'(G)$ and $\beta'(G)$ are related in precisely the same manner as are α and β .

- 1 Matchings and edge coverings are not related to one another as simply as are independent sets and coverings
- 2 The complement of a matching need not be an edge cover.
- 3 The complement of an edge covering is not necessarily a matching.
- 4 However, it so happens that the parameters $\alpha'(G)$ and $\beta'(G)$ are related in precisely the same manner as are α and β .


Example



Edge Cover : $K = \{ab, cd, be\}$

Matching : $\neq E \setminus K = \{ac, ad, bd, de\}$

Example



Edge Cover : $K = \{ab, cd, be\}$

Matching : $\neq E \setminus K = \{ac, ad, bd, de\}$

Gallai's Theorem

Theorem

If $\delta(G) > 0$, then $\alpha'(G) + \beta'(G) = n = |V(G)|$.

Proof.

Proof consists of two steps.

1 $\alpha'(G) + \beta'(G) \leq n = |V(G)|$

2 $\alpha'(G) + \beta'(G) \geq n = |V(G)|$



Gallai's Theorem

Theorem

If $\delta(G) > 0$, then $\alpha'(G) + \beta'(G) = n = |V(G)|$.

Proof.

Proof consists of two steps.

1 $\alpha'(G) + \beta'(G) \leq n = |V(G)|$

2 $\alpha'(G) + \beta'(G) \geq n = |V(G)|$



Claim : $\alpha'(G) + \beta'(G) \leq n = |V(G)|$

- 1 Let M be a maximum matching.
- 2 Let U be the set of all M -unsaturated vertices.
- 3 Then, $|U| = n - 2|M|$.
- 4 Let $U = \{v_1, v_2, \dots, v_p\}$, where $p = n - 2|M|$.
- 5 Let e_i be an edge incident to v_i , $i = 1, \dots, p$.
- 6 such an edge exists, since $\delta(G) > 0$.
- 7 Then $M \cup \{e_1, e_2, \dots, e_p\}$ is an edge-cover.
- 8 So,
$$\beta'(G) \leq |M| + p \leq |M| + (n - 2|M|) = n - |M| = n - \alpha'(G)$$
- 9 Hence, $\alpha'(G) + \beta'(G) \leq n$

Claim : $\alpha'(G) + \beta'(G) \leq n = |V(G)|$

1 Let M be a maximum matching.

2 Let U be the set of all M -unsaturated vertices.

3 Then, $|U| = n - 2|M|$.

4 Let $U = \{v_1, v_2, \dots, v_p\}$, where $p = n - 2|M|$.

5 Let e_i be an edge incident to v_i , $i = 1, \dots, p$.

6 such an edge exists, since $\delta(G) > 0$.

7 Then $M \cup \{e_1, e_2, \dots, e_p\}$ is an edge-cover.

8 So,

$$\beta'(G) \leq |M| + p \leq |M| + (n - 2|M|) = n - |M| = n - \alpha'(G)$$

9 Hence, $\alpha'(G) + \beta'(G) \leq n$

Claim : $\alpha'(G) + \beta'(G) \leq n = |V(G)|$

1 Let M be a maximum matching.

2 Let U be the set of all M -unsaturated vertices.

3 Then, $|U| = n - 2|M|$.

4 Let $U = \{v_1, v_2, \dots, v_p\}$, where $p = n - 2|M|$.

5 Let e_i be an edge incident to v_i , $i = 1, \dots, p$.

6 such an edge exists, since $\delta(G) > 0$.

7 Then $M \cup \{e_1, e_2, \dots, e_p\}$ is an edge-cover.

8 So,

$$\beta'(G) \leq |M| + p \leq |M| + (n - 2|M|) = n - |M| = n - \alpha'(G)$$

9 Hence, $\alpha'(G) + \beta'(G) \leq n$

Claim : $\alpha'(G) + \beta'(G) \leq n = |V(G)|$

1 Let M be a maximum matching.

2 Let U be the set of all M -unsaturated vertices.

3 Then, $|U| = n - 2|M|$.

4 Let $U = \{v_1, v_2, \dots, v_p\}$, where $p = n - 2|M|$.

5 Let e_i be an edge incident to v_i , $i = 1, \dots, p$.

6 such an edge exists, since $\delta(G) > 0$.

7 Then $M \cup \{e_1, e_2, \dots, e_p\}$ is an edge-cover.

8 So,

$$\beta'(G) \leq |M| + p \leq |M| + (n - 2|M|) = n - |M| = n - \alpha'(G)$$

9 Hence, $\alpha'(G) + \beta'(G) \leq n$

Claim : $\alpha'(G) + \beta'(G) \leq n = |V(G)|$

- 1** Let M be a maximum matching.
- 2** Let U be the set of all M -unsaturated vertices.
- 3** Then, $|U| = n - 2|M|$.
- 4** Let $U = \{v_1, v_2, \dots, v_p\}$, where $p = n - 2|M|$.
- 5** Let e_i be an edge incident to v_i , $i = 1, \dots, p$.
- 6** such an edge exists, since $\delta(G) > 0$.
- 7** Then $M \cup \{e_1, e_2, \dots, e_p\}$ is an edge-cover.
- 8** So,
$$\beta'(G) \leq |M| + p \leq |M| + (n - 2|M|) = n - |M| = n - \alpha'(G)$$
- 9** Hence, $\alpha'(G) + \beta'(G) \leq n$

Claim : $\alpha'(G) + \beta'(G) \leq n = |V(G)|$

- 1** Let M be a maximum matching.
- 2** Let U be the set of all M -unsaturated vertices.
- 3** Then, $|U| = n - 2|M|$.
- 4** Let $U = \{v_1, v_2, \dots, v_p\}$, where $p = n - 2|M|$.
- 5** Let e_i be an edge incident to v_i , $i = 1, \dots, p$.
- 6** such an edge exists, since $\delta(G) > 0$.
- 7** Then $M \cup \{e_1, e_2, \dots, e_p\}$ is an edge-cover.
- 8** So,
$$\beta'(G) \leq |M| + p \leq |M| + (n - 2|M|) = n - |M| = n - \alpha'(G)$$
- 9** Hence, $\alpha'(G) + \beta'(G) \leq n$

Claim : $\alpha'(G) + \beta'(G) \leq n = |V(G)|$

- 1** Let M be a maximum matching.
- 2** Let U be the set of all M -unsaturated vertices.
- 3** Then, $|U| = n - 2|M|$.
- 4** Let $U = \{v_1, v_2, \dots, v_p\}$, where $p = n - 2|M|$.
- 5** Let e_i be an edge incident to v_i , $i = 1, \dots, p$.
- 6** such an edge exists, since $\delta(G) > 0$.
- 7** Then $M \cup \{e_1, e_2, \dots, e_p\}$ is an edge-cover.
- 8** So,
$$\beta'(G) \leq |M| + p \leq |M| + (n - 2|M|) = n - |M| = n - \alpha'(G)$$
- 9** Hence, $\alpha'(G) + \beta'(G) \leq n$

Claim : $\alpha'(G) + \beta'(G) \leq n = |V(G)|$

- 1** Let M be a maximum matching.
- 2** Let U be the set of all M -unsaturated vertices.
- 3** Then, $|U| = n - 2|M|$.
- 4** Let $U = \{v_1, v_2, \dots, v_p\}$, where $p = n - 2|M|$.
- 5** Let e_i be an edge incident to v_i , $i = 1, \dots, p$.
- 6** such an edge exists, since $\delta(G) > 0$.
- 7** Then $M \cup \{e_1, e_2, \dots, e_p\}$ is an edge-cover.
- 8** So,
$$\beta'(G) \leq |M| + p \leq |M| + (n - 2|M|) = n - |M| = n - \alpha'(G)$$
- 9** Hence, $\alpha'(G) + \beta'(G) \leq n$

Claim : $\alpha'(G) + \beta'(G) \leq n = |V(G)|$

- 1** Let M be a maximum matching.
- 2** Let U be the set of all M -unsaturated vertices.
- 3** Then, $|U| = n - 2|M|$.
- 4** Let $U = \{v_1, v_2, \dots, v_p\}$, where $p = n - 2|M|$.
- 5** Let e_i be an edge incident to v_i , $i = 1, \dots, p$.
- 6** such an edge exists, since $\delta(G) > 0$.
- 7** Then $M \cup \{e_1, e_2, \dots, e_p\}$ is an edge-cover.
- 8** So,
$$\beta'(G) \leq |M| + p \leq |M| + (n - 2|M|) = n - |M| = n - \alpha'(G)$$
- 9** Hence, $\alpha'(G) + \beta'(G) \leq n$

Claim : $\alpha'(G) + \beta'(G) \leq n = |V(G)|$

- 1** Let M be a maximum matching.
- 2** Let U be the set of all M -unsaturated vertices.
- 3** Then, $|U| = n - 2|M|$.
- 4** Let $U = \{v_1, v_2, \dots, v_p\}$, where $p = n - 2|M|$.
- 5** Let e_i be an edge incident to v_i , $i = 1, \dots, p$.
- 6** such an edge exists, since $\delta(G) > 0$.
- 7** Then $M \cup \{e_1, e_2, \dots, e_p\}$ is an edge-cover.
- 8** So,
$$\beta'(G) \leq |M| + p \leq |M| + (n - 2|M|) = n - |M| = n - \alpha'(G)$$
- 9** Hence, $\alpha'(G) + \beta'(G) \leq n$

Claim : $\alpha'(G) + \beta'(G) \geq n = |V(G)|$

- 1 Let F be a minimum edge cover.
- 2 Let H be the spanning subgraph of G with edge set F .
- 3 Let M_H be a maximum matching in H .
- 4 Let U be the set of M_H unsaturated vertices in H .
- 5 As before, $|U| = n - 2|M_H|$.
- 6 Let $U = \{v_1, v_2, \dots, v_p\}$ where $p = n - 2|M_H|$.

Claim : $\alpha'(G) + \beta'(G) \geq n = |V(G)|$

- 1** Let F be a minimum edge cover.
- 2 Let H be the spanning subgraph of G with edge set F .
- 3 Let M_H be a maximum matching in H .
- 4 Let U be the set of M_H unsaturated vertices in H .
- 5 As before, $|U| = n - 2|M_H|$.
- 6 Let $U = \{v_1, v_2, \dots, v_p\}$ where $p = n - 2|M_H|$.

Claim : $\alpha'(G) + \beta'(G) \geq n = |V(G)|$

- 1** Let F be a minimum edge cover.
- 2** Let H be the spanning subgraph of G with edge set F .
- 3** Let M_H be a maximum matching in H .
- 4** Let U be the set of M_H unsaturated vertices in H .
- 5** As before, $|U| = n - 2|M_H|$.
- 6** Let $U = \{v_1, v_2, \dots, v_p\}$ where $p = n - 2|M_H|$.

Claim : $\alpha'(G) + \beta'(G) \geq n = |V(G)|$

- 1** Let F be a minimum edge cover.
- 2** Let H be the spanning subgraph of G with edge set F .
- 3** Let M_H be a maximum matching in H .
- 4** Let U be the set of M_H unsaturated vertices in H .
- 5** As before, $|U| = n - 2|M_H|$.
- 6** Let $U = \{v_1, v_2, \dots, v_p\}$ where $p = n - 2|M_H|$.

Claim : $\alpha'(G) + \beta'(G) \geq n = |V(G)|$

- 1** Let F be a minimum edge cover.
- 2** Let H be the spanning subgraph of G with edge set F .
- 3** Let M_H be a maximum matching in H .
- 4** Let U be the set of M_H unsaturated vertices in H .
- 5** As before, $|U| = n - 2|M_H|$.
- 6** Let $U = \{v_1, v_2, \dots, v_p\}$ where $p = n - 2|M_H|$.

Claim : $\alpha'(G) + \beta'(G) \geq n = |V(G)|$

- 1** Let F be a minimum edge cover.
- 2** Let H be the spanning subgraph of G with edge set F .
- 3** Let M_H be a maximum matching in H .
- 4** Let U be the set of M_H unsaturated vertices in H .
- 5** As before, $|U| = n - 2|M_H|$.
- 6** Let $U = \{v_1, v_2, \dots, v_p\}$ where $p = n - 2|M_H|$.

Claim : $\alpha'(G) + \beta'(G) \geq n = |V(G)|$

- 1** Let F be a minimum edge cover.
- 2** Let H be the spanning subgraph of G with edge set F .
- 3** Let M_H be a maximum matching in H .
- 4** Let U be the set of M_H unsaturated vertices in H .
- 5** As before, $|U| = n - 2|M_H|$.
- 6** Let $U = \{v_1, v_2, \dots, v_p\}$ where $p = n - 2|M_H|$.

- 1 Since, F is an edge-cover, it contains an edge e_i incident with v_i , $1 \leq i \leq p$.
- 2 Since, M_H is a maximum matching, U is an independent set in H .
- 3 Therefore $e_i \neq e_j$, if $v_i \neq v_j$.
- 4 Hence, e_1, e_2, \dots, e_p are all distinct edges.
- 5 It follows that
$$|F| \geq |M_H| + p = |M_H| + n - 2|M_H| = n - |M_H|.$$
- 6 Hence, $\beta'(G) = |F| \geq n - 2|M_H| \geq n - \alpha'(G)$, since $\alpha'(G) \geq |M_H|$.
- 7 So, $\alpha'(G) + \beta'(G) \geq n$



- 1 Since, F is an edge-cover, it contains an edge e_i incident with v_i , $1 \leq i \leq p$.
- 2 Since, M_H is a maximum matching, U is an independent set in H .
- 3 Therefore $e_i \neq e_j$, if $v_i \neq v_j$.
- 4 Hence, e_1, e_2, \dots, e_p are all distinct edges.
- 5 It follows that
$$|F| \geq |M_H| + p = |M_H| + n - 2|M_H| = n - |M_H|.$$
- 6 Hence, $\beta'(G) = |F| \geq n - 2|M_H| \geq n - \alpha'(G)$, since $\alpha'(G) \geq |M_H|$.
- 7 So, $\alpha'(G) + \beta'(G) \geq n$



- 1 Since, F is an edge-cover, it contains an edge e_i incident with v_i , $1 \leq i \leq p$.
- 2 Since, M_H is a maximum matching, U is an independent set in H .
- 3 Therefore $e_i \neq e_j$, if $v_i \neq v_j$.
- 4 Hence, e_1, e_2, \dots, e_p are all distinct edges.
- 5 It follows that
$$|F| \geq |M_H| + p = |M_H| + n - 2|M_H| = n - |M_H|.$$
- 6 Hence, $\beta'(G) = |F| \geq n - 2|M_H| \geq n - \alpha'(G)$, since $\alpha'(G) \geq |M_H|$.
- 7 So, $\alpha'(G) + \beta'(G) \geq n$



- 1 Since, F is an edge-cover, it contains an edge e_i incident with v_i , $1 \leq i \leq p$.
- 2 Since, M_H is a maximum matching, U is an independent set in H .
- 3 Therefore $e_i \neq e_j$, if $v_i \neq v_j$.
- 4 Hence, e_1, e_2, \dots, e_p are all distinct edges.
- 5 It follows that
$$|F| \geq |M_H| + p = |M_H| + n - 2|M_H| = n - |M_H|.$$
- 6 Hence, $\beta'(G) = |F| \geq n - 2|M_H| \geq n - \alpha'(G)$, since $\alpha'(G) \geq |M_H|$.
- 7 So, $\alpha'(G) + \beta'(G) \geq n$



- 1 Since, F is an edge-cover, it contains an edge e_i incident with v_i , $1 \leq i \leq p$.
- 2 Since, M_H is a maximum matching, U is an independent set in H .
- 3 Therefore $e_i \neq e_j$, if $v_i \neq v_j$.
- 4 Hence, e_1, e_2, \dots, e_p are all distinct edges.
- 5 It follows that
$$|F| \geq |M_H| + p = |M_H| + n - 2|M_H| = n - |M_H|.$$
- 6 Hence, $\beta'(G) = |F| \geq n - 2|M_H| \geq n - \alpha'(G)$, since $\alpha'(G) \geq |M_H|$.
- 7 So, $\alpha'(G) + \beta'(G) \geq n$



- 1 Since, F is an edge-cover, it contains an edge e_i incident with v_i , $1 \leq i \leq p$.
- 2 Since, M_H is a maximum matching, U is an independent set in H .
- 3 Therefore $e_i \neq e_j$, if $v_i \neq v_j$.
- 4 Hence, e_1, e_2, \dots, e_p are all distinct edges.
- 5 It follows that
$$|F| \geq |M_H| + p = |M_H| + n - 2|M_H| = n - |M_H|.$$
- 6 Hence, $\beta'(G) = |F| \geq n - 2|M_H| \geq n - \alpha'(G)$, since $\alpha'(G) \geq |M_H|$.
- 7 So, $\alpha'(G) + \beta'(G) \geq n$



- 1 Since, F is an edge-cover, it contains an edge e_i incident with v_i , $1 \leq i \leq p$.
- 2 Since, M_H is a maximum matching, U is an independent set in H .
- 3 Therefore $e_i \neq e_j$, if $v_i \neq v_j$.
- 4 Hence, e_1, e_2, \dots, e_p are all distinct edges.
- 5 It follows that
$$|F| \geq |M_H| + p = |M_H| + n - 2|M_H| = n - |M_H|.$$
- 6 Hence, $\beta'(G) = |F| \geq n - 2|M_H| \geq n - \alpha'(G)$, since $\alpha'(G) \geq |M_H|$.
- 7 So, $\alpha'(G) + \beta'(G) \geq n$



Theorem

Theorem

In a bipartite graph G with $\delta > 0$, the number of vertices in a maximum independent set is equal to the number of edges in a minimum edge covering.

Proof.

- 1 $\alpha(G) + \beta(G) = n$
- 2 $\alpha'(G) + \beta'(G) = n$
- 3 G is bipartite.
- 4 In a bipartite graph, the number of edges in a maximum matching is equal to the number of vertices in a minimum covering. (Proved earlier)
- 5 which implies, $\alpha' = \beta$.
- 6 Hence, $\alpha(G) = \beta'(G)$.

Theorem

Theorem

In a bipartite graph G with $\delta > 0$, the number of vertices in a maximum independent set is equal to the number of edges in a minimum edge covering.

Proof.

1 $\alpha(G) + \beta(G) = n$

2 $\alpha'(G) + \beta'(G) = n$

3 G is bipartite.

4 In a bipartite graph, the number of edges in a maximum matching is equal to the number of vertices in a minimum covering. (Proved earlier)

5 which implies, $\alpha' = \beta$.

6 Hence, $\alpha(G) = \beta'(G)$.

Theorem

Theorem

In a bipartite graph G with $\delta > 0$, the number of vertices in a maximum independent set is equal to the number of edges in a minimum edge covering.

Proof.

1 $\alpha(G) + \beta(G) = n$

2 $\alpha'(G) + \beta'(G) = n$

3 G is bipartite.

4 In a bipartite graph, the number of edges in a maximum matching is equal to the number of vertices in a minimum covering. (Proved earlier)

5 which implies, $\alpha' = \beta$.

6 Hence, $\alpha(G) = \beta'(G)$.

Theorem

Theorem

In a bipartite graph G with $\delta > 0$, the number of vertices in a maximum independent set is equal to the number of edges in a minimum edge covering.

Proof.

1 $\alpha(G) + \beta(G) = n$

2 $\alpha'(G) + \beta'(G) = n$

3 G is bipartite.

4 In a bipartite graph, the number of edges in a maximum matching is equal to the number of vertices in a minimum covering. (Proved earlier)

5 which implies, $\alpha' = \beta$.

6 Hence, $\alpha(G) = \beta'(G)$.

Theorem

Theorem

In a bipartite graph G with $\delta > 0$, the number of vertices in a maximum independent set is equal to the number of edges in a minimum edge covering.

Proof.

1 $\alpha(G) + \beta(G) = n$

2 $\alpha'(G) + \beta'(G) = n$

3 G is bipartite.

4 In a bipartite graph, the number of edges in a maximum matching is equal to the number of vertices in a minimum covering. (Proved earlier)

5 which implies, $\alpha' = \beta$.

6 Hence, $\alpha(G) = \beta'(G)$.

Theorem

Theorem

In a bipartite graph G with $\delta > 0$, the number of vertices in a maximum independent set is equal to the number of edges in a minimum edge covering.

Proof.

- 1 $\alpha(G) + \beta(G) = n$
- 2 $\alpha'(G) + \beta'(G) = n$
- 3 G is bipartite.
- 4 In a bipartite graph, the number of edges in a maximum matching is equal to the number of vertices in a minimum covering. (Proved earlier)
- 5 which implies, $\alpha' = \beta$.
- 6 Hence, $\alpha(G) = \beta'(G)$.

Theorem

Theorem

In a bipartite graph G with $\delta > 0$, the number of vertices in a maximum independent set is equal to the number of edges in a minimum edge covering.

Proof.

- 1 $\alpha(G) + \beta(G) = n$
- 2 $\alpha'(G) + \beta'(G) = n$
- 3 G is bipartite.
- 4 In a bipartite graph, the number of edges in a maximum matching is equal to the number of vertices in a minimum covering. (Proved earlier)
- 5 which implies, $\alpha' = \beta$.
- 6 Hence, $\alpha(G) = \beta'(G)$.