

Euler Tours

October 2, 2024

Definition

- 1** A trail in a graph which contains all its edges is called an Eulerian trail.
 - 2** A closed Eulerian trial is called an Eulerian Tour.
 - 3** A graph is called an Eulerian graph if it contains a Eulerian tour.
-
- 1** An Eulerian graph is necessarily connected.
 - 2** If G is Eulerian then one can choose an Eulerian trail starting and ending from any given vertex.

Definition

- 1** A trail in a graph which contains all its edges is called an Eulerian trail.
 - 2** A closed Eulerian trial is called an Eulerian Tour.
 - 3** A graph is called an Eulerian graph if it contains a Eulerian tour.
-
- 1** An Eulerian graph is necessarily connected.
 - 2** If G is Eulerian then one can choose an Eulerian trail starting and ending from any given vertex.

Definition

- 1** A trail in a graph which contains all its edges is called an Eulerian trail.
 - 2** A closed Eulerian trial is called an Eulerian Tour.
 - 3** A graph is called an **Eulerian graph** if it contains a Eulerian tour.
-
- 1** An Eulerian graph is necessarily connected.
 - 2** If G is Eulerian then one can choose an Eulerian trail starting and ending from any given vertex.


Definition

- 1** A trail in a graph which contains all its edges is called an Eulerian trail.
 - 2** A closed Eulerian trial is called an Eulerian Tour.
 - 3** A graph is called an **Eulerian graph** if it contains a Eulerian tour.
-
- 1** An Eulerian graph is necessarily connected.
 - 2** If G is Eulerian then one can choose an Eulerian trail starting and ending from any given vertex.

Definition

- 1** A trail in a graph which contains all its edges is called an Eulerian trail.
 - 2** A closed Eulerian trial is called an Eulerian Tour.
 - 3** A graph is called an **Eulerian graph** if it contains a Eulerian tour.
-
- 1** An Eulerian graph is necessarily connected.
 - 2** If G is Eulerian then one can choose an Eulerian trail starting and ending from any given vertex.


Example



Example of Euler Graph

Example

Non-Eulerian Graph



Theorem

Theorem

A connected graph G is Eulerian iff every vertex has even degree.

Proof.

" \Rightarrow " Assume G is Eulerian. We show that every vertex has even degree.

- 1 Let $W = (v_0, e_1, v_1, e_2, v_2, \dots, v_{t-1}, e_t, v_t (= v_0))$ be an Euler tour in G .
- 2 If v_i is an internal vertex ($\neq v_0$) of W appearing k times, then $\deg_G(v_i) = 2k$.
- 3 If v_0 appears r times internally, then $\deg_G(v_0) = 2r + 2$.



G is a graph such that every vertex is of even degree. To prove that G is Eulerian.

- 1 By induction on number of edges m .
- 2 **Base Case** : $m \leq 2$: Easy verification.
- 3 Assume true for all $m \leq k - 1$.
- 4 Consider a graph G with k edges.
- 5 x - any vertex of G .
- 6 y, z - vertices (need not be distinct) adjacent to x .
- 7 Delete edges xy and xz and add a new edge yz .
- 8 The new graph H has $\leq m - 1$ edges and its every vertex has even degree.
- 9 H may be connected or disconnected.

G is a graph such that every vertex is of even degree. To prove that G is Eulerian.

- 1 By induction on number of edges m .
- 2 **Base Case :** $m \leq 2$: Easy verification.
- 3 Assume true for all $m \leq k - 1$.
- 4 Consider a graph G with k edges.
- 5 x - any vertex of G .
- 6 y, z - vertices (need not be distinct) adjacent to x .
- 7 Delete edges xy and xz and add a new edge yz .
- 8 The new graph H has $\leq m - 1$ edges and its every vertex has even degree.
- 9 H may be connected or disconnected.

G is a graph such that every vertex is of even degree. To prove that G is Eulerian.

- 1 By induction on number of edges m .
- 2 **Base Case :** $m \leq 2$: Easy verification.
- 3 Assume true for all $m \leq k - 1$.
- 4 Consider a graph G with k edges.
- 5 x - any vertex of G .
- 6 y, z - vertices (need not be distinct) adjacent to x .
- 7 Delete edges xy and xz and add a new edge yz .
- 8 The new graph H has $\leq m - 1$ edges and its every vertex has even degree.
- 9 H may be connected or disconnected.

G is a graph such that every vertex is of even degree. To prove that G is Eulerian.

- 1 By induction on number of edges m .
- 2 **Base Case :** $m \leq 2$: Easy verification.
- 3 Assume true for all $m \leq k - 1$.
- 4 Consider a graph G with k edges.
- 5 x - any vertex of G .
- 6 y, z - vertices (need not be distinct) adjacent to x .
- 7 Delete edges xy and xz and add a new edge yz .
- 8 The new graph H has $\leq m - 1$ edges and its every vertex has even degree.
- 9 H may be connected or disconnected.

G is a graph such that every vertex is of even degree. To prove that G is Eulerian.

- 1 By induction on number of edges m .
- 2 **Base Case :** $m \leq 2$: Easy verification.
- 3 Assume true for all $m \leq k - 1$.
- 4 Consider a graph G with k edges.
- 5 x - any vertex of G .
- 6 y, z - vertices (need not be distinct) adjacent to x .
- 7 Delete edges xy and xz and add a new edge yz .
- 8 The new graph H has $\leq m - 1$ edges and its every vertex has even degree.
- 9 H may be connected or disconnected.

G is a graph such that every vertex is of even degree. To prove that G is Eulerian.

- 1 By induction on number of edges m .
- 2 **Base Case :** $m \leq 2$: Easy verification.
- 3 Assume true for all $m \leq k - 1$.
- 4 Consider a graph G with k edges.
- 5 x - any vertex of G .
- 6 y, z - vertices (need not be distinct) adjacent to x .
- 7 Delete edges xy and xz and add a new edge yz .
- 8 The new graph H has $\leq m - 1$ edges and its every vertex has even degree.
- 9 H may be connected or disconnected.

G is a graph such that every vertex is of even degree. To prove that G is Eulerian.

- 1 By induction on number of edges m .
- 2 **Base Case :** $m \leq 2$: Easy verification.
- 3 Assume true for all $m \leq k - 1$.
- 4 Consider a graph G with k edges.
- 5 x - any vertex of G .
- 6 y, z - vertices (need not be distinct) adjacent to x .
- 7 Delete edges xy and xz and add a new edge yz .
- 8 The new graph H has $\leq m - 1$ edges and its every vertex has even degree.
- 9 H may be connected or disconnected.

G is a graph such that every vertex is of even degree. To prove that G is Eulerian.

- 1 By induction on number of edges m .
- 2 **Base Case** : $m \leq 2$: Easy verification.
- 3 Assume true for all $m \leq k - 1$.
- 4 Consider a graph G with k edges.
- 5 x - any vertex of G .
- 6 y, z - vertices (need not be distinct) adjacent to x .
- 7 Delete edges xy and xz and add a new edge yz .
- 8 The new graph H has $\leq m - 1$ edges and its every vertex has even degree.
- 9 H may be connected or disconnected.

We consider two cases.

Case 1 : H is connected.

- 1 By induction hypothesis, H contains an Eulerian tour.

$$W = (v_0, e_1, v_1, e_2, v_2, \dots, y, f = yz, z, \dots, v_0)$$

- 2 Then, by replacing edge f with yx, x, xz in W we get,

$$W^* = (v_0, e_1, v_1, e_2, v_2, \dots, y, yx, x, xz, z, \dots, v_0)$$

- 3 W^* an Euler tour in G .

We consider two cases.

Case 1 : H is connected.

- 1 By induction hypothesis, H contains an Eulerian tour.

$$W = (v_0, e_1, v_1, e_2, v_2, \dots, y, f = yz, z, \dots, v_0)$$

- 2 Then, by replacing edge f with yx, x, xz in W we get,

$$W^* = (v_0, e_1, v_1, e_2, v_2, \dots, y, yx, x, xz, z, \dots, v_0)$$

- 3 W^* an Euler tour in G .

We consider two cases.

Case 1 : H is connected.

- 1** By induction hypothesis, H contains an Eulerian tour.

$$W = (v_0, e_1, v_1, e_2, v_2, \dots, y, f = yz, z, \dots, v_0)$$

- 2** Then, by replacing edge f with yx, x, xz in W we get,

$$W^* = (v_0, e_1, v_1, e_2, v_2, \dots, y, yx, x, xz, z, \dots, v_0)$$

- 3** W^* an Euler tour in G .

Case 2 : H is disconnected.

- 1 Let C and D be two components of H such that $x \in C$ and edge $f = yz \in D$.
- 2 $|E(C)| \leq m - 1$ and $|E(D)| \leq m - 1$.
- 3 By induction, C and D contain Euler tours.
- 4 $W_1 = (x, \dots, x)$ in C .
- 5 $W_2 = (v_0, e_1, v_1, e_2, v_2, \dots, y, f, z, \dots, v_0)$ in D .
- 6 Then, $W^* = (v_0, e_1, v_1, e_2, v_2, \dots, y, yx, W_1, xz, z, \dots, v_0)$ is an Eulerian tour in G .

Case 2 : H is disconnected.

- 1** Let C and D be two components of H such that $x \in C$ and edge $f = yz \in D$.
- 2** $|E(C)| \leq m - 1$ and $|E(D)| \leq m - 1$.
- 3** By induction, C and D contain Euler tours.
- 4** $W_1 = (x, \dots, x)$ in C .
- 5** $W_2 = (v_0, e_1, v_1, e_2, v_2, \dots, y, f, z, \dots, v_0)$ in D .
- 6** Then, $W^* = (v_0, e_1, v_1, e_2, v_2, \dots, y, yx, W_1, xz, z, \dots, v_0)$ is an Eulerian tour in G .

Case 2 : H is disconnected.

- 1** Let C and D be two components of H such that $x \in C$ and edge $f = yz \in D$.
- 2** $|E(C)| \leq m - 1$ and $|E(D)| \leq m - 1$.
- 3** By induction, C and D contain Euler tours.
- 4** $W_1 = (x, \dots, x)$ in C .
- 5** $W_2 = (v_0, e_1, v_1, e_2, v_2, \dots, y, f, z, \dots, v_0)$ in D .
- 6** Then, $W^* = (v_0, e_1, v_1, e_2, v_2, \dots, y, yx, W_1, xz, z, \dots, v_0)$ is an Eulerian tour in G .

Case 2 : H is disconnected.

- 1** Let C and D be two components of H such that $x \in C$ and edge $f = yz \in D$.
- 2** $|E(C)| \leq m - 1$ and $|E(D)| \leq m - 1$.
- 3** By induction, C and D contain Euler tours.
- 4** $W_1 = (x, \dots, x)$ in C .
- 5** $W_2 = (v_0, e_1, v_1, e_2, v_2, \dots, y, f, z, \dots, v_0)$ in D .
- 6** Then, $W^* = (v_0, e_1, v_1, e_2, v_2, \dots, y, yx, W_1, xz, z, \dots, v_0)$ is an Eulerian tour in G .

Case 2 : H is disconnected.

- 1** Let C and D be two components of H such that $x \in C$ and edge $f = yz \in D$.
- 2** $|E(C)| \leq m - 1$ and $|E(D)| \leq m - 1$.
- 3** By induction, C and D contain Euler tours.
- 4** $W_1 = (x, \dots, x)$ in C .
- 5** $W_2 = (v_0, e_1, v_1, e_2, v_2, \dots, y, f, z, \dots, v_0)$ in D .
- 6** Then, $W^* = (v_0, e_1, v_1, e_2, v_2, \dots, y, yx, W_1, xz, z, \dots, v_0)$ is an Eulerian tour in G .

Case 2 : H is disconnected.

- 1** Let C and D be two components of H such that $x \in C$ and edge $f = yz \in D$.
- 2** $|E(C)| \leq m - 1$ and $|E(D)| \leq m - 1$.
- 3** By induction, C and D contain Euler tours.
- 4** $W_1 = (x, \dots, x)$ in C .
- 5** $W_2 = (v_0, e_1, v_1, e_2, v_2, \dots, y, f, z, \dots, v_0)$ in D .
- 6** Then, $W^* = (v_0, e_1, v_1, e_2, v_2, \dots, y, yx, W_1, xz, z, \dots, v_0)$ is an Eulerian tour in G .

Case 2 : H is disconnected.

- 1 Let C and D be two components of H such that $x \in C$ and edge $f = yz \in D$.
- 2 $|E(C)| \leq m - 1$ and $|E(D)| \leq m - 1$.
- 3 By induction, C and D contain Euler tours.
- 4 $W_1 = (x, \dots, x)$ in C .
- 5 $W_2 = (v_0, e_1, v_1, e_2, v_2, \dots, y, f, z, \dots, v_0)$ in D .
- 6 Then, $W^* = (v_0, e_1, v_1, e_2, v_2, \dots, y, yx, W_1, xz, z, \dots, v_0)$ is an Eulerian tour in G .

Motivation : TSP

Travelling salesman Problem

Given n cities and distance between any two cities, a traveling salesman wishes to start from one of these cities, visits each city exactly once and comes back to the starting point. Design an algorithm to find a shortest route.

Motivation : TSP

Graph theoretical model of TSP


Given a weighted graph G on n vertices, design an algorithm to find a cycle with minimum weight containing all the vertices of G .

- 1** A path (cycle) in a graph containing all its vertices is called a Hamilton path (respectively Hamilton cycle).
- 2** A graph is called a **Hamilton graph** or **Hamiltonian** if it contains a Hamilton cycle.
- 3** A graph is Hamiltonian iff its underlying simple graph is Hamilton. Hence, the study of Hamilton graphs is limited to simple graphs.

- 1 A path (cycle) in a graph containing all its vertices is called a Hamilton path (respectively Hamilton cycle).
- 2 A graph is called a **Hamilton graph** or **Hamiltonian** if it contains a Hamilton cycle.
- 3 A graph is Hamiltonian iff its underlying simple graph is Hamilton. Hence, the study of Hamilton graphs is limited to simple graphs.

- 1** A path (cycle) in a graph containing all its vertices is called a Hamilton path (respectively Hamilton cycle).
- 2** A graph is called a **Hamilton graph** or **Hamiltonian** if it contains a Hamilton cycle.
- 3** A graph is Hamiltonian iff its underlying simple graph is Hamilton. Hence, the study of Hamilton graphs is limited to simple graphs.

Example



Hamilton problems(Open)

- 1** Find necessary and sufficient conditions for a graph to be Hamiltonian.
- 2** Design a polynomial-time algorithm to generate a Hamilton cycle in a given graph G or declare that G has no Hamilton cycle.

Hamilton problems(Open)

- 1** Find necessary and sufficient conditions for a graph to be Hamiltonian.
- 2** Design a polynomial-time algorithm to generate a Hamilton cycle in a given graph G or declare that G has no Hamilton cycle.

Necessary conditions and sufficient conditions

Theorem

(Necessary condition). If G is Hamiltonian, then

$$\omega(G - S) \leq |S| \text{ for every } S \subseteq V(G)$$

Proof.

Two way counting technique □

Necessary conditions and sufficient conditions

Theorem

(Necessary condition). If G is Hamiltonian, then


$$\omega(G - S) \leq |S| \text{ for every } S \subseteq V(G)$$

Proof.


Two way counting technique




- 1 Let C be a Hamilton cycle in G and let $S \subseteq V(G)$ and $|S| = s$.
- 2 Consider the set of edges $F \subseteq E(C)$ with one end in S and another end in $G - S$.
- 3 Every vertex in G is incident with two edges of C .




- 1 Let C be a Hamilton cycle in G and let $S \subseteq V(G)$ and $|S| = s$.
- 2 Consider the set of edges $F \subseteq E(C)$ with one end in S and another end in $G - S$.
- 3 Every vertex in G is incident with two edges of C .



- 1** Let C be a Hamilton cycle in G and let $S \subseteq V(G)$ and $|S| = s$.
- 2** Consider the set of edges $F \subseteq E(C)$ with one end in S and another end in $G - S$.
- 3** Every vertex in G is incident with two edges of C .



- 1** Let C be a Hamilton cycle in G and let $S \subseteq V(G)$ and $|S| = s$.
- 2** Consider the set of edges $F \subseteq E(C)$ with one end in S and another end in $G - S$.
- 3** Every vertex in G is incident with two edges of C .



1 Therefore, every vertex in S is incident with at most two edges of F .

2 $\implies |F| \leq 2s = 2|S|.$

3 Also, every component of $G - S$ is incident with at least two edges of F .

4 Hence, $|F| \geq 2\omega(G - S).$

5 $\omega(G - S) \leq \frac{|F|}{2} \leq |S|.$

6 $\implies \omega(G - S) \leq |S|$ for all $S \subseteq V(G)$



- 1 Therefore, every vertex in S is incident with at most two edges of F .
- 2 $\implies |F| \leq 2s = 2|S|$.
- 3 Also, every component of $G - S$ is incident with at least two edges of F .
- 4 Hence, $|F| \geq 2\omega(G - S)$.
- 5 $\omega(G - S) \leq \frac{|F|}{2} \leq |S|$.
- 6 $\implies \omega(G - S) \leq |S|$ for all $S \subseteq V(G)$ □

- 1 Therefore, every vertex in S is incident with at most two edges of F .
- 2 $\implies |F| \leq 2s = 2|S|$.
- 3 Also, every component of $G - S$ is incident with at least two edges of F .
- 4 Hence, $|F| \geq 2\omega(G - S)$.
- 5 $\omega(G - S) \leq \frac{|F|}{2} \leq |S|$.
- 6 $\implies \omega(G - S) \leq |S|$ for all $S \subseteq V(G)$ □


- 1 Therefore, every vertex in S is incident with at most two edges of F .
- 2 $\implies |F| \leq 2s = 2|S|$.
- 3 Also, every component of $G - S$ is incident with at least two edges of F .
- 4 Hence, $|F| \geq 2\omega(G - S)$.
- 5 $\omega(G - S) \leq \frac{|F|}{2} \leq |S|$.
- 6 $\implies \omega(G - S) \leq |S|$ for all $S \subseteq V(G)$ □

- 1 Therefore, every vertex in S is incident with at most two edges of F .
- 2 $\implies |F| \leq 2s = 2|S|$.
- 3 Also, every component of $G - S$ is incident with at least two edges of F .
- 4 Hence, $|F| \geq 2\omega(G - S)$.
- 5 $\omega(G - S) \leq \frac{|F|}{2} \leq |S|$.
- 6 $\implies \omega(G - S) \leq |S|$ for all $S \subseteq V(G)$ □

- 1 Therefore, every vertex in S is incident with at most two edges of F .
- 2 $\implies |F| \leq 2s = 2|S|$.
- 3 Also, every component of $G - S$ is incident with at least two edges of F .
- 4 Hence, $|F| \geq 2\omega(G - S)$.
- 5 $\omega(G - S) \leq \frac{|F|}{2} \leq |S|$.
- 6 $\implies \omega(G - S) \leq |S|$ for all $S \subseteq V(G)$ □

Example


One can use previous theorem to show that a given graph is **Not Hamiltonian**.



For $S = \{c, d\}$, $G - S$ has three components.

Example


One can use previous theorem to show that a given graph is **Not Hamiltonian**.



For $S = \{c, d\}$, $G - S$ has three components.

Example

One can use previous theorem to show that a given graph is **Not Hamiltonian**.



For $S = \{c, d\}$, $G - S$ has three components.

Example-Petersen Graph

Previous result is not applicable to all graphs.




Figure: Petersen Graph

For any $S \subseteq V(G)$, $\omega(G - S) \leq |S|$.

But G is not Hamiltonian.

Example-Petersen Graph

Previous result is not applicable to all graphs.




Figure: Petersen Graph

For any $S \subseteq V(G)$, $\omega(G - S) \leq |S|$.
But G is not Hamiltonian.

Example-Petersen Graph

Previous result is not applicable to all graphs.




Figure: Petersen Graph

For any $S \subseteq V(G)$, $\omega(G - S) \leq |S|$.
But G is not Hamiltonian.

Definition

A graph G is called maximal non-Hamiltonian if it satisfies the following.

- 1 G is non-Hamiltonian.
- 2 $G + uv$ is Hamiltonian for every pair of non-adjacent vertices u and v .

Definition

A graph G is called maximal non-Hamiltonian if it satisfies the following.

- 1 G is non-Hamiltonian.
- 2 $G + uv$ is Hamiltonian for every pair of non-adjacent vertices u and v .

Definition

A graph G is called maximal non-Hamiltonian if it satisfies the following.

- 1 G is non-Hamiltonian.
- 2 $G + uv$ is Hamiltonian for every pair of non-adjacent vertices u and v .

Dirac's Theorem

Theorem

If G is a simple graph with $n (\geq 3)$ vertices and $\delta(G) \geq \frac{n}{2}$ then G is Hamiltonian.

Proof.

- 1 Proof by Contradiction.
- 2 Let G be such that, G is maximal non-Hamiltonian and $\delta(G) \geq \frac{n}{2}$.
- 3 G is not complete as $n \geq 3$.
- 4 Choose $u, v \in V(G)$ and u and v are non-adjacent.
- 5 $G + uv$ is Hamiltonian.
- 6 Every Hamilton cycle of $G + uv$ contains the edge uv .
- 7 Let $P : u = v_1 - v_2 - \cdots - v_n = v$ be a Hamilton Path in G .



Dirac's Theorem

Theorem

If G is a simple graph with $n (\geq 3)$ vertices and $\delta(G) \geq \frac{n}{2}$ then G is Hamiltonian.

Proof.

- 1** Proof by Contradiction.
- 2** Let G be such that, G is maximal non-Hamiltonian and $\delta(G) \geq \frac{n}{2}$.
- 3** G is not complete as $n \geq 3$.
- 4** Choose $u, v \in V(G)$ and u and v are non-adjacent.
- 5** $G + uv$ is Hamiltonian.
- 6** Every Hamilton cycle of $G + uv$ contains the edge uv .
- 7** Let $P : u = v_1 - v_2 - \cdots - v_n = v$ be a Hamilton Path in G .



Dirac's Theorem

Theorem

If G is a simple graph with $n (\geq 3)$ vertices and $\delta(G) \geq \frac{n}{2}$ then G is Hamiltonian.

Proof.

- 1** Proof by Contradiction.
- 2** Let G be such that, G is maximal non-Hamiltonian and $\delta(G) \geq \frac{n}{2}$.
- 3** G is not complete as $n \geq 3$.
- 4** Choose $u, v \in V(G)$ and u and v are non-adjacent.
- 5** $G + uv$ is Hamiltonian.
- 6** Every Hamilton cycle of $G + uv$ contains the edge uv .
- 7** Let $P : u = v_1 - v_2 - \cdots - v_n = v$ be a Hamilton Path in G .



Dirac's Theorem

Theorem

If G is a simple graph with n (≥ 3) vertices and $\delta(G) \geq \frac{n}{2}$ then G is Hamiltonian.

Proof.

- 1** Proof by Contradiction.
- 2** Let G be such that, G is maximal non-Hamiltonian and $\delta(G) \geq \frac{n}{2}$.
- 3** G is not complete as $n \geq 3$.
- 4** Choose $u, v \in V(G)$ and u and v are non-adjacent.
- 5** $G + uv$ is Hamiltonian.
- 6** Every Hamilton cycle of $G + uv$ contains the edge uv .
- 7** Let $P : u = v_1 - v_2 - \cdots - v_n = v$ be a Hamilton Path in G .



Dirac's Theorem

Theorem

If G is a simple graph with $n (\geq 3)$ vertices and $\delta(G) \geq \frac{n}{2}$ then G is Hamiltonian.

Proof.

- 1 Proof by Contradiction.
- 2 Let G be such that, G is maximal non-Hamiltonian and $\delta(G) \geq \frac{n}{2}$.
- 3 G is not complete as $n \geq 3$.
- 4 Choose $u, v \in V(G)$ and u and v are non-adjacent.
- 5 $G + uv$ is Hamiltonian.
- 6 Every Hamilton cycle of $G + uv$ contains the edge uv .
- 7 Let $P : u = v_1 - v_2 - \cdots - v_n = v$ be a Hamilton Path in G .



Dirac's Theorem

Theorem

If G is a simple graph with $n (\geq 3)$ vertices and $\delta(G) \geq \frac{n}{2}$ then G is Hamiltonian.

Proof.

- 1 Proof by Contradiction.
- 2 Let G be such that, G is maximal non-Hamiltonian and $\delta(G) \geq \frac{n}{2}$.
- 3 G is not complete as $n \geq 3$.
- 4 Choose $u, v \in V(G)$ and u and v are non-adjacent.
- 5 $G + uv$ is Hamiltonian.
- 6 Every Hamilton cycle of $G + uv$ contains the edge uv .
- 7 Let $P : u = v_1 - v_2 - \cdots - v_n = v$ be a Hamilton Path in G .



Dirac's Theorem

Theorem

If G is a simple graph with $n (\geq 3)$ vertices and $\delta(G) \geq \frac{n}{2}$ then G is Hamiltonian.

Proof.

- 1 Proof by Contradiction.
- 2 Let G be such that, G is maximal non-Hamiltonian and $\delta(G) \geq \frac{n}{2}$.
- 3 G is not complete as $n \geq 3$.
- 4 Choose $u, v \in V(G)$ and u and v are non-adjacent.
- 5 $G + uv$ is Hamiltonian.
- 6 Every Hamilton cycle of $G + uv$ contains the edge uv .
- 7 Let $P : u = v_1 - v_2 - \cdots - v_n = v$ be a Hamilton Path in G .



Dirac's Theorem

Theorem

If G is a simple graph with n (≥ 3) vertices and $\delta(G) \geq \frac{n}{2}$ then G is Hamiltonian.

Proof.

- 1 Proof by Contradiction.
- 2 Let G be such that, G is maximal non-Hamiltonian and $\delta(G) \geq \frac{n}{2}$.
- 3 G is not complete as $n \geq 3$.
- 4 Choose $u, v \in V(G)$ and u and v are non-adjacent.
- 5 $G + uv$ is Hamiltonian.
- 6 Every Hamilton cycle of $G + uv$ contains the edge uv .
- 7 Let $P : u = v_1 - v_2 - \cdots - v_n = v$ be a Hamilton Path in G .



Proof Contd.

Proof.

Define

$$S = \{v_i : uv_{i+1} \in E(G)\}$$
$$T = \{v_i : v_iv \in E(G)\}$$



- 1 $v_n \notin S \cup T$.
- 2 $|S \cup T| < n$
- 3 $S \cap T = \emptyset$.
- 4 If not, let $v_j \in S \cap T$, then

$$v_1 v_2 \cdots v_j v_n v_{n-1} \cdots v_{j+1} v_1$$

is a Hamilton cycle of G , contrary to our assumption.

- 5 $\deg(u) + \deg(v) = |S| + |T| = |S \cup T| - |S \cap T| < n$.
- 6 A contradiction since $\delta(G) \geq \frac{n}{2}$.



Proof.

Define

$$S = \{v_i : uv_{i+1} \in E(G)\}$$
$$T = \{v_i : v_iv \in E(G)\}$$



- 1 $v_n \notin S \cup T$.
- 2 $|S \cup T| < n$
- 3 $S \cap T = \emptyset$.
- 4 If not, let $v_j \in S \cap T$, then

$$v_1 v_2 \cdots v_j v_n v_{n-1} \cdots v_{j+1} v_1$$

is a Hamilton cycle of G , contrary to our assumption.

- 5 $\deg(u) + \deg(v) = |S| + |T| = |S \cup T| - |S \cap T| < n$.
- 6 A contradiction since $\delta(G) \geq \frac{n}{2}$.



Proof Contd.

Proof.

Define

$$S = \{v_i : uv_{i+1} \in E(G)\}$$
$$T = \{v_i : v_iv \in E(G)\}$$



1 $v_n \notin S \cup T.$

2 $|S \cup T| < n$

3 $S \cap T = \emptyset.$

4 If not, let $v_j \in S \cap T$, then

$$v_1 v_2 \cdots v_j v_n v_{n-1} \cdots v_{j+1} v_1$$

is a Hamilton cycle of G , contrary to our assumption.

5 $\deg(u) + \deg(v) = |S| + |T| = |S \cup T| - |S \cap T| < n.$

6 A contradiction since $\delta(G) \geq \frac{n}{2}.$



Proof.

Define

$$S = \{v_i : uv_{i+1} \in E(G)\}$$
$$T = \{v_i : v_iv \in E(G)\}$$



1 $v_n \notin S \cup T.$

2 $|S \cup T| < n$

3 $S \cap T = \emptyset.$

4 If not, let $v_j \in S \cap T$, then

$$v_1 v_2 \cdots v_j v_n v_{n-1} \cdots v_{j+1} v_1$$

is a Hamilton cycle of G , contrary to our assumption.

5 $\deg(u) + \deg(v) = |S| + |T| = |S \cup T| - |S \cap T| < n.$

6 A contradiction since $\delta(G) \geq \frac{n}{2}.$



Proof Contd.

Proof.

Define

$$S = \{v_i : uv_{i+1} \in E(G)\}$$
$$T = \{v_i : v_iv \in E(G)\}$$



1 $v_n \notin S \cup T.$

2 $|S \cup T| < n$

3 $S \cap T = \emptyset.$

4 If not, let $v_j \in S \cap T$, then

$$v_1 v_2 \cdots v_j v_n v_{n-1} \cdots v_{j+1} v_1$$

is a Hamilton cycle of G , contrary to our assumption.

5 $\deg(u) + \deg(v) = |S| + |T| = |S \cup T| - |S \cap T| < n.$

6 A contradiction since $\delta(G) \geq \frac{n}{2}.$



Proof.

Define

$$S = \{v_i : uv_{i+1} \in E(G)\}$$
$$T = \{v_i : v_iv \in E(G)\}$$



- 1 $v_n \notin S \cup T$.
- 2 $|S \cup T| < n$
- 3 $S \cap T = \emptyset$.
- 4 If not, let $v_j \in S \cap T$, then

$$v_1 v_2 \cdots v_j v_n v_{n-1} \cdots v_{j+1} v_1$$

is a Hamilton cycle of G , contrary to our assumption.

- 5 $\deg(u) + \deg(v) = |S| + |T| = |S \cup T| - |S \cap T| < n$.
- 6 A contradiction since $\delta(G) \geq \frac{n}{2}$.



Proof.

Define

$$S = \{v_i : uv_{i+1} \in E(G)\}$$
$$T = \{v_i : v_iv \in E(G)\}$$



- 1 $v_n \notin S \cup T$.
- 2 $|S \cup T| < n$
- 3 $S \cap T = \emptyset$.
- 4 If not, let $v_j \in S \cap T$, then

$$v_1 v_2 \cdots v_j v_n v_{n-1} \cdots v_{j+1} v_1$$

is a Hamilton cycle of G , contrary to our assumption.

- 5 $\deg(u) + \deg(v) = |S| + |T| = |S \cup T| - |S \cap T| < n$.
- 6 A contradiction since $\delta(G) \geq \frac{n}{2}$.



Proof.

Define

$$S = \{v_i : uv_{i+1} \in E(G)\}$$
$$T = \{v_i : v_iv \in E(G)\}$$



- 1 $v_n \notin S \cup T$.
- 2 $|S \cup T| < n$
- 3 $S \cap T = \emptyset$.
- 4 If not, let $v_j \in S \cap T$, then

$$v_1 v_2 \cdots v_j v_n v_{n-1} \cdots v_{j+1} v_1$$

is a Hamilton cycle of G , contrary to our assumption.

- 5 $\deg(u) + \deg(v) = |S| + |T| = |S \cup T| - |S \cap T| < n$.
- 6 A contradiction since $\delta(G) \geq \frac{n}{2}$.



Theorem

If a graph G on n (≥ 3) is such that

$$\deg(u) + \deg(v) \geq n$$

for every pair of non-adjacent vertices u and v then G is Hamiltonian.

Note : We did not prove this in class. We only proved the consequence of it. (only the consequence in your text book :)).

Consequence of Ore's theorem

Theorem

If a graph G on n (≥ 3) is such that

$$\deg(u) + \deg(v) \geq n$$

for a pair of non-adjacent vertices u and v then G is Hamiltonian iff $G + uv$ is Hamiltonian.

Proof.

- 1 Let $u, v \in G$ be such that $\deg_G(u) + \deg_G(v) \geq n$.
- 2 If $uv \in G$, then there is nothing to prove.
- 3 Assume thus that $uv \notin G$.



Consequence of Ore's theorem

Theorem

If a graph G on n (≥ 3) is such that

$$\deg(u) + \deg(v) \geq n$$

for a pair of non-adjacent vertices u and v then G is Hamiltonian iff $G + uv$ is Hamiltonian.

Proof.

- 1 Let $u, v \in G$ be such that $\deg_G(u) + \deg_G(v) \geq n$.
- 2 If $uv \in G$, then there is nothing to prove.
- 3 Assume thus that $uv \notin G$.



Consequence of Ore's theorem

Theorem

If a graph G on n (≥ 3) is such that

$$\deg(u) + \deg(v) \geq n$$

for a pair of non-adjacent vertices u and v then G is Hamiltonian iff $G + uv$ is Hamiltonian.

Proof.

- 1** Let $u, v \in G$ be such that $\deg_G(u) + \deg_G(v) \geq n$.
- 2** If $uv \in G$, then there is nothing to prove.
- 3** Assume thus that $uv \notin G$.



Consequence of Ore's theorem

Theorem

If a graph G on n (≥ 3) is such that

$$\deg(u) + \deg(v) \geq n$$

for a pair of non-adjacent vertices u and v then G is Hamiltonian iff $G + uv$ is Hamiltonian.

Proof.

- 1** Let $u, v \in G$ be such that $\deg_G(u) + \deg_G(v) \geq n$.
- 2** If $uv \in G$, then there is nothing to prove.
- 3** Assume thus that $uv \notin G$.



Consequence of Ore's theorem

Theorem

If a graph G on n (≥ 3) is such that

$$\deg(u) + \deg(v) \geq n$$

for a pair of non-adjacent vertices u and v then G is Hamiltonian iff $G + uv$ is Hamiltonian.

Proof.

- 1** Let $u, v \in G$ be such that $\deg_G(u) + \deg_G(v) \geq n$.
- 2** If $uv \in G$, then there is nothing to prove.
- 3** Assume thus that $uv \notin G$.



This is trivial.

- 1 Given G is Hamiltonian.
- 2 G has a Hamilton cycle C (say)
- 3 C is also a Hamilton cycle of $G + uv$.
- 4 $\Rightarrow G + uv$ is Hamiltonian.

This is trivial.

- 1** Given G is Hamiltonian.
- 2 G has a Hamilton cycle C (say)
- 3 C is also a Hamilton cycle of $G + uv$.
- 4 $\Rightarrow G + uv$ is Hamiltonian.

This is trivial.

- 1 Given G is Hamiltonian.
- 2 G has a Hamilton cycle C (say)
- 3 C is also a Hamilton cycle of $G + uv$.
- 4 $\Rightarrow G + uv$ is Hamiltonian.

This is trivial.

- 1 Given G is Hamiltonian.
- 2 G has a Hamilton cycle C (say)
- 3 C is also a Hamilton cycle of $G + uv$.
- 4 $\Rightarrow G + uv$ is Hamiltonian.

This is trivial.

- 1 Given G is Hamiltonian.
- 2 G has a Hamilton cycle C (say)
- 3 C is also a Hamilton cycle of $G + uv$.
- 4 $\Rightarrow G + uv$ is Hamiltonian.

This is trivial.

- 1 Given G is Hamiltonian.
- 2 G has a Hamilton cycle C (say)
- 3 C is also a Hamilton cycle of $G + uv$.
- 4 $\Rightarrow G + uv$ is Hamiltonian.

Suppose given that $G + uv$ is Hamiltonian.

We show that G is Hamiltonian.

- 1 Denote $e = uv$.
- 2 $G + e$ has a Hamilton cycle C .
- 3 If C do not contain e , then C is a Hamilton cycle of G .
- 4 If C contains e .
- 5 Let $C = u = v_1 - v_2 - \cdots - v_n = v - u$.
- 6 Then for $P = u = v_1 - v_2 - \cdots - v_n = v$, $C = P + e$.
- 7 P is a Hamilton path of G .

Proof: Converse

Suppose given that $G + uv$ is Hamiltonian.
We show that G is Hamiltonian.

- 1 Denote $e = uv$.
- 2 $G + e$ has a Hamilton cycle C .
- 3 If C do not contain e , then C is a Hamilton cycle of G .
- 4 If C contains e .
- 5 Let $C = u = v_1 - v_2 - \cdots - v_n = v - u$.
- 6 Then for $P = u = v_1 - v_2 - \cdots - v_n = v$, $C = P + e$.
- 7 P is a Hamilton path of G .

Proof: Converse

Suppose given that $G + uv$ is Hamiltonian.
We show that G is Hamiltonian.

- 1 Denote $e = uv$.
- 2 $G + e$ has a Hamilton cycle C .
- 3 If C do not contain e , then C is a Hamilton cycle of G .
- 4 If C contains e .
- 5 Let $C = u = v_1 - v_2 - \cdots - v_n = v - u$.
- 6 Then for $P = u = v_1 - v_2 - \cdots - v_n = v$, $C = P + e$.
- 7 P is a Hamilton path of G .

Proof: Converse

Suppose given that $G + uv$ is Hamiltonian.
We show that G is Hamiltonian.

- 1 Denote $e = uv$.
- 2 $G + e$ has a Hamilton cycle C .
- 3 If C do not contain e , then C is a Hamilton cycle of G .
- 4 If C contains e .
- 5 Let $C = u = v_1 - v_2 - \cdots - v_n = v - u$.
- 6 Then for $P = u = v_1 - v_2 - \cdots - v_n = v$, $C = P + e$.
- 7 P is a Hamilton path of G .

Proof: Converse

Suppose given that $G + uv$ is Hamiltonian.
We show that G is Hamiltonian.

- 1 Denote $e = uv$.
- 2 $G + e$ has a Hamilton cycle C .
- 3 If C do not contain e , then C is a Hamilton cycle of G .
- 4 If C contains e .
- 5 Let $C = u = v_1 - v_2 - \cdots - v_n = v - u$.
- 6 Then for $P = u = v_1 - v_2 - \cdots - v_n = v$, $C = P + e$.
- 7 P is a Hamilton path of G .

Proof: Converse

Suppose given that $G + uv$ is Hamiltonian.
We show that G is Hamiltonian.

- 1 Denote $e = uv$.
- 2 $G + e$ has a Hamilton cycle C .
- 3 If C do not contain e , then C is a Hamilton cycle of G .
- 4 If C contains e .
- 5 Let $C = u = v_1 - v_2 - \cdots - v_n = v - u$.
- 6 Then for $P = u = v_1 - v_2 - \cdots - v_n = v$, $C = P + e$.
- 7 P is a Hamilton path of G .

Proof: Converse

Suppose given that $G + uv$ is Hamiltonian.
We show that G is Hamiltonian.

- 1 Denote $e = uv$.
- 2 $G + e$ has a Hamilton cycle C .
- 3 If C do not contain e , then C is a Hamilton cycle of G .
- 4 If C contains e .
- 5 Let $C = u = v_1 - v_2 - \cdots - v_n = v - u$.
- 6 Then for $P = u = v_1 - v_2 - \cdots - v_n = v$, $C = P + e$.
- 7 P is a Hamilton path of G .

Proof: Converse

Suppose given that $G + uv$ is Hamiltonian.
We show that G is Hamiltonian.

- 1 Denote $e = uv$.
- 2 $G + e$ has a Hamilton cycle C .
- 3 If C do not contain e , then C is a Hamilton cycle of G .
- 4 If C contains e .
- 5 Let $C = u = v_1 - v_2 - \cdots - v_n = v - u$.
- 6 Then for $P = u = v_1 - v_2 - \cdots - v_n = v$, $C = P + e$.
- 7 P is a Hamilton path of G .

Suppose given that $G + uv$ is Hamiltonian.
We show that G is Hamiltonian.

- 1 Denote $e = uv$.
- 2 $G + e$ has a Hamilton cycle C .
- 3 If C do not contain e , then C is a Hamilton cycle of G .
- 4 If C contains e .
- 5 Let $C = u = v_1 - v_2 - \cdots - v_n = v - u$.
- 6 Then for $P = u = v_1 - v_2 - \cdots - v_n = v$, $C = P + e$.
- 7 P is a Hamilton path of G .

Claim : There exists an i with $1 < i < n$ such that $uv_i \in G$ and $v_{i-1}v \in G$.

- 1** u and v not adjacent in G .
- 2** $\deg_G(u) + \deg_G(v) \geq n$.
- 3** $\implies (n-1) - \deg_G(v) \leq \deg_G(u) - 1$.
- 4** Atmost $\deg_G(u) - 1$ vertices other than v are not adjacent to v .
- 5** Let $S = \{v_{i-1} : uv_i \in E(G)\}$ in the Hamilton path P .
- 6** $v \notin S$ and $|S| = \deg_G(u)$.
- 7** Therefore, at least one vertex say v_k of S is adjacent to v_n .

Claim : There exists an i with $1 < i < n$ such that $uv_i \in G$ and $v_{i-1}v \in G$.

- 1** u and v not adjacent in G .
- 2** $\deg_G(u) + \deg_G(v) \geq n$.
- 3** $\implies (n-1) - \deg_G(v) \leq \deg_G(u) - 1$.
- 4** Atmost $\deg_G(u) - 1$ vertices other than v are not adjacent to v .
- 5** Let $S = \{v_{i-1} : uv_i \in E(G)\}$ in the Hamilton path P .
- 6** $v \notin S$ and $|S| = \deg_G(u)$.
- 7** Therefore, at least one vertex say v_k of S is adjacent to v_n .

Claim : There exists an i with $1 < i < n$ such that $uv_i \in G$ and $v_{i-1}v \in G$.

- 1** u and v not adjacent in G .
- 2** $\deg_G(u) + \deg_G(v) \geq n$.
- 3** $\implies (n - 1) - \deg_G(v) \leq \deg_G(u) - 1$.
- 4** Atmost $\deg_G(u) - 1$ vertices other than v are not adjacent to v .
- 5** Let $S = \{v_{i-1} : uv_i \in E(G)\}$ in the Hamilton path P .
- 6** $v \notin S$ and $|S| = \deg_G(u)$.
- 7** Therefore, at least one vertex say v_k of S is adjacent to v_n .

Claim : There exists an i with $1 < i < n$ such that $uv_i \in G$ and $v_{i-1}v \in G$.

- 1 u and v not adjacent in G .
- 2 $\deg_G(u) + \deg_G(v) \geq n$.
- 3 $\implies (n - 1) - \deg_G(v) \leq \deg_G(u) - 1$.
- 4 Atmost $\deg_G(u) - 1$ vertices other than v are not adjacent to v .
- 5 Let $S = \{v_{i-1} : uv_i \in E(G)\}$ in the Hamilton path P .
- 6 $v \notin S$ and $|S| = \deg_G(u)$.
- 7 Therefore, at least one vertex say v_k of S is adjacent to v_n .

Claim : There exists an i with $1 < i < n$ such that $uv_i \in G$ and $v_{i-1}v \in G$.

- 1** u and v not adjacent in G .
- 2** $\deg_G(u) + \deg_G(v) \geq n$.
- 3** $\implies (n-1) - \deg_G(v) \leq \deg_G(u) - 1$.
- 4** Atmost $\deg_G(u) - 1$ vertices other than v are not adjacent to v .
- 5** Let $S = \{v_{i-1} : uv_i \in E(G)\}$ in the Hamilton path P .
- 6** $v \notin S$ and $|S| = \deg_G(u)$.
- 7** Therefore, at least one vertex say v_k of S is adjacent to v_n .

Claim : There exists an i with $1 < i < n$ such that $uv_i \in G$ and $v_{i-1}v \in G$.

- 1** u and v not adjacent in G .
- 2** $\deg_G(u) + \deg_G(v) \geq n$.
- 3** $\implies (n-1) - \deg_G(v) \leq \deg_G(u) - 1$.
- 4** Atmost $\deg_G(u) - 1$ vertices other than v are not adjacent to v .
- 5** Let $S = \{v_{i-1} : uv_i \in E(G)\}$ in the Hamilton path P .
- 6** $v \notin S$ and $|S| = \deg_G(u)$.
- 7** Therefore, at least one vertex say v_k of S is adjacent to v_n .

Claim : There exists an i with $1 < i < n$ such that $uv_i \in G$ and $v_{i-1}v \in G$.

- 1** u and v not adjacent in G .
- 2** $\deg_G(u) + \deg_G(v) \geq n$.
- 3** $\implies (n-1) - \deg_G(v) \leq \deg_G(u) - 1$.
- 4** Atmost $\deg_G(u) - 1$ vertices other than v are not adjacent to v .
- 5** Let $S = \{v_{i-1} : uv_i \in E(G)\}$ in the Hamilton path P .
- 6** $v \notin S$ and $|S| = \deg_G(u)$.
- 7** Therefore, at least one vertex say v_k of S is adjacent to v_n .

1 By the previous claim, there exists v_k such that $e_1 = uv_{k+1}, e_2 = v_kv \in E(G)$.

2 Then, from the Hamilton path

$$P = u = v_1 - v - 2 - \cdots - v_k - v_{k+1} - \cdots - v_n = v$$

, we construct a Hamilton Cycle.

3 $C = u = v_1 - v_2 - \cdots - v_k - v = v_n - v_{n-1} - \cdots - v_{k+1} - v_1 = u$ is a Hamilton cycle in G . \square

- 1 By the previous claim, there exists v_k such that $e_1 = uv_{k+1}, e_2 = v_kv \in E(G)$.
- 2 Then, from the Hamilton path

$$P = u = v_1 - v - 2 - \cdots - v_k - v_{k+1} - \cdots - v_n = v$$

, we construct a Hamilton Cycle.

- 3 $C = u = v_1 - v_2 - \cdots - v_k - v = v_n - v_{n-1} - \cdots - v_{k+1} - v_1 = u$ is a Hamilton cycle in G . □

- 1 By the previous claim, there exists v_k such that $e_1 = uv_{k+1}, e_2 = v_kv \in E(G)$.
- 2 Then, from the Hamilton path

$$P = u = v_1 - v - 2 - \cdots - v_k - v_{k+1} - \cdots - v_n = v$$

, we construct a Hamilton Cycle.

- 3 $C = u = v_1 - v_2 - \cdots - v_k - v = v_n - v_{n-1} - \cdots - v_{k+1} - v_1 = u$ is a Hamilton cycle in G . □

Definition

For a graph G , define inductively a sequence G_0, G_1, \dots, G_k of graphs such that

$$G_0 = G \text{ and } G_{i+1} = G_i + uv$$

where u and v are any vertices such that

- 1** $uv \notin G_i$ and
- 2** $\deg_{G_i}(u) + \deg_{G_i}(v) \geq n$.

- 1 This procedure stops when no new edges can be added to G_k for some k , that is, in G_k , for all $u, v \in G$ either $uv \in G_k$ or $\deg_{G_k}(u) + \deg_{G_k}(v) < n$.
- 2 The result of this procedure is the closure of G , and it is denoted by $cl(G)(= G_k)$.
- 3 In each step of the construction of $cl(G)$ there are usually alternatives which edge uv is to be added to the graph, and therefore the above procedure is not deterministic.
- 4 However, the final result $cl(G)$ is independent of the choices.
- 5 Example

- 1 This procedure stops when no new edges can be added to G_k for some k , that is, in G_k , for all $u, v \in G$ either $uv \in G_k$ or $\deg_{G_k}(u) + \deg_{G_k}(v) < n$.
- 2 The result of this procedure is the closure of G , and it is denoted by $cl(G)(= G_k)$.
- 3 In each step of the construction of $cl(G)$ there are usually alternatives which edge uv is to be added to the graph, and therefore the above procedure is not deterministic.
- 4 However, the final result $cl(G)$ is independent of the choices.
- 5 Example

- 1 This procedure stops when no new edges can be added to G_k for some k , that is, in G_k , for all $u, v \in G$ either $uv \in G_k$ or $\deg_{G_k}(u) + \deg_{G_k}(v) < n$.
- 2 The result of this procedure is the closure of G , and it is denoted by $cl(G)(= G_k)$.
- 3 In each step of the construction of $cl(G)$ there are usually alternatives which edge uv is to be added to the graph, and therefore the above procedure is not deterministic.
- 4 However, the final result $cl(G)$ is independent of the choices.
- 5 Example

- 1 This procedure stops when no new edges can be added to G_k for some k , that is, in G_k , for all $u, v \in G$ either $uv \in G_k$ or $\deg_{G_k}(u) + \deg_{G_k}(v) < n$.
- 2 The result of this procedure is the closure of G , and it is denoted by $cl(G)(= G_k)$.
- 3 In each step of the construction of $cl(G)$ there are usually alternatives which edge uv is to be added to the graph, and therefore the above procedure is not deterministic.
- 4 However, the final result $cl(G)$ is independent of the choices.

5 Example

- 1 This procedure stops when no new edges can be added to G_k for some k , that is, in G_k , for all $u, v \in G$ either $uv \in G_k$ or $\deg_{G_k}(u) + \deg_{G_k}(v) < n$.
- 2 The result of this procedure is the closure of G , and it is denoted by $cl(G)(= G_k)$.
- 3 In each step of the construction of $cl(G)$ there are usually alternatives which edge uv is to be added to the graph, and therefore the above procedure is not deterministic.
- 4 However, the final result $cl(G)$ is independent of the choices.
- 5 Example

Lemma

The closure $cl(G)$ is uniquely defined for all graphs G of order $n \geq 3$.

Proof.

Suppose there are two ways to close G , say

$$H = G + \{e_1, \dots, e_r\} \text{ and } H' = G + \{f_1, \dots, f_s\}$$

where the edges are added in the given orders.

We show that

$$\{e_1, \dots, e_r\} = \{f_1, \dots, f_s\}$$



Lemma

The closure $cl(G)$ is uniquely defined for all graphs G of order $n \geq 3$.

Proof.

Suppose there are two ways to close G , say

$$H = G + \{e_1, \dots, e_r\} \text{ and } H' = G + \{f_1, \dots, f_s\}$$

where the edges are added in the given orders.

We show that

$$\{e_1, \dots, e_r\} = \{f_1, \dots, f_s\}$$



Lemma

The closure $cl(G)$ is uniquely defined for all graphs G of order $n \geq 3$.

Proof.

Suppose there are two ways to close G , say

$$H = G + \{e_1, \dots, e_r\} \text{ and } H' = G + \{f_1, \dots, f_s\}$$

where the edges are added in the given orders.

We show that

$$\{e_1, \dots, e_r\} = \{f_1, \dots, f_s\}$$



Proof Contd.

- 1 Let $H_i = G + \{e_1, \dots, e_i\}$ and $H'_i = G + \{f_1, \dots, f_i\}$.
- 2 For the initial values, we have $G = H_0 = H'_0$.
- 3 Let $e_k = uv \in H_k$ be the first edge such that $e_k \notin H'$
- 4 Now, $e_k \notin H_{k-1}$ we have

$$\deg_{H_{k-1}}(u) + \deg_{H_{k-1}}(v) \geq n,$$

- 5 $H_{k-1} \subseteq H'$ (Subgraph).
- 6 $\deg_{H'}(u) + \deg_{H'}(v) \geq n$, which means that $e = uv$ must be in H' ; a contradiction.
- 7 Therefore $H \subseteq H'$.
- 8 Similarly we can show that $H \subseteq H'$.
- 9 Hence, $H = H'$. □

Proof Contd.

- 1 Let $H_i = G + \{e_1, \dots, e_i\}$ and $H'_i = G + \{f_1, \dots, f_i\}$.
- 2 For the initial values, we have $G = H_0 = H'_0$.
- 3 Let $e_k = uv \in H_k$ be the first edge such that $e_k \notin H'$
- 4 Now, $e_k \notin H_{k-1}$ we have

$$\deg_{H_{k-1}}(u) + \deg_{H_{k-1}}(v) \geq n,$$

- 5 $H_{k-1} \subseteq H'$ (Subgraph).
- 6 $\deg_{H'}(u) + \deg_{H'}(v) \geq n$, which means that $e = uv$ must be in H' ; a contradiction.
- 7 Therefore $H \subseteq H'$.
- 8 Similarly we can show that $H \subseteq H'$.
- 9 Hence, $H = H'$. □

Proof Contd.

- 1 Let $H_i = G + \{e_1, \dots, e_i\}$ and $H'_i = G + \{f_1, \dots, f_i\}$.
- 2 For the initial values, we have $G = H_0 = H'_0$.
- 3 Let $e_k = uv \in H_k$ be the first edge such that $e_k \notin H'$
- 4 Now, $e_k \notin H_{k-1}$ we have

$$\deg_{H_{k-1}}(u) + \deg_{H_{k-1}}(v) \geq n,$$

- 5 $H_{k-1} \subseteq H'$ (Subgraph).
- 6 $\deg_{H'}(u) + \deg_{H'}(v) \geq n$, which means that $e = uv$ must be in H' ; a contradiction.
- 7 Therefore $H \subseteq H'$.
- 8 Similarly we can show that $H \subseteq H'$.
- 9 Hence, $H = H'$. □

- 1 Let $H_i = G + \{e_1, \dots, e_i\}$ and $H'_i = G + \{f_1, \dots, f_i\}$.
- 2 For the initial values, we have $G = H_0 = H'_0$.
- 3 Let $e_k = uv \in H_k$ be the first edge such that $e_k \notin H'$
- 4 Now, $e_k \notin H_{k-1}$ we have

$$\deg_{H_{k-1}}(u) + \deg_{H_{k-1}}(v) \geq n,$$

- 5 $H_{k-1} \subseteq H'$ (Subgraph).
- 6 $\deg_{H'}(u) + \deg_{H'}(v) \geq n$, which means that $e = uv$ must be in H' ; a contradiction.
- 7 Therefore $H \subseteq H'$.
- 8 Similarly we can show that $H \subseteq H'$.
- 9 Hence, $H = H'$. □

Proof Contd.

- 1 Let $H_i = G + \{e_1, \dots, e_i\}$ and $H'_i = G + \{f_1, \dots, f_i\}$.
- 2 For the initial values, we have $G = H_0 = H'_0$.
- 3 Let $e_k = uv \in H_k$ be the first edge such that $e_k \notin H'$
- 4 Now, $e_k \notin H_{k-1}$ we have

$$\deg_{H_{k-1}}(u) + \deg_{H_{k-1}}(v) \geq n,$$

- 5 $H_{k-1} \subseteq H'$ (Subgraph).
- 6 $\deg_{H'}(u) + \deg_{H'}(v) \geq n$, which means that $e = uv$ must be in H' ; a contradiction.
- 7 Therefore $H \subseteq H'$.
- 8 Similarly we can show that $H \subseteq H'$.
- 9 Hence, $H = H'$. □

- 1 Let $H_i = G + \{e_1, \dots, e_i\}$ and $H'_i = G + \{f_1, \dots, f_i\}$.
- 2 For the initial values, we have $G = H_0 = H'_0$.
- 3 Let $e_k = uv \in H_k$ be the first edge such that $e_k \notin H'$
- 4 Now, $e_k \notin H_{k-1}$ we have

$$\deg_{H_{k-1}}(u) + \deg_{H_{k-1}}(v) \geq n,$$

- 5 $H_{k-1} \subseteq H'$ (Subgraph).
- 6 $\deg_{H'}(u) + \deg_{H'}(v) \geq n$, which means that $e = uv$ must be in H' ; a contradiction.
- 7 Therefore $H \subseteq H'$.
- 8 Similarly we can show that $H \subseteq H'$.
- 9 Hence, $H = H'$. □

Proof Contd.

- 1 Let $H_i = G + \{e_1, \dots, e_i\}$ and $H'_i = G + \{f_1, \dots, f_i\}$.
- 2 For the initial values, we have $G = H_0 = H'_0$.
- 3 Let $e_k = uv \in H_k$ be the first edge such that $e_k \notin H'$
- 4 Now, $e_k \notin H_{k-1}$ we have

$$\deg_{H_{k-1}}(u) + \deg_{H_{k-1}}(v) \geq n,$$

- 5 $H_{k-1} \subseteq H'$ (Subgraph).
- 6 $\deg_{H'}(u) + \deg_{H'}(v) \geq n$, which means that $e = uv$ must be in H' ; a contradiction.
- 7 Therefore $H \subseteq H'$.
- 8 Similarly we can show that $H \subseteq H'$.
- 9 Hence, $H = H'$. □

- 1 Let $H_i = G + \{e_1, \dots, e_i\}$ and $H'_i = G + \{f_1, \dots, f_i\}$.
- 2 For the initial values, we have $G = H_0 = H'_0$.
- 3 Let $e_k = uv \in H_k$ be the first edge such that $e_k \notin H'$
- 4 Now, $e_k \notin H_{k-1}$ we have

$$\deg_{H_{k-1}}(u) + \deg_{H_{k-1}}(v) \geq n,$$

- 5 $H_{k-1} \subseteq H'$ (Subgraph).
- 6 $\deg_{H'}(u) + \deg_{H'}(v) \geq n$, which means that $e = uv$ must be in H' ; a contradiction.
- 7 Therefore $H \subseteq H'$.
- 8 Similarly we can show that $H \subseteq H'$.
- 9 Hence, $H = H'$. □

- 1 Let $H_i = G + \{e_1, \dots, e_i\}$ and $H'_i = G + \{f_1, \dots, f_i\}$.
- 2 For the initial values, we have $G = H_0 = H'_0$.
- 3 Let $e_k = uv \in H_k$ be the first edge such that $e_k \notin H'$
- 4 Now, $e_k \notin H_{k-1}$ we have

$$\deg_{H_{k-1}}(u) + \deg_{H_{k-1}}(v) \geq n,$$

- 5 $H_{k-1} \subseteq H'$ (Subgraph).
- 6 $\deg_{H'}(u) + \deg_{H'}(v) \geq n$, which means that $e = uv$ must be in H' ; a contradiction.
- 7 Therefore $H \subseteq H'$.
- 8 Similarly we can show that $H \subseteq H'$.
- 9 Hence, $H = H'$. □

Theorem

Theorem

Let G be a graph such that $|V(G)| = n \geq 3$.

- 1 G is Hamiltonian if and only if its closure $cl(G)$ is Hamiltonian.
- 2 If $cl(G)$ is a complete graph, then G is Hamiltonian.

Proof.

Exercise. □