

Coloring

October 17, 2024

Motivation :

- 1** The concept is useful to model many scheduling problems.
- 2** It also arises in many circuit board problems where the wires connecting a device have to be of different color.

Motivation :

- 1** The concept is useful to model many scheduling problems.
- 2** It also arises in many circuit board problems where the wires connecting a device have to be of different color.

Definition

- 1 A **k -vertex colouring** of G is an assignment of k colours, $1, 2, \dots, k$, to the vertices of G .
- 2 The colouring is proper if no two distinct adjacent vertices have the same colour.
- 3 A proper **k -vertex colouring** of a loopless graph G is a partition (V_1, V_2, \dots, V_t) of V into k (possibly empty) independent sets.
- 4 G is **k -vertex-colourable** if G has a proper k -vertex colouring.

Definition

- 1 A **k -vertex colouring** of G is an assignment of k colours, $1, 2, \dots, k$, to the vertices of G .
- 2 The colouring is proper if no two distinct adjacent vertices have the same colour.
- 3 A proper k -vertex colouring of a loopless graph G is a partition (V_1, V_2, \dots, V_t) of V into k (possibly empty) independent sets.
- 4 G is k -vertex-colourable if G has a proper k -vertex colouring.

Definition

- 1 A **k -vertex colouring** of G is an assignment of k colours, $1, 2, \dots, k$, to the vertices of G .
- 2 The colouring is proper if no two distinct adjacent vertices have the same colour.
- 3 A **proper k -vertex colouring** of a loopless graph G is a partition (V_1, V_2, \dots, V_t) of V into k (possibly empty) independent sets.
- 4 G is **k -vertex-colourable** if G has a proper k -vertex colouring.

Definition

- 1 A **k -vertex colouring** of G is an assignment of k colours, $1, 2, \dots, k$, to the vertices of G .
- 2 The colouring is proper if no two distinct adjacent vertices have the same colour.
- 3 A **proper k -vertex colouring** of a loopless graph G is a partition (V_1, V_2, \dots, V_t) of V into k (possibly empty) independent sets.
- 4 G is **k -vertex-colourable** if G has a proper k -vertex colouring.

- 1 A simple graph is 1-colourable if and only if it is empty (No vertices or totally disconnected).
- 2 A simple graph 2-colourable if and only if it is bipartite.
- 3 The chromatic number, $\chi(G)$, of G is the minimum k for which G is k -colourable.
- 4 If $\chi(G) = k$, G is said to be k -chromatic.

- 1 A simple graph is 1-colourable if and only if it is empty (No vertices or totally disconnected).
- 2 A simple graph 2-colourable if and only if it is bipartite.
- 3 The chromatic number, $\chi(G)$, of G is the minimum k for which G is k -colourable.
- 4 If $\chi(G) = k$, G is said to be k -chromatic.

- 1 A simple graph is 1-colourable if and only if it is empty (No vertices or totally disconnected).
- 2 A simple graph 2-colourable if and only if it is bipartite.
- 3 The chromatic number, $\chi(G)$, of G is the minimum k for which G is k -colourable.
- 4 If $\chi(G) = k$, G is said to be k -chromatic.

- 1 A simple graph is 1-colourable if and only if it is empty (No vertices or totally disconnected).
- 2 A simple graph 2-colourable if and only if it is bipartite.
- 3 The chromatic number, $\chi(G)$, of G is the minimum k for which G is k -colourable.
- 4 If $\chi(G) = k$, G is said to be **k -chromatic**.

- 1 We refer to a 'proper vertex colouring' as, simply, a colouring.
- 2 We refer to a 'proper k -vertex colouring' as a k -colouring.
- 3 we shall similarly abbreviate ' k -vertex-colourable' to k -colourable.
- 4 Clearly, a graph is k -colourable if and only if its underlying simple graph is k -colourable.
- 5 Therefore, in discussing colourings, we shall restrict ourselves to simple graphs.

- 1 We refer to a 'proper vertex colouring' as, simply, a colouring.
- 2 We refer to a 'proper k -vertex colouring' as a k -colouring.
- 3 we shall similarly abbreviate ' k -vertex-colourable' to k -colourable.
- 4 Clearly, a graph is k -colourable if and only if its underlying simple graph is k -colourable.
- 5 Therefore, in discussing colourings, we shall restrict ourselves to simple graphs.

- 1 We refer to a 'proper vertex colouring' as, simply, a colouring.
- 2 We refer to a 'proper k -vertex colouring' as a k -colouring.
- 3 we shall similarly abbreviate ' k -vertex-colourable' to k -colourable.
- 4 Clearly, a graph is k -colourable if and only if its underlying simple graph is k -colourable.
- 5 Therefore, in discussing colourings, we shall restrict ourselves to simple graphs.

- 1 We refer to a 'proper vertex colouring' as, simply, a colouring.
- 2 We refer to a 'proper k -vertex colouring' as a k -colouring.
- 3 we shall similarly abbreviate ' k -vertex-colourable' to k -colourable.
- 4 Clearly, a graph is k -colourable if and only if its underlying simple graph is k -colourable.
- 5 Therefore, in discussing colourings, we shall restrict ourselves to simple graphs.

- 1 We refer to a 'proper vertex colouring' as, simply, a colouring.
- 2 We refer to a 'proper k -vertex colouring' as a k -colouring.
- 3 we shall similarly abbreviate ' k -vertex-colourable' to k -colourable.
- 4 Clearly, a graph is k -colourable if and only if its underlying simple graph is k -colourable.
- 5 Therefore, in discussing colourings, we shall restrict ourselves to simple graphs.

Example




Figure: Petersen




Figure: Proper coloring

We study the properties of a special class of graphs called **critical graphs**. (Introduced by Dirac in 1952).

Definition

- 1 We say that a graph G is critical if $\chi(H) < \chi(G)$ for every proper subgraph H of G .
- 2 A k -critical graph is one that is k -chromatic and critical.
- 3 Every k -chromatic graph has a k -critical subgraph.

We study the properties of a special class of graphs called **critical graphs**. (Introduced by Dirac in 1952).

Definition

- 1** We say that a graph G is critical if $\chi(H) < \chi(G)$ for every proper subgraph H of G .
- 2** A k -critical graph is one that is k -chromatic and critical.
- 3** Every k -chromatic graph has a k -critical subgraph.

We study the properties of a special class of graphs called **critical graphs**. (Introduced by Dirac in 1952).

Definition

- 1** We say that a graph G is critical if $\chi(H) < \chi(G)$ for every proper subgraph H of G .
- 2** A k -critical graph is one that is k -chromatic and critical.
- 3** Every k -chromatic graph has a k -critical subgraph.

We study the properties of a special class of graphs called **critical graphs**. (Introduced by Dirac in 1952).

Definition

- 1** We say that a graph G is critical if $\chi(H) < \chi(G)$ for every proper subgraph H of G .
- 2** A k -critical graph is one that is k -chromatic and critical.
- 3** Every k -chromatic graph has a k -critical subgraph.

- 1 A 1-critical graph has chromatic number 1, so must be an empty graph E_n .
- 2 Which n ? If $n > 1$, then on the removal of any vertex, we still have an empty graph with chromatic number 1, and so the graph is not 1-critical.
- 3 But if $n = 1$, when we remove the only vertex we get a graph which has no vertices, and so has chromatic number 0.
- 4 So the only 1-critical graph is $E_1 = K_1$.

- 1 A 1-critical graph has chromatic number 1, so must be an empty graph E_n .
- 2 Which n ? If $n > 1$, then on the removal of any vertex, we still have an empty graph with chromatic number 1, and so the graph is not 1-critical.
- 3 But if $n = 1$, when we remove the only vertex we get a graph which has no vertices, and so has chromatic number 0.
- 4 So the only 1-critical graph is $E_1 = K_1$.

- 1 A 1-critical graph has chromatic number 1, so must be an empty graph E_n .
- 2 Which n ? If $n > 1$, then on the removal of any vertex, we still have an empty graph with chromatic number 1, and so the graph is not 1-critical.
- 3 But if $n = 1$, when we remove the only vertex we get a graph which has no vertices, and so has chromatic number 0.
- 4 So the only 1-critical graph is $E_1 = K_1$.

- 1 A 1-critical graph has chromatic number 1, so must be an empty graph E_n .
- 2 Which n ? If $n > 1$, then on the removal of any vertex, we still have an empty graph with chromatic number 1, and so the graph is not 1-critical.
- 3 But if $n = 1$, when we remove the only vertex we get a graph which has no vertices, and so has chromatic number 0.
- 4 So the only 1-critical graph is $E_1 = K_1$.

2-critical graphs

- 1** A 2-critical graph has chromatic number 2.
- 2 so it must be a bipartite graph with at least one edge.
- 3 On deleting any vertex, we must have an empty graph (the only graphs with chromatic number 1).
- 4 So every vertex must be adjacent to every edge.
- 5 The only graph with this property is K_2 .
- 6 so K_2 is the only 2-critical graph.

Exercise : Classify all 3-critical graphs.

2-critical graphs

- 1 A 2-critical graph has chromatic number 2.
- 2 so it must be a bipartite graph with at least one edge.
- 3 On deleting any vertex, we must have an empty graph (the only graphs with chromatic number 1).
- 4 So every vertex must be adjacent to every edge.
- 5 The only graph with this property is K_2 .
- 6 so K_2 is the only 2-critical graph.

Exercise : Classify all 3-critical graphs.

2-critical graphs

- 1 A 2-critical graph has chromatic number 2.
- 2 so it must be a bipartite graph with at least one edge.
- 3 On deleting any vertex, we must have an empty graph (the only graphs with chromatic number 1).
- 4 So every vertex must be adjacent to every edge.
- 5 The only graph with this property is K_2 .
- 6 so K_2 is the only 2-critical graph.

Exercise : Classify all 3-critical graphs.

2-critical graphs

- 1 A 2-critical graph has chromatic number 2.
- 2 so it must be a bipartite graph with at least one edge.
- 3 On deleting any vertex, we must have an empty graph (the only graphs with chromatic number 1).
- 4 So every vertex must be adjacent to every edge.
- 5 The only graph with this property is K_2 .
- 6 so K_2 is the only 2-critical graph.

Exercise : Classify all 3-critical graphs.

2-critical graphs

- 1 A 2-critical graph has chromatic number 2.
- 2 so it must be a bipartite graph with at least one edge.
- 3 On deleting any vertex, we must have an empty graph (the only graphs with chromatic number 1).
- 4 So every vertex must be adjacent to every edge.
- 5 The only graph with this property is K_2 .
- 6 so K_2 is the only 2-critical graph.


Exercise : Classify all 3-critical graphs.

2-critical graphs

- 1** A 2-critical graph has chromatic number 2.
- 2** so it must be a bipartite graph with at least one edge.
- 3** On deleting any vertex, we must have an empty graph (the only graphs with chromatic number 1).
- 4** So every vertex must be adjacent to every edge.
- 5** The only graph with this property is K_2 .
- 6** so K_2 is the only 2-critical graph.


Exercise : Classify all 3-critical graphs.

Examples




- 1 K_3 is 3-critical.
- 2 K_4 - bipartite, hence 2-chromatic, but k -critical for $k = ?$.
- 3 $K - 5$ - not bipartite..3-chromatic, k -critical, $k = ?$

Examples




- 1** K_3 is 3-critical.
- 2** K_4 - bipartite, hence 2-chromatic, but k -critical for $k = ?$.
- 3** K_5 - not bipartite..3-chromatic, k -critical, $k = ?$

Examples



- 1** K_3 is 3-critical.
- 2** K_4 - bipartite, hence 2-chromatic, but k -critical for $k = ?$.
- 3** K_5 - not bipartite..3-chromatic, k -critical, $k = ?$

Examples



- 1** K_3 is 3-critical.
- 2** K_4 - bipartite, hence 2-chromatic, but k -critical for $k = ?$.
- 3** $K - 5$ - not bipartite..3-chromatic, k -critical, $k = ?$

Properties

Lemma

Every k -critical graph is connected.

Proof.

Easy by definition.



Properties

Lemma

Every k -critical graph is connected.

Proof.

Easy by definition.



Lemma

If G is k -critical, then $\delta \geq k - 1$.

Proof.

- 1 By contradiction.
- 2 Suppose $\delta(G) \leq k - 2$.
- 3 Let v be a vertex of degree δ in G .
- 4 Since G is k -critical, $G - v$ is $(k - 1)$ -colourable.
- 5 Let $(V_1, V_2, \dots, V_{k-1})$ be a $(k - 1)$ -colouring of $G - v$.
- 6 By definition, v is adjacent in G to $\delta < k - 1$ vertices.



Lemma

If G is k -critical, then $\delta \geq k - 1$.

Proof.

- 1** By contradiction.
- 2 Suppose $\delta(G) \leq k - 2$.
- 3 Let v be a vertex of degree δ in G .
- 4 Since G is k -critical, $G - v$ is $(k - 1)$ -colourable.
- 5 Let $(V_1, V_2, \dots, V_{k-1})$ be a $(k - 1)$ -colouring of $G - v$.
- 6 By definition, v is adjacent in G to $\delta < k - 1$ vertices.



Lemma

If G is k -critical, then $\delta \geq k - 1$.

Proof.

- 1 By contradiction.
- 2 Suppose $\delta(G) \leq k - 2$.
- 3 Let v be a vertex of degree δ in G .
- 4 Since G is k -critical, $G - v$ is $(k - 1)$ -colourable.
- 5 Let $(V_1, V_2, \dots, V_{k-1})$ be a $(k - 1)$ -colouring of $G - v$.
- 6 By definition, v is adjacent in G to $\delta < k - 1$ vertices.



Lemma

If G is k -critical, then $\delta \geq k - 1$.

Proof.

- 1 By contradiction.
- 2 Suppose $\delta(G) \leq k - 2$.
- 3 Let v be a vertex of degree δ in G .
- 4 Since G is k -critical, $G - v$ is $(k - 1)$ -colourable.
- 5 Let $(V_1, V_2, \dots, V_{k-1})$ be a $(k - 1)$ -colouring of $G - v$.
- 6 By definition, v is adjacent in G to $\delta < k - 1$ vertices.



Lemma

If G is k -critical, then $\delta \geq k - 1$.

Proof.

- 1 By contradiction.
- 2 Suppose $\delta(G) \leq k - 2$.
- 3 Let v be a vertex of degree δ in G .
- 4 Since G is k -critical, $G - v$ is $(k - 1)$ -colourable.
- 5 Let $(V_1, V_2, \dots, V_{k-1})$ be a $(k - 1)$ -colouring of $G - v$.
- 6 By definition, v is adjacent in G to $\delta < k - 1$ vertices.



Lemma

If G is k -critical, then $\delta \geq k - 1$.

Proof.

- 1 By contradiction.
- 2 Suppose $\delta(G) \leq k - 2$.
- 3 Let v be a vertex of degree δ in G .
- 4 Since G is k -critical, $G - v$ is $(k - 1)$ -colourable.
- 5 Let $(V_1, V_2, \dots, V_{k-1})$ be a $(k - 1)$ - colouring of $G - v$.
- 6 By definition, v is adjacent in G to $\delta < k - 1$ vertices.



Lemma

If G is k -critical, then $\delta \geq k - 1$.

Proof.

- 1 By contradiction.
- 2 Suppose $\delta(G) \leq k - 2$.
- 3 Let v be a vertex of degree δ in G .
- 4 Since G is k -critical, $G - v$ is $(k - 1)$ -colourable.
- 5 Let $(V_1, V_2, \dots, V_{k-1})$ be a $(k - 1)$ - colouring of $G - v$.
- 6 By definition, v is adjacent in G to $\delta < k - 1$ vertices.



- 1 Therefore v must be nonadjacent in G to every vertex of some V_i .
- 2 But then $(V_1, V_2, \dots, V_i \cup \{v\}, \dots, V_{k-1})$ is a $(k-1)$ -colouring of G , a contradiction.
- 3 Thus, $\delta > k-1$. □

- 1 Therefore v must be nonadjacent in G to every vertex of some V_i .
- 2 But then $(V_1, V_2, \dots, V_i \cup \{v\}, \dots, V_{k-1})$ is a $(k - 1)$ -colouring of G , a contradiction.
- 3 Thus, $\delta > k - 1$. □

- 1 Therefore v must be nonadjacent in G to every vertex of some V_i .
- 2 But then $(V_1, V_2, \dots, V_i \cup \{v\}, \dots, V_{k-1})$ is a $(k - 1)$ -colouring of G , a contradiction.
- 3 Thus, $\delta > k - 1$. □

Corollary

Corollary

Every k -chromatic graph has at least k vertices of degree at least $k - 1$.

Proof.

- 1 Let G be a k -chromatic graph.
- 2 Let H be a k -critical subgraph of G .
- 3 By previous Lemma, each vertex of H has degree at least $k - 1$ in H , and hence also in G .
- 4 The corollary now follows since H , being k -chromatic, clearly has at least k vertices.



Corollary

Corollary

Every k -chromatic graph has at least k vertices of degree at least $k - 1$.

Proof.

- 1** Let G be a k -chromatic graph.
- 2** Let H be a k -critical subgraph of G .
- 3** By previous Lemma, each vertex of H has degree at least $k - 1$ in H , and hence also in G .
- 4** The corollary now follows since H , being k -chromatic, clearly has at least k vertices.



Corollary

Corollary

Every k -chromatic graph has at least k vertices of degree at least $k - 1$.

Proof.

- 1** Let G be a k -chromatic graph.
- 2** Let H be a k -critical subgraph of G .
- 3** By previous Lemma, each vertex of H has degree at least $k - 1$ in H , and hence also in G .
- 4** The corollary now follows since H , being k -chromatic, clearly has at least k vertices.



Corollary

Corollary

Every k -chromatic graph has at least k vertices of degree at least $k - 1$.

Proof.

- 1** Let G be a k -chromatic graph.
- 2** Let H be a k -critical subgraph of G .
- 3** By previous Lemma, each vertex of H has degree at least $k - 1$ in H , and hence also in G .
- 4** The corollary now follows since H , being k -chromatic, clearly has at least k vertices.



Corollary

Corollary

Every k -chromatic graph has at least k vertices of degree at least $k - 1$.

Proof.

- 1** Let G be a k -chromatic graph.
- 2** Let H be a k -critical subgraph of G .
- 3** By previous Lemma, each vertex of H has degree at least $k - 1$ in H , and hence also in G .
- 4** The corollary now follows since H , being k -chromatic, clearly has at least k vertices.



Corollary

For any graph G ,

$$\chi \leq \Delta + 1$$

Definition

- 1** Let S be a vertex cut of a connected graph G .
- 2** Let the components of $G - S$ have vertex sets V_1, V_2, \dots, V_n .
- 3** Then the subgraphs $G_i = G[V_i \cup S]$ are called the S -components of G .
- 4** We say that colourings of G_1, G_2, \dots, G_n agree on S if, for every $v \in S$, vertex v is assigned the same colour in each of the colourings.

Definition

- 1** Let S be a vertex cut of a connected graph G .
- 2** Let the components of $G - S$ have vertex sets V_1, V_2, \dots, V_n .
- 3** Then the subgraphs $G_i = G[V_i \cup S]$ are called the S -components of G .
- 4** We say that colourings of G_1, G_2, \dots, G_n agree on S if, for every $v \in S$, vertex v is assigned the same colour in each of the colourings.

Definition

- 1** Let S be a vertex cut of a connected graph G .
- 2** Let the components of $G - S$ have vertex sets V_1, V_2, \dots, V_n .
- 3** Then the subgraphs $G_i = G[V_i \cup S]$ are called the S -components of G .
- 4** We say that colourings of G_1, G_2, \dots, G_n agree on S if, for every $v \in S$, vertex v is assigned the same colour in each of the colourings.

Definition

- 1** Let S be a vertex cut of a connected graph G .
- 2** Let the components of $G - S$ have vertex sets V_1, V_2, \dots, V_n .
- 3** Then the subgraphs $G_i = G[V_i \cup S]$ are called the S -components of G .
- 4** We say that colourings of G_1, G_2, \dots, G_n agree on S if, for every $v \in S$, vertex v is assigned the same colour in each of the colourings.

Example




Figure 8.3. (a) G ; (b) the $\{u, v\}$ -components of G

Theorem

Theorem

In a critical graph, no vertex cut is a clique.

Proof.

- 1 By contradiction.
- 2 Let G be a k -critical graph.
- 3 Suppose that G has a vertex cut S that is a clique.
- 4 Denote the S -components of G by G_1, G_2, \dots, G_n .
- 5 Since G is k -critical, each G_i is $(k - 1)$ -colourable.



Theorem

Theorem

In a critical graph, no vertex cut is a clique.

Proof.

- 1** By contradiction.
- 2** Let G be a k -critical graph.
- 3** Suppose that G has a vertex cut S that is a clique.
- 4** Denote the S -components of G by G_1, G_2, \dots, G_n .
- 5** Since G is k -critical, each G_i is $(k - 1)$ -colourable.



Theorem

Theorem

In a critical graph, no vertex cut is a clique.

Proof.

- 1** By contradiction.
- 2** Let G be a k -critical graph.
- 3** Suppose that G has a vertex cut S that is a clique.
- 4** Denote the S -components of G by G_1, G_2, \dots, G_n .
- 5** Since G is k -critical, each G_i is $(k - 1)$ -colourable.



Theorem

Theorem

In a critical graph, no vertex cut is a clique.

Proof.

- 1** By contradiction.
- 2** Let G be a k -critical graph.
- 3** Suppose that G has a vertex cut S that is a clique.
- 4** Denote the S -components of G by G_1, G_2, \dots, G_n .
- 5** Since G is k -critical, each G_i is $(k - 1)$ -colourable.



Theorem

Theorem

In a critical graph, no vertex cut is a clique.

Proof.

- 1 By contradiction.
- 2 Let G be a k -critical graph.
- 3 Suppose that G has a vertex cut S that is a clique.
- 4 Denote the S -components of G by G_1, G_2, \dots, G_n .
- 5 Since G is k -critical, each G_i is $(k - 1)$ -colourable.



Theorem

Theorem

In a critical graph, no vertex cut is a clique.

Proof.

- 1** By contradiction.
- 2** Let G be a k -critical graph.
- 3** Suppose that G has a vertex cut S that is a clique.
- 4** Denote the S -components of G by G_1, G_2, \dots, G_n .
- 5** Since G is k -critical, each G_i is $(k - 1)$ -colourable.




- 1** Since S is a clique, the vertices in S must receive distinct colours in any $(k - 1)$ -colouring of G_i .
- 2 There are $(k - 1)$ -colourings of G_1, G_2, \dots, G_n which agree on S .
- 3 But these colourings together yield a $(k - 1)$ -colouring of G , a contradiction. □

- 1 Since S is a clique, the vertices in S must receive distinct colours in any $(k - 1)$ -colouring of G_i .
- 2 There are $(k - 1)$ -colourings of G_1, G_2, \dots, G_n which agree on S .
- 3 But these colourings together yield a $(k - 1)$ -colouring of G , a contradiction. □

- 1** Since S is a clique, the vertices in S must receive distinct colours in any $(k - 1)$ -colouring of G_i .
- 2** There are $(k - 1)$ -colourings of G_1, G_2, \dots, G_n which agree on S .
- 3** But these colourings together yield a $(k - 1)$ -colouring of G , a contradiction. □

Example



Is G critical? Why?

Corollary

Corollary

Every critical graph G is a block.

Proof.

- 1 If v is a cut vertex, then $\{v\}$ is a vertex cut.
- 2 $\{v\}$ (trivially,) is a clique.
- 3 Given that G is k -critical for some k .
- 4 By previous theorem, G does not have a Clique.
- 5 A single cut-vertex is a clique.
- 6 So, G does not have a cut-vertex.
- 7 $\Rightarrow G$ is a block.



Corollary

Corollary

Every critical graph G is a block.

Proof.

- 1** If v is a cut vertex, then $\{v\}$ is a vertex cut.
- 2 $\{v\}$ (trivially,) is a clique.
- 3 Given that G is k -critical for some k .
- 4 By previous theorem, G does not have a Clique.
- 5 A single cut-vertex is a clique.
- 6 So, G does not have a cut-vertex.
- 7 $\Rightarrow G$ is a block.



Corollary

Corollary

Every critical graph G is a block.

Proof.

- 1** If v is a cut vertex, then $\{v\}$ is a vertex cut.
- 2** $\{v\}$ (trivially,) is a clique.
- 3** Given that G is k -critical for some k .
- 4** By previous theorem, G does not have a Cut-Vertex.
- 5** A single cut-vertex is a clique.
- 6** So, G does not have a cut-vertex.
- 7** $\Rightarrow G$ is a block.



Corollary

Corollary

Every critical graph G is a block.

Proof.

- 1** If v is a cut vertex, then $\{v\}$ is a vertex cut.
- 2** $\{v\}$ (trivially,) is a clique.
- 3** Given that G is k -critical for some k .
- 4** By previous theorem, G does not have a Cut-Vertex.
- 5** A single cut-vertex is a clique.
- 6** So, G does not have a cut-vertex.
- 7** $\Rightarrow G$ is a block.



Corollary

Corollary

Every critical graph G is a block.

Proof.

- 1** If v is a cut vertex, then $\{v\}$ is a vertex cut.
- 2** $\{v\}$ (trivially,) is a clique.
- 3** Given that G is k -critical for some k .
- 4** By previous theorem, G does not have a Clique.
- 5** A single cut-vertex is a clique.
- 6** So, G does not have a cut-vertex.
- 7** $\Rightarrow G$ is a block.



Corollary

Corollary

Every critical graph G is a block.

Proof.

- 1** If v is a cut vertex, then $\{v\}$ is a vertex cut.
- 2** $\{v\}$ (trivially,) is a clique.
- 3** Given that G is k -critical for some k .
- 4** By previous theorem, G does not have a Clique.
- 5** A single cut-vertex is a clique.
- 6** So, G does not have a cut-vertex.
- 7** $\Rightarrow G$ is a block.



Corollary

Corollary

Every critical graph G is a block.

Proof.

- 1** If v is a cut vertex, then $\{v\}$ is a vertex cut.
- 2** $\{v\}$ (trivially,) is a clique.
- 3** Given that G is k -critical for some k .
- 4** By previous theorem, G does not have a Cut-Vertex.
- 5** A single cut-vertex is a clique.
- 6** So, G does not have a cut-vertex.
- 7** $\Rightarrow G$ is a block.



Corollary

Corollary

Every critical graph G is a block.

Proof.

- 1** If v is a cut vertex, then $\{v\}$ is a vertex cut.
- 2** $\{v\}$ (trivially,) is a clique.
- 3** Given that G is k -critical for some k .
- 4** By previous theorem, G does not have a Cut-Vertex.
- 5** A single cut-vertex is a clique.
- 6** So, G does not have a cut-vertex.
- 7** $\Rightarrow G$ is a block.



Theorem

If G is a connected simple graph and is neither an odd cycle nor a complete graph, then $\chi(G) \leq \Delta$.

- 1 Proof...in Advanced course.
- 2 Use critical graphs/existence of a critical subgraph.
- 3 Not a strong result.
- 4 Implies that only a complete graph or an odd cycle have $\chi(G) = \Delta + 1$.

Theorem

If G is a connected simple graph and is neither an odd cycle nor a complete graph, then $\chi(G) \leq \Delta$.

- 1 Proof...in Advanced course.
- 2 Use critical graphs/existence of a critical subgraph.
- 3 Not a strong result.
- 4 Implies that only a complete graph or an odd cycle have $\chi(G) = \Delta + 1$.

Theorem

If G is a connected simple graph and is neither an odd cycle nor a complete graph, then $\chi(G) \leq \Delta$.

- 1** Proof...in Advanced course.
- 2** Use critical graphs/existence of a critical subgraph.
- 3** Not a strong result.
- 4** Implies that only a complete graph or an odd cycle have $\chi(G) = \Delta + 1$.

Theorem

If G is a connected simple graph and is neither an odd cycle nor a complete graph, then $\chi(G) \leq \Delta$.

- 1** Proof...in Advanced course.
- 2** Use critical graphs/existence of a critical subgraph.
- 3** Not a strong result.
- 4** Implies that only a complete graph or an odd cycle have $\chi(G) = \Delta + 1$.

Theorem

If G is a connected simple graph and is neither an odd cycle nor a complete graph, then $\chi(G) \leq \Delta$.

- 1** Proof...in Advanced course.
- 2** Use critical graphs/existence of a critical subgraph.
- 3** Not a strong result.
- 4** Implies that only a complete graph or an odd cycle have $\chi(G) = \Delta + 1$.

Definition

- 1** A k -edge colouring \mathcal{C} of a loopless graph G is an assignment of k colours, $1, 2, \dots, k$, to the edges of G .
- 2** The colouring \mathcal{C} is proper if no two adjacent edges have the same colour.
- 3** A k -edge colouring can be thought of as a partition (E_1, E_2, \dots, E_k) of E , where E_i denotes the (possibly empty) subset of E assigned colour i .
- 4** A proper k -edge colouring is then a k -edge colouring (E_1, E_2, \dots, E_k) in which each subset E_i is a matching.

Definition

- 1** A k -edge colouring \mathcal{C} of a loopless graph G is an assignment of k colours, $1, 2, \dots, k$, to the edges of G .
- 2** The colouring \mathcal{C} is proper if no two adjacent edges have the same colour.
- 3** A k -edge colouring can be thought of as a partition (E_1, E_2, \dots, E_k) of E , where E_i denotes the (possibly empty) subset of E assigned colour i .
- 4** A proper k -edge colouring is then a k -edge colouring (E_1, E_2, \dots, E_k) in which each subset E_i is a matching.


Definition

- 1** A k -edge colouring \mathcal{C} of a loopless graph G is an assignment of k colours, $1, 2, \dots, k$, to the edges of G .
- 2** The colouring \mathcal{C} is proper if no two adjacent edges have the same colour.
- 3** A k -edge colouring can be thought of as a partition (E_1, E_2, \dots, E_k) of E , where E_i denotes the (possibly empty) subset of E assigned colour i .
- 4** A proper k -edge colouring is then a k -edge colouring (E_1, E_2, \dots, E_k) in which each subset E_i is a matching.

Definition

- 1 A k -edge colouring \mathcal{C} of a loopless graph G is an assignment of k colours, $1, 2, \dots, k$, to the edges of G .
- 2 The colouring \mathcal{C} is proper if no two adjacent edges have the same colour.
- 3 A k -edge colouring can be thought of as a partition (E_1, E_2, \dots, E_k) of E , where E_i denotes the (possibly empty) subset of E assigned colour i .
- 4 A proper k -edge colouring is then a k -edge colouring (E_1, E_2, \dots, E_k) in which each subset E_i is a matching.


Example



- 1 Yellow = $E_1 = \{34\}$
- 2 Brown = $E_2 = \{13, 42\}$
- 3 Blue = $E_3 = \{23\}$
- 4 Green = $E_4 = \{14\}$
- 5 Red = $E_5 = \{23\}$

Note : Even though we have a proper coloring with 5 colors, the coloring is not optimal.

Example



- 1** Yellow = $E_1 = \{34\}$
- 2** Brown = $E_2 = \{13, 42\}$
- 3** Blue = $E_3 = \{23\}$
- 4** Green = $E_4 = \{14\}$
- 5** Red = $E_5 = \{23\}$

Note : Even though we have a proper coloring with 5 colors, the coloring is not optimal.

Definition

- 1** G is k -edge colourable if G has a proper k -edge-colouring.
- 2 Trivially, every loopless graph G is $|E(G)|$ -edge-colourable.
- 3 If G is k -edge-colourable, then G is also l -edge-colourable for every $l > k$.
- 4 The edge chromatic number $\chi'(G)$, of a loopless graph G , is the minimum k for which G is k -edge colourable.
- 5 G is k -edge-chromatic if $\chi'(G) = k$.
- 6 In any proper edge colouring, the edges incident with any one vertex must be assigned different colours. It follows that

$$\chi'(G) \geq \Delta(G)$$

Definition

- 1** G is k -edge colourable if G has a proper k -edge-colouring.
- 2** Trivially, every loopless graph G is $|E(G)|$ -edge-colourable.
- 3** If G is k -edge-colourable, then G is also l -edge-colourable for every $l > k$.
- 4** The edge chromatic number $\chi'(G)$, of a loopless graph G , is the minimum k for which G is k -edge colourable.
- 5** G is k -edge-chromatic if $\chi'(G) = k$.
- 6** In any proper edge colouring, the edges incident with any one vertex must be assigned different colours. It follows that

$$\chi'(G) \geq \Delta(G)$$

Definition

- 1** G is k -edge colourable if G has a proper k -edge-colouring.
- 2** Trivially, every loopless graph G is $|E(G)|$ -edge-colourable.
- 3** If G is k -edge-colourable, then G is also l -edge-colourable for every $l > k$.
- 4** The edge chromatic number $\chi'(G)$, of a loopless graph G , is the minimum k for which G is k -edge colourable.
- 5** G is k -edge-chromatic if $\chi'(G) = k$.
- 6** In any proper edge colouring, the edges incident with any one vertex must be assigned different colours. It follows that

$$\chi'(G) \geq \Delta(G)$$

Definition

- 1** G is k -edge colourable if G has a proper k -edge-colouring.
- 2** Trivially, every loopless graph G is $|E(G)|$ -edge-colourable.
- 3** If G is k -edge-colourable, then G is also l -edge-colourable for every $l > k$.
- 4** The edge chromatic number $\chi'(G)$, of a loopless graph G , is the minimum k for which G is k -edge colourable.
- 5** G is k -edge-chromatic if $\chi'(G) = k$.
- 6** In any proper edge colouring, the edges incident with any one vertex must be assigned different colours. It follows that

$$\chi'(G) \geq \Delta(G)$$

Definition

- 1** G is k -edge colourable if G has a proper k -edge-colouring.
- 2** Trivially, every loopless graph G is $|E(G)|$ -edge-colourable.
- 3** If G is k -edge-colourable, then G is also l -edge-colourable for every $l > k$.
- 4** The edge chromatic number $\chi'(G)$, of a loopless graph G , is the minimum k for which G is k -edge colourable.
- 5** G is k -edge-chromatic if $\chi'(G) = k$.
- 6** In any proper edge colouring, the edges incident with any one vertex must be assigned different colours. It follows that

$$\chi'(G) \geq \Delta(G)$$

Definition

- 1** G is k -edge colourable if G has a proper k -edge-colouring.
- 2** Trivially, every loopless graph G is $|E(G)|$ -edge-colourable.
- 3** If G is k -edge-colourable, then G is also l -edge-colourable for every $l > k$.
- 4** The edge chromatic number $\chi'(G)$, of a loopless graph G , is the minimum k for which G is k -edge colourable.
- 5** G is k -edge-chromatic if $\chi'(G) = k$.
- 6** In any proper edge colouring, the edges incident with any one vertex must be assigned different colours. It follows that

$$\chi'(G) \geq \Delta(G)$$

Gupta(1966)-Vizing(1964) Theorem

Theorem

If G is simple, then either $\chi' = \Delta$ or $\chi' = \Delta + 1$.

No Proof...(Advanced Course)

Classification based on edge coloring

Vizing's theorem divides the class of simple graphs into two classes.

Definition

A simple graph G is said to be of Class-1 if $\chi'(G) = \Delta(G)$ and it is said to be of Class-2 if $\chi'(G) = \Delta(G) + 1$.


Classification based on edge coloring

Vizing's theorem divides the class of simple graphs into two classes.


Definition

A simple graph G is said to be of Class-1 if $\chi'(G) = \Delta(G)$ and it is said to be of Class-2 if $\chi'(G) = \Delta(G) + 1$.


Example-1



I



II




III

1 $\chi'(C_3) = 3 = \Delta(G) + 1$, Hence, Class-2.


2 $\chi'(C_4) = 2 = \Delta(G)$, Hence, Class-1.

3 $\chi'(C_5) = 3 = \Delta(G) + 1$, Hence, Class-2.


Example-1



I



II




III

1 $\chi'(C_3) = 3 = \Delta(G) + 1$, Hence, Class-2.

2 $\chi'(C_4) = 2 = \Delta(G)$, Hence, Class-1.

3 $\chi'(C_5) = 3 = \Delta(G) + 1$, Hence, Class-2.

Example-1




1 $\chi'(C_3) = 3 = \Delta(G) + 1$, Hence, Class-2.

2 $\chi'(C_4) = 2 = \Delta(G)$, Hence, Class-1.

3 $\chi'(C_5) = 3 = \Delta(G) + 1$, Hence, Class-2.

Example-1



1 $\chi'(C_3) = 3 = \Delta(G) + 1$, Hence, Class-2.

2 $\chi'(C_4) = 2 = \Delta(G)$, Hence, Class-1.

3 $\chi'(C_5) = 3 = \Delta(G) + 1$, Hence, Class-2.

König's Theorem

Theorem

For any bipartite graph G , then $\chi'(G) = \Delta$.

Is the converse true?

Before proving we need the following Lemma.

König's Theorem

Theorem

For any bipartite graph G , then $\chi'(G) = \Delta$.

Is the converse true?

Before proving we need the following Lemma.

Lemma

Lemma

Let G be a bipartite graphs and let $d = \Delta(G)$. Then G is a subgraph of a d -regular bipartite graph H .

Proof.

- 1 Let $G[X, Y]$ be the bipartition of $V(G)$, where $X = \{x_1, x_2, \dots, x_s\}$ and $Y = \{y_1, y_2, \dots, y_t\}$.
- 2 Let $G_1[X_1, Y_1]$ be an isomorphic copy of G , where $X_1 = \{x'_1, x'_2, \dots, x'_s\}$ and $Y_1 = \{y'_1, y'_2, \dots, y'_t\}$.
- 3 Construct the bipartite graph $H_1[X \cup Y_1, X_1 \cup Y]$ by adding the following edges:
- 4 Join x_i and x'_i ($1 \leq i \leq s$), if $\deg_G(x_i) < \Delta(G)$.
- 5 Join y_i and y'_i ($1 \leq i \leq t$), if $\deg_G(y_i) < \Delta(G)$.



Lemma

Lemma

Let G be a bipartite graphs and let $d = \Delta(G)$. Then G is a subgraph of a d -regular bipartite graph H .

Proof.

- 1** Let $G[X, Y]$ be the bipartition of $V(G)$, where $X = \{x_1, x_2, \dots, x_s\}$ and $Y = \{y_1, y_2, \dots, y_t\}$.
- 2** Let $G_1[X_1, Y_1]$ be an isomorphic copy of G , where $X_1 = \{x'_1, x'_2, \dots, x'_s\}$ and $Y_1 = \{y'_1, y'_2, \dots, y'_t\}$.
- 3** Construct the bipartite graph $H_1[X \cup Y_1, X_1 \cup Y]$ by adding the following edges:
- 4** Join x_i and x'_i ($1 \leq i \leq s$), if $\deg_G(x_i) < \Delta(G)$.
- 5** Join y_i and y'_i ($1 \leq i \leq t$), if $\deg_G(y_i) < \Delta(G)$.



Lemma

Lemma

Let G be a bipartite graphs and let $d = \Delta(G)$. Then G is a subgraph of a d -regular bipartite graph H .

Proof.

- 1** Let $G[X, Y]$ be the bipartition of $V(G)$, where $X = \{x_1, x_2, \dots, x_s\}$ and $Y = \{y_1, y_2, \dots, y_t\}$.
- 2** Let $G_1[X_1, Y_1]$ be an isomorphic copy of G , where $X_1 = \{x'_1, x'_2, \dots, x'_s\}$ and $Y_1 = \{y'_1, y'_2, \dots, y'_t\}$.
- 3** Construct the bipartite graph $H_1[X \cup Y_1, X_1 \cup Y]$ by adding the following edges:
- 4** Join x_i and x'_i ($1 \leq i \leq s$), if $\deg_G(x_i) < \Delta(G)$.
- 5** Join y_i and y'_i ($1 \leq i \leq t$), if $\deg_G(y_i) < \Delta(G)$.



Lemma

Lemma

Let G be a bipartite graphs and let $d = \Delta(G)$. Then G is a subgraph of a d -regular bipartite graph H .

Proof.

- 1** Let $G[X, Y]$ be the bipartition of $V(G)$, where $X = \{x_1, x_2, \dots, x_s\}$ and $Y = \{y_1, y_2, \dots, y_t\}$.
- 2** Let $G_1[X_1, Y_1]$ be an isomorphic copy of G , where $X_1 = \{x'_1, x'_2, \dots, x'_s\}$ and $Y_1 = \{y'_1, y'_2, \dots, y'_t\}$.
- 3** Construct the bipartite graph $H_1[X \cup Y_1, X_1 \cup Y]$ by adding the following edges:
 - 4** Join x_i and x'_i ($1 \leq i \leq s$), if $\deg_G(x_i) < \Delta(G)$.
 - 5** Join y_i and y'_i ($1 \leq i \leq t$), if $\deg_G(y_i) < \Delta(G)$.



Lemma

Lemma

Let G be a bipartite graphs and let $d = \Delta(G)$. Then G is a subgraph of a d -regular bipartite graph H .

Proof.

- 1** Let $G[X, Y]$ be the bipartition of $V(G)$, where $X = \{x_1, x_2, \dots, x_s\}$ and $Y = \{y_1, y_2, \dots, y_t\}$.
- 2** Let $G_1[X_1, Y_1]$ be an isomorphic copy of G , where $X_1 = \{x'_1, x'_2, \dots, x'_s\}$ and $Y_1 = \{y'_1, y'_2, \dots, y'_t\}$.
- 3** Construct the bipartite graph $H_1[X \cup Y_1, X_1 \cup Y]$ by adding the following edges:
- 4** Join x_i and x'_i ($1 \leq i \leq s$), if $\deg_G(x_i) < \Delta(G)$.
- 5** Join y_i and y'_i ($1 \leq i \leq t$), if $\deg_G(y_i) < \Delta(G)$.



Lemma



Lemma

Let G be a bipartite graphs and let $d = \Delta(G)$. Then G is a subgraph of a d -regular bipartite graph H .

Proof.

- 1** Let $G[X, Y]$ be the bipartition of $V(G)$, where $X = \{x_1, x_2, \dots, x_s\}$ and $Y = \{y_1, y_2, \dots, y_t\}$.
- 2** Let $G_1[X_1, Y_1]$ be an isomorphic copy of G , where $X_1 = \{x'_1, x'_2, \dots, x'_s\}$ and $Y_1 = \{y'_1, y'_2, \dots, y'_t\}$.
- 3** Construct the bipartite graph $H_1[X \cup Y_1, X_1 \cup Y]$ by adding the following edges:
- 4** Join x_i and x'_i ($1 \leq i \leq s$), if $\deg_G(x_i) < \Delta(G)$.
- 5** Join y_i and y'_i ($1 \leq i \leq t$), if $\deg_G(y_i) < \Delta(G)$.





(a) $G \rightarrow H_1$

- 1 Then $\Delta(H_1) = \Delta(G)$ and $\delta(H_1) = \delta(G) + 1$.
- 2 If $\delta(G) + 1 = \Delta(G)$, then H_1 is a $\Delta(G)$ -regular bipartite graph with G as an induced subgraph.
- 3 If not, We repeat the above process to construct H_2 from H_1 .
- 4 Continue the procedure till we find a bipartite graph H_k that is $\Delta(G)$ -regular.
- 5 G is an induced subgraph of H_k . □


- 1 Then $\Delta(H_1) = \Delta(G)$ and $\delta(H_1) = \delta(G) + 1$.
- 2 If $\delta(G) + 1 = \Delta(G)$, then H_1 is a $\Delta(G)$ -regular bipartite graph with G as an induced subgraph.
- 3 If not, We repeat the above process to construct H_2 from H_1 .
- 4 Continue the procedure till we find a bipartite graph H_k that is $\Delta(G)$ -regular.
- 5 G is an induced subgraph of H_k . □

- 1 Then $\Delta(H_1) = \Delta(G)$ and $\delta(H_1) = \delta(G) + 1$.
- 2 If $\delta(G) + 1 = \Delta(G)$, then H_1 is a $\Delta(G)$ -regular bipartite graph with G as an induced subgraph.
- 3 If not, We repeat the above process to construct H_2 from H_1 .
- 4 Continue the procedure till we find a bipartite graph H_k that is $\Delta(G)$ -regular.
- 5 G is an induced subgraph of H_k . □

- 1 Then $\Delta(H_1) = \Delta(G)$ and $\delta(H_1) = \delta(G) + 1$.
- 2 If $\delta(G) + 1 = \Delta(G)$, then H_1 is a $\Delta(G)$ -regular bipartite graph with G as an induced subgraph.
- 3 If not, We repeat the above process to construct H_2 from H_1 .
- 4 Continue the procedure till we find a bipartite graph H_k that is $\Delta(G)$ -regular.
- 5 G is an induced subgraph of H_k . □


- 1 Then $\Delta(H_1) = \Delta(G)$ and $\delta(H_1) = \delta(G) + 1$.
- 2 If $\delta(G) + 1 = \Delta(G)$, then H_1 is a $\Delta(G)$ -regular bipartite graph with G as an induced subgraph.
- 3 If not, We repeat the above process to construct H_2 from H_1 .
- 4 Continue the procedure till we find a bipartite graph H_k that is $\Delta(G)$ -regular.
- 5 G is an induced subgraph of H_k . □

Proof Contd.



H_1 with vertices relabeled.

(b) $H_1 \rightarrow H_2$



H_2^I

Proof of Konig's Theorem

Theorem

For any bipartite graph G , then $\chi'(G) = \Delta$.

Proof.

- 1 We know that (By observation) $\chi'(G) \geq \Delta(G)$.
- 2 We only need to show $\chi'(G) \leq \Delta(G)$.
- 3 We prove by induction on $\Delta(G)$.
- 4 Base case : $\Delta = 0$.. trivially true.
- 5 $\Delta(G) = 1$. $G = K_2$, if connected or all components of G are K_2 's and isolated vertices.
- 6 Need only one color to color the edges of G .
- 7 $\chi'(G) = 1 = \Delta(G)$.
- 8 Induction Hypothesis : Assume true for $\Delta(G) = l$.



Proof of Konig's Theorem

Theorem

For any bipartite graph G , then $\chi'(G) = \Delta$.

Proof.

- 1** We know that (By observation) $\chi'(G) \geq \Delta(G)$.
- 2 We only need to show $\chi'(G) \leq \Delta(G)$.
- 3 We prove by induction on $\Delta(G)$.
- 4 Base case : $\Delta = 0$.. trivially true.
- 5 $\Delta(G) = 1$. $G = K_2$, if connected or all components of G are K_2 's and isolated vertices.
- 6 Need only one color to color the edges of G .
- 7 $\chi'(G) = 1 = \Delta(G)$.
- 8 Induction Hypothesis : Assume true for $\Delta(G) = l$.



Proof of Konig's Theorem

Theorem

For any bipartite graph G , then $\chi'(G) = \Delta$.

Proof.

- 1** We know that (By observation) $\chi'(G) \geq \Delta(G)$.
- 2** We only need to show $\chi'(G) \leq \Delta(G)$.
- 3** We prove by induction on $\Delta(G)$.
- 4** Base case : $\Delta = 0$.. trivially true.
- 5** $\Delta(G) = 1$. $G = K_2$, if connected or all components of G are K_2 's and isolated vertices.
- 6** Need only one color to color the edges of G .
- 7** $\chi'(G) = 1 = \Delta(G)$.
- 8** Induction Hypothesis : Assume true for $\Delta(G) = l$.



Proof of Konig's Theorem

Theorem

For any bipartite graph G , then $\chi'(G) = \Delta$.

Proof.

- 1** We know that (By observation) $\chi'(G) \geq \Delta(G)$.
- 2** We only need to show $\chi'(G) \leq \Delta(G)$.
- 3** We prove by induction on $\Delta(G)$.
- 4** Base case : $\Delta = 0$.. trivially true.
- 5** $\Delta(G) = 1$. $G = K_2$, if connected or all components of G are K_2 's and isolated vertices.
- 6** Need only one color to color the edges of G .
- 7** $\chi'(G) = 1 = \Delta(G)$.
- 8** Induction Hypothesis : Assume true for $\Delta(G) = l$.



Proof of Konig's Theorem

Theorem

For any bipartite graph G , then $\chi'(G) = \Delta$.

Proof.

- 1** We know that (By observation) $\chi'(G) \geq \Delta(G)$.
- 2** We only need to show $\chi'(G) \leq \Delta(G)$.
- 3** We prove by induction on $\Delta(G)$.
- 4** Base case : $\Delta = 0$.. trivially true.
- 5** $\Delta(G) = 1$. $G = K_2$, if connected or all components of G are K_2 's and isolated vertices.
- 6** Need only one color to color the edges of G .
- 7** $\chi'(G) = 1 = \Delta(G)$.
- 8** Induction Hypothesis : Assume true for $\Delta(G) = l$.



Proof of Konig's Theorem

Theorem

For any bipartite graph G , then $\chi'(G) = \Delta$.

Proof.

- 1** We know that (By observation) $\chi'(G) \geq \Delta(G)$.
- 2** We only need to show $\chi'(G) \leq \Delta(G)$.
- 3** We prove by induction on $\Delta(G)$.
- 4** Base case : $\Delta = 0$.. trivially true.
- 5** $\Delta(G) = 1$. $G = K_2$, if connected or all components of G are K_2 's and isolated vertices.
- 6** Need only one color to color the edges of G .
- 7** $\chi'(G) = 1 = \Delta(G)$.
- 8** Induction Hypothesis : Assume true for $\Delta(G) = l$.



Proof of Konig's Theorem

Theorem

For any bipartite graph G , then $\chi'(G) = \Delta$.

Proof.

- 1** We know that (By observation) $\chi'(G) \geq \Delta(G)$.
- 2** We only need to show $\chi'(G) \leq \Delta(G)$.
- 3** We prove by induction on $\Delta(G)$.
- 4** Base case : $\Delta = 0$.. trivially true.
- 5** $\Delta(G) = 1$. $G = K_2$, if connected or all components of G are K_2 's and isolated vertices.
- 6** Need only one color to color the edges of G .
- 7** $\chi'(G) = 1 = \Delta(G)$.
- 8** Induction Hypothesis : Assume true for $\Delta(G) = l$.



Proof of Konig's Theorem

Theorem

For any bipartite graph G , then $\chi'(G) = \Delta$.

Proof.

- 1** We know that (By observation) $\chi'(G) \geq \Delta(G)$.
- 2** We only need to show $\chi'(G) \leq \Delta(G)$.
- 3** We prove by induction on $\Delta(G)$.
- 4** Base case : $\Delta = 0$.. trivially true.
- 5** $\Delta(G) = 1$. $G = K_2$, if connected or all components of G are K_2 's and isolated vertices.
- 6** Need only one color to color the edges of G .
- 7** $\chi'(G) = 1 = \Delta(G)$.
- 8** Induction Hypothesis : Assume true for $\Delta(G) = l$.



Proof of Konig's Theorem

Theorem

For any bipartite graph G , then $\chi'(G) = \Delta$.

Proof.

- 1** We know that (By observation) $\chi'(G) \geq \Delta(G)$.
- 2** We only need to show $\chi'(G) \leq \Delta(G)$.
- 3** We prove by induction on $\Delta(G)$.
- 4** Base case : $\Delta = 0$.. trivially true.
- 5** $\Delta(G) = 1$. $G = K_2$, if connected or all components of G are K_2 's and isolated vertices.
- 6** Need only one color to color the edges of G .
- 7** $\chi'(G) = 1 = \Delta(G)$.
- 8** Induction Hypothesis : Assume true for $\Delta(G) = l$.



Proof.

- 1** Let G be a bipartite graph with $\Delta(G) = l + 1$.
- 2** By Previous lemma, there exists a H_k , which is $\Delta(G)$ -regular, bipartite and G is an induced subgraph of H_k .
- 3** By Hall's matching theorem. H_k contains a Perfect Matching M .
- 4** $H_k - M$ has a maximum degree $\Delta(G) - 1 = l$.
- 5** By induction, $H_k - M$ has a proper $\Delta - 1$ edge coloring.
- 6** Give the Δ color to all edges of M .
- 7** Combining together we get a Δ -edge coloring of G .



Proof.

- 1 Let G be a bipartite graph with $\Delta(G) = l + 1$.
- 2 By Previous lemma, there exists a H_k , which is $\Delta(G)$ -regular, bipartite and G is an induced subgraph of H_k .
- 3 By Hall's matching theorem. H_k contains a Perfect Matching M .
- 4 $H_k - M$ has a maximum degree $\Delta(G) - 1 = l$.
- 5 By induction, $H_k - M$ has a proper $\Delta - 1$ edge coloring.
- 6 Give the Δ color to all edges of M .
- 7 Combining together we get a Δ -edge coloring of G .



Proof.

- 1** Let G be a bipartite graph with $\Delta(G) = l + 1$.
- 2** By Previous lemma, there exists a H_k , which is $\Delta(G)$ -regular, bipartite and G is an induced subgraph of H_k .
- 3** By Hall's matching theorem. H_k contains a Perfect Matching M .
- 4** $H_k - M$ has a maximum degree $\Delta(G) - 1 = l$.
- 5** By induction, $H_k - M$ has a proper $\Delta - 1$ edge coloring.
- 6** Give the Δ color to all edges of M .
- 7** Combining together we get a Δ -edge coloring of G .



Proof.

- 1** Let G be a bipartite graph with $\Delta(G) = l + 1$.
- 2** By Previous lemma, there exists a H_k , which is $\Delta(G)$ -regular, bipartite and G is an induced subgraph of H_k .
- 3** By Hall's matching theorem. H_k contains a Perfect Matching M .
- 4** $H_k - M$ has a maximum degree $\Delta(G) - 1 = l$.
- 5** By induction, $H_k - M$ has a proper $\Delta - 1$ edge coloring.
- 6** Give the Δ color to all edges of M .
- 7** Combining together we get a Δ -edge coloring of G .



Proof.

- 1** Let G be a bipartite graph with $\Delta(G) = l + 1$.
- 2** By Previous lemma, there exists a H_k , which is $\Delta(G)$ -regular, bipartite and G is an induced subgraph of H_k .
- 3** By Hall's matching theorem. H_k contains a Perfect Matching M .
- 4** $H_k - M$ has a maximum degree $\Delta(G) - 1 = l$.
- 5** By induction, $H_k - M$ has a proper $\Delta - 1$ edge coloring.
- 6** Give the Δ color to all edges of M .
- 7** Combining together we get a Δ -edge coloring of G .



Proof.

- 1** Let G be a bipartite graph with $\Delta(G) = l + 1$.
- 2** By Previous lemma, there exists a H_k , which is $\Delta(G)$ -regular, bipartite and G is an induced subgraph of H_k .
- 3** By Hall's matching theorem. H_k contains a Perfect Matching M .
- 4** $H_k - M$ has a maximum degree $\Delta(G) - 1 = l$.
- 5** By induction, $H_k - M$ has a proper $\Delta - 1$ edge coloring.
- 6** Give the Δ color to all edges of M .
- 7** Combining together we get a Δ -edge coloring of G .



Proof.

- 1** Let G be a bipartite graph with $\Delta(G) = l + 1$.
- 2** By Previous lemma, there exists a H_k , which is $\Delta(G)$ -regular, bipartite and G is an induced subgraph of H_k .
- 3** By Hall's matching theorem. H_k contains a Perfect Matching M .
- 4** $H_k - M$ has a maximum degree $\Delta(G) - 1 = l$.
- 5** By induction, $H_k - M$ has a proper $\Delta - 1$ edge coloring.
- 6** Give the Δ color to all edges of M .
- 7** Combining together we get a Δ -edge coloring of G .



We derive a few sufficient conditions for a graph to be of Class-2.

Theorem

For any graph G ,

$$\chi'(G) \geq \left\lceil \frac{|E(G)|}{\alpha'(G)} \right\rceil$$

Proof.

- 1 Let $C = (M_1, \dots, M_{\chi'})$ be a χ' -edge-coloring of G .
- 2 Then, $|E(G)| = |M_1| + |M_2| + \dots + |M_{\chi'}| \leq |\alpha'(G)| + |\alpha'(G)| + \dots + |\alpha'(G)| = \alpha'(G) \times \chi'(G)$.
- 3 " \Rightarrow " $\chi'(G) \geq \left\lceil \frac{|E(G)|}{\alpha'(G)} \right\rceil$



We derive a few sufficient conditions for a graph to be of Class-2.

Theorem

For any graph G ,

$$\chi'(G) \geq \left\lceil \frac{|E(G)|}{\alpha'(G)} \right\rceil$$

Proof.

- 1 Let $C = (M_1, \dots, M_{\chi'})$ be a χ' -edge-coloring of G .
- 2 Then, $|E(G)| = |M_1| + |M_2| + \dots + |M_{\chi'}| \leq |\alpha'(G)| + |\alpha'(G)| + \dots + |\alpha'(G)| = \alpha'(G) \times \chi'(G)$.
- 3 " \Rightarrow " $\chi'(G) \geq \left\lceil \frac{|E(G)|}{\alpha'(G)} \right\rceil$



We derive a few sufficient conditions for a graph to be of Class-2.

Theorem

For any graph G ,

$$\chi'(G) \geq \left\lceil \frac{|E(G)|}{\alpha'(G)} \right\rceil$$

Proof.

1 Let $C = (M_1, \dots, M_{\chi'})$ be a χ' -edge-coloring of G .

2 Then, $|E(G)| = |M_1| + |M_2| + \dots + |M_{\chi'}| \leq |\alpha'(G)| + |\alpha'(G)| + \dots + |\alpha'(G)| = \alpha'(G) \times \chi'(G)$.

3 " \Rightarrow " $\chi'(G) \geq \left\lceil \frac{|E(G)|}{\alpha'(G)} \right\rceil$



We derive a few sufficient conditions for a graph to be of Class-2.

Theorem

For any graph G ,

$$\chi'(G) \geq \left\lceil \frac{|E(G)|}{\alpha'(G)} \right\rceil$$

Proof.

1 Let $C = (M_1, \dots, M_{\chi'})$ be a χ' -edge-coloring of G .

2 Then, $|E(G)| = |M_1| + |M_2| + \dots + |M_{\chi'}| \leq$
 $|\alpha'(G)| + |\alpha'(G)| + \dots + |\alpha'(G)| = \alpha'(G) \times \chi'(G).$

3 " \Rightarrow " $\chi'(G) \geq \left\lceil \frac{|E(G)|}{\alpha'(G)} \right\rceil$



We derive a few sufficient conditions for a graph to be of Class-2.

Theorem

For any graph G ,

$$\chi'(G) \geq \left\lceil \frac{|E(G)|}{\alpha'(G)} \right\rceil$$

Proof.

- 1** Let $C = (M_1, \dots, M_{\chi'})$ be a χ' -edge-coloring of G .
- 2** Then, $|E(G)| = |M_1| + |M_2| + \dots + |M_{\chi'}| \leq |\alpha'(G)| + |\alpha'(G)| + \dots + |\alpha'(G)| = \alpha'(G) \times \chi'(G)$.
- 3** " \Rightarrow " $\chi'(G) \geq \left\lceil \frac{|E(G)|}{\alpha'(G)} \right\rceil$



Corollary

If G is a simple graph with $|E(G)| > \Delta(G) \times \chi'(G)$, then G is a Class-2 graph.

Proof.

1 By previous theorem,

$$\chi'(G) \geq \left\lceil \frac{|E(G)|}{\alpha'(G)} \right\rceil$$

$$2 \quad \chi'(G) \geq \left\lceil \frac{|E(G)|}{\alpha'(G)} \right\rceil > \left\lceil \frac{\Delta(G) \times \alpha'(G)}{\alpha'(G)} \right\rceil = \Delta(G)$$

$$3 \quad \text{By Gupta-Vizing, } \chi'(G) = \Delta(G) + 1$$

4 Hence G is of class 2.



Corollary

If G is a simple graph with $|E(G)| > \Delta(G) \times \chi'(G)$, then G is a Class-2 graph.

Proof.

1 By previous theorem,

$$\chi'(G) \geq \left\lceil \frac{|E(G)|}{\alpha'(G)} \right\rceil$$

2 $\chi'(G) \geq \left\lceil \frac{|E(G)|}{\alpha'(G)} \right\rceil > \left\lceil \frac{\Delta(G) \times \alpha'(G)}{\alpha'(G)} \right\rceil = \Delta(G)$

3 By Gupta-Vizing, $\chi'(G) = \Delta(G) + 1$

4 Hence G is of class 2.



Corollary

If G is a simple graph with $|E(G)| > \Delta(G) \times \chi'(G)$, then G is a Class-2 graph.

Proof.

1 By previous theorem,

$$\chi'(G) \geq \left\lceil \frac{|E(G)|}{\alpha'(G)} \right\rceil$$

2 $\chi'(G) \geq \left\lceil \frac{|E(G)|}{\alpha'(G)} \right\rceil > \left\lceil \frac{\Delta(G) \times \alpha'(G)}{\alpha'(G)} \right\rceil = \Delta(G)$

3 By Gupta-Vizing, $\chi'(G) = \Delta(G) + 1$

4 Hence G is of class 2.



Corollary

If G is a simple graph with $|E(G)| > \Delta(G) \times \chi'(G)$, then G is a Class-2 graph.

Proof.

1 By previous theorem,

$$\chi'(G) \geq \left\lceil \frac{|E(G)|}{\alpha'(G)} \right\rceil$$

2 $\chi'(G) \geq \left\lceil \frac{|E(G)|}{\alpha'(G)} \right\rceil > \left\lceil \frac{\Delta(G) \times \alpha'(G)}{\alpha'(G)} \right\rceil = \Delta(G)$

3 By Gupta-Vizing, $\chi'(G) = \Delta(G) + 1$

4 Hence G is of class 2.



Corollary

If G is a simple graph with $|E(G)| > \Delta(G) \times \chi'(G)$, then G is a Class-2 graph.

Proof.

1 By previous theorem,

$$\chi'(G) \geq \left\lceil \frac{|E(G)|}{\alpha'(G)} \right\rceil$$

2 $\chi'(G) \geq \left\lceil \frac{|E(G)|}{\alpha'(G)} \right\rceil > \left\lceil \frac{\Delta(G) \times \alpha'(G)}{\alpha'(G)} \right\rceil = \Delta(G)$

3 By Gupta-Vizing, $\chi'(G) = \Delta(G) + 1$

4 Hence G is of class 2.

