



Connectivity

September 2, 2024




- Deletion of any edge disconnects the graph.

Examples




- Deletion of any edge does not disconnect the graph.
- Deletion of vertex disconnects the graph.

Examples



- ▶ Deletion of any edge does not disconnect the graph.
- ▶ Deletion of vertex disconnects the graph.

Examples



- No cut edges and no cut vertices.

- ▶ Each successive graph is more strongly connected than the previous one.
- ▶ We define two concepts that measures the extent to which a given graph is connected.
- ▶ (Vertex)-Connectivity and Edge-Connectivity.

- ▶ Each successive graph is more strongly connected than the previous one.
- ▶ We define two concepts that measures the extent to which a given graph is connected.
- ▶ (Vertex)-Connectivity and Edge-Connectivity.

- ▶ Each successive graph is more strongly connected than the previous one.
- ▶ We define two concepts that measures the extent to which a given graph is connected.
- ▶ (Vertex)-Connectivity and Edge-Connectivity.

Definition

A vertex subset $W \subseteq V(G)$ is called a vertex-cut if $G \setminus W$ is disconnected or $G \setminus W$ is trivial.

- 1 $\kappa(G) = \min\{|W| : W \subseteq V(G) \text{ and } W \text{ is a vertex-cut}\}$
- 2 $\kappa(G)$ is called the vertex-connectivity of G .
- 3 That is, $\kappa(G)$ is the minimum number of vertices whose deletion disconnects G or results in a graph with a single vertex.
- 4 Any vertex-cut $K \subseteq V(G)$ such that $|K| = \kappa(G)$ is called a minimum vertex-cut of G .
- 5 A graph G is said to be k -vertex-connected, if $\kappa(G) \geq k$.
- 6 All non-trivial connected graphs are 1-connected.

Definition

A vertex subset $W \subseteq V(G)$ is called a vertex-cut if $G \setminus W$ is disconnected or $G \setminus W$ is trivial.

- 1 $\kappa(G) = \min\{|W| : W \subseteq V(G) \text{ and } W \text{ is a vertex-cut}\}$
- 2 $\kappa(G)$ is called the vertex-connectivity of G .
- 3 That is, $\kappa(G)$ is the minimum number of vertices whose deletion disconnects G or results in a graph with a single vertex.
- 4 Any vertex-cut $K \subseteq V(G)$ such that $|K| = \kappa(G)$ is called a minimum vertex-cut of G .
- 5 A graph G is said to be k -vertex-connected, if $\kappa(G) \geq k$.
- 6 All non-trivial connected graphs are 1-connected.

Definition

A vertex subset $W \subseteq V(G)$ is called a vertex-cut if $G \setminus W$ is disconnected or $G \setminus W$ is trivial.

- 1 $\kappa(G) = \min\{|W| : W \subseteq V(G) \text{ and } W \text{ is a vertex-cut}\}$
- 2 $\kappa(G)$ is called the vertex-connectivity of G .
- 3 That is, $\kappa(G)$ is the minimum number of vertices whose deletion disconnects G or results in a graph with a single vertex.
- 4 Any vertex-cut $K \subseteq V(G)$ such that $|K| = \kappa(G)$ is called a minimum vertex-cut of G .
- 5 A graph G is said to be k -vertex-connected, if $\kappa(G) \geq k$.
- 6 All non-trivial connected graphs are 1-connected.

Definition

A vertex subset $W \subseteq V(G)$ is called a vertex-cut if $G \setminus W$ is disconnected or $G \setminus W$ is trivial.

- 1 $\kappa(G) = \min\{|W| : W \subseteq V(G) \text{ and } W \text{ is a vertex-cut}\}$
- 2 $\kappa(G)$ is called the vertex-connectivity of G .
- 3 That is, $\kappa(G)$ is the minimum number of vertices whose deletion disconnects G or results in a graph with a single vertex.
- 4 Any vertex-cut $K \subseteq V(G)$ such that $|K| = \kappa(G)$ is called a minimum vertex-cut of G .
- 5 A graph G is said to be k -vertex-connected, if $\kappa(G) \geq k$.
- 6 All non-trivial connected graphs are 1-connected.

Definition

A vertex subset $W \subseteq V(G)$ is called a vertex-cut if $G \setminus W$ is disconnected or $G \setminus W$ is trivial.

- 1 $\kappa(G) = \min\{|W| : W \subseteq V(G) \text{ and } W \text{ is a vertex-cut}\}$
- 2 $\kappa(G)$ is called the vertex-connectivity of G .
- 3 That is, $\kappa(G)$ is the minimum number of vertices whose deletion disconnects G or results in a graph with a single vertex.
- 4 Any vertex-cut $K \subseteq V(G)$ such that $|K| = \kappa(G)$ is called a minimum vertex-cut of G .
- 5 A graph G is said to be k -vertex-connected, if $\kappa(G) \geq k$.
- 6 All non-trivial connected graphs are 1-connected.

Definition

A vertex subset $W \subseteq V(G)$ is called a vertex-cut if $G \setminus W$ is disconnected or $G \setminus W$ is trivial.


- 1 $\kappa(G) = \min\{|W| : W \subseteq V(G) \text{ and } W \text{ is a vertex-cut}\}$
- 2 $\kappa(G)$ is called the vertex-connectivity of G .
- 3 That is, $\kappa(G)$ is the minimum number of vertices whose deletion disconnects G or results in a graph with a single vertex.
- 4 Any vertex-cut $K \subseteq V(G)$ such that $|K| = \kappa(G)$ is called a minimum vertex-cut of G .
- 5 A graph G is said to be k -vertex-connected, if $\kappa(G) \geq k$.
- 6 All non-trivial connected graphs are 1-connected.

Definition

A vertex subset $W \subseteq V(G)$ is called a vertex-cut if $G \setminus W$ is disconnected or $G \setminus W$ is trivial.


- 1 $\kappa(G) = \min\{|W| : W \subseteq V(G) \text{ and } W \text{ is a vertex-cut}\}$
- 2 $\kappa(G)$ is called the vertex-connectivity of G .
- 3 That is, $\kappa(G)$ is the minimum number of vertices whose deletion disconnects G or results in a graph with a single vertex.
- 4 Any vertex-cut $K \subseteq V(G)$ such that $|K| = \kappa(G)$ is called a minimum vertex-cut of G .
- 5 A graph G is said to be k -vertex-connected, if $\kappa(G) \geq k$.
- 6 All non-trivial connected graphs are 1-connected.

Examples



- ▶ Vertex cut : $\{a, d\}$ or $\{b, d\}$ or $\{a, b, d\}$

Examples



- Vertex cut : $\{a, d\}$ or $\{b, d\}$ or $\{a, b, d\}$

Definition

An **edge cut** is a subset $F \subseteq E(G)$ of the form $[S, S']$ where V is partitioned into $S \neq \emptyset$ and $S' \neq \emptyset$ such that each edge in F has one end in S and other in S' .

Or Equivalently(???), for a non-trivial Graph,

Definition

An edge subset $F \subseteq E(G)$ is called a **disconnecting set** if $G \setminus F$ is disconnected or $G \setminus F$ is a single vertex (trivial) graph.

Definition

An **edge cut** is a subset $F \subseteq E(G)$ of the form $[S, S']$ where V is partitioned into $S \neq \emptyset$ and $S' \neq \emptyset$ such that each edge in F has one end in S and other in S' .

Or Equivalently(???), for a non-trivial Graph,

Definition

An edge subset $F \subseteq E(G)$ is called a **disconnecting set** if $G \setminus F$ is disconnected or $G \setminus F$ is a single vertex (trivial) graph.

Remark :

- 1 Every edge cut is a disconnecting set.
- 2 Every minimal disconnecting set is an edge cut.
- 3 For a disconnecting set F , let H be a component of $G - F$. Then $[V(H), V(G) - V(H)]$ is an edge cut.

Remark :

- 1 Every edge cut is a disconnecting set.
- 2 Every minimal disconnecting set is an edge cut.
- 3 For a disconnecting set F , let H be a component of $G - F$. Then $[V(H), V(G) - V(H)]$ is an edge cut.

Remark :

- 1 Every edge cut is a disconnecting set.
- 2 Every minimal disconnecting set is an edge cut.
- 3 For a disconnecting set F , let H be a component of $G - F$. Then $[V(H), V(G) - V(H)]$ is an edge cut.

- 1 A k -edge cut is an edge cut of k elements.
- 2 $\kappa'(G) = \min\{|F| : F \subseteq E(G) \text{ and } F \text{ is an edge-cut}\}$
- 3 $\kappa'(G)$ is called the edge-connectivity of G .
- 4 That is, $\kappa'(G)$ is the minimum number of edges whose deletion disconnects G or results in a graph with a single vertex.
- 5 Any edge-cut $F \subseteq E(G)$ such that $|F| = \kappa'(G)$ is called a minimum edge-cut of G .
- 6 A graph G is said to be k -edge-connected, if $\kappa'(G) \geq k$.

- 1 A k -edge cut is an edge cut of k elements.
- 2 $\kappa'(G) = \min\{|F| : F \subseteq E(G) \text{ and } F \text{ is an edge-cut}\}$
- 3 $\kappa'(G)$ is called the edge-connectivity of G .
- 4 That is, $\kappa'(G)$ is the minimum number of edges whose deletion disconnects G or results in a graph with a single vertex.
- 5 Any edge-cut $F \subseteq E(G)$ such that $|F| = \kappa'(G)$ is called a minimum edge-cut of G .
- 6 A graph G is said to be k -edge-connected, if $\kappa'(G) \geq k$.


- 1 A k -edge cut is an edge cut of k elements.
- 2 $\kappa'(G) = \min\{|F| : F \subseteq E(G) \text{ and } F \text{ is an edge-cut}\}$
- 3 $\kappa'(G)$ is called the edge-connectivity of G .
- 4 That is, $\kappa'(G)$ is the minimum number of edges whose deletion disconnects G or results in a graph with a single vertex.
- 5 Any edge-cut $F \subseteq E(G)$ such that $|F| = \kappa'(G)$ is called a minimum edge-cut of G .
- 6 A graph G is said to be k -edge-connected, if $\kappa'(G) \geq k$.

- 1 A k -edge cut is an edge cut of k elements.
- 2 $\kappa'(G) = \min\{|F| : F \subseteq E(G) \text{ and } F \text{ is an edge-cut}\}$
- 3 $\kappa'(G)$ is called the edge-connectivity of G .
- 4 That is, $\kappa'(G)$ is the minimum number of edges whose deletion disconnects G or results in a graph with a single vertex.
- 5 Any edge-cut $F \subseteq E(G)$ such that $|F| = \kappa'(G)$ is called a minimum edge-cut of G .
- 6 A graph G is said to be k -edge-connected, if $\kappa'(G) \geq k$.

- 1 A k -edge cut is an edge cut of k elements.
- 2 $\kappa'(G) = \min\{|F| : F \subseteq E(G) \text{ and } F \text{ is an edge-cut}\}$
- 3 $\kappa'(G)$ is called the edge-connectivity of G .
- 4 That is, $\kappa'(G)$ is the minimum number of edges whose deletion disconnects G or results in a graph with a single vertex.
- 5 Any edge-cut $F \subseteq E(G)$ such that $|F| = \kappa'(G)$ is called a minimum edge-cut of G .
- 6 A graph G is said to be k -edge-connected, if $\kappa'(G) \geq k$.


- 1 A k -edge cut is an edge cut of k elements.
- 2 $\kappa'(G) = \min\{|F| : F \subseteq E(G) \text{ and } F \text{ is an edge-cut}\}$
- 3 $\kappa'(G)$ is called the edge-connectivity of G .
- 4 That is, $\kappa'(G)$ is the minimum number of edges whose deletion disconnects G or results in a graph with a single vertex.
- 5 Any edge-cut $F \subseteq E(G)$ such that $|F| = \kappa'(G)$ is called a minimum edge-cut of G .
- 6 A graph G is said to be k -edge-connected, if $\kappa'(G) \geq k$.

Example



► Edge cut : $\{be, de\}$ or $\{ac, cd\}$ or $\{ab, ac, ad\}$

Example



- Edge cut : $\{be, de\}$ or $\{ac, cd\}$ or $\{ab, ac, ad\}$

- 1 $\kappa'(G) = 0$ if and only if G is disconnected or G is trivial.
- 2 $\kappa'(G) = 1$ if and only if G is connected and G has a cut-edge.
- 3 $\kappa'(G) \leq \delta(G)$; this follows since by deleting all the edges incident with a vertex of minimum degree, we disconnect the graph.
- 4 A graph G may have many minimum edge-cuts but $\kappa'(G)$ is unique.
- 5 If G is k -edge-connected, then it is t -edge-connected, for every t , $1 \leq t \leq k$.
- 6 If F is a minimum edge-cut, then $G \setminus F$ contains exactly two components A and B such that $[V(A), V(B)] = F$.

- 1 $\kappa'(G) = 0$ if and only if G is disconnected or G is trivial.
- 2 $\kappa'(G) = 1$ if and only if G is connected and G has a cut-edge.
- 3 $\kappa'(G) \leq \delta(G)$; this follows since by deleting all the edges incident with a vertex of minimum degree, we disconnect the graph.
- 4 A graph G may have many minimum edge-cuts but $\kappa'(G)$ is unique.
- 5 If G is k -edge-connected, then it is t -edge-connected, for every t , $1 \leq t \leq k$.
- 6 If F is a minimum edge-cut, then $G \setminus F$ contains exactly two components A and B such that $[V(A), V(B)] = F$.

- 1 $\kappa'(G) = 0$ if and only if G is disconnected or G is trivial.
- 2 $\kappa'(G) = 1$ if and only if G is connected and G has a cut-edge.
- 3 $\kappa'(G) \leq \delta(G)$; this follows since by deleting all the edges incident with a vertex of minimum degree, we disconnect the graph.
- 4 A graph G may have many minimum edge-cuts but $\kappa'(G)$ is unique.
- 5 If G is k -edge-connected, then it is t -edge-connected, for every t , $1 \leq t \leq k$.
- 6 If F is a minimum edge-cut, then $G \setminus F$ contains exactly two components A and B such that $[V(A), V(B)] = F$.

Properties-Edge cut

- 1 $\kappa'(G) = 0$ if and only if G is disconnected or G is trivial.
- 2 $\kappa'(G) = 1$ if and only if G is connected and G has a cut-edge.
- 3 $\kappa'(G) \leq \delta(G)$; this follows since by deleting all the edges incident with a vertex of minimum degree, we disconnect the graph.
- 4 A graph G may have many minimum edge-cuts but $\kappa'(G)$ is unique.
- 5 If G is k -edge-connected, then it is t -edge-connected, for every t , $1 \leq t \leq k$.
- 6 If F is a minimum edge-cut, then $G \setminus F$ contains exactly two components A and B such that $[V(A), V(B)] = F$.

- 1 $\kappa'(G) = 0$ if and only if G is disconnected or G is trivial.
- 2 $\kappa'(G) = 1$ if and only if G is connected and G has a cut-edge.
- 3 $\kappa'(G) \leq \delta(G)$; this follows since by deleting all the edges incident with a vertex of minimum degree, we disconnect the graph.
- 4 A graph G may have many minimum edge-cuts but $\kappa'(G)$ is unique.
- 5 If G is k -edge-connected, then it is t -edge-connected, for every t , $1 \leq t \leq k$.
- 6 If F is a minimum edge-cut, then $G \setminus F$ contains exactly two components A and B such that $[V(A), V(B)] = F$.

- 1 $\kappa'(G) = 0$ if and only if G is disconnected or G is trivial.
- 2 $\kappa'(G) = 1$ if and only if G is connected and G has a cut-edge.
- 3 $\kappa'(G) \leq \delta(G)$; this follows since by deleting all the edges incident with a vertex of minimum degree, we disconnect the graph.
- 4 A graph G may have many minimum edge-cuts but $\kappa'(G)$ is unique.
- 5 If G is k -edge-connected, then it is t -edge-connected, for every t , $1 \leq t \leq k$.
- 6 If F is a minimum edge-cut, then $G \setminus F$ contains exactly two components A and B such that $[V(A), V(B)] = F$.

Theorem

For any graph G with at least two vertices,

$$\kappa(G) \leq \kappa'(G) \leq \delta(G)$$

Proof.

1 We have observed : $\kappa'(G) \leq \delta(G)$.

2 Only need to prove $\kappa(G) \leq \kappa'(G)$.

3 If G is disconnected or trivial,

$$\kappa(G) = \kappa'(G) = 0 \leq \delta(G)$$



Theorem

For any graph G with at least two vertices,

$$\kappa(G) \leq \kappa'(G) \leq \delta(G)$$

Proof.

- 1** We have observed : $\kappa'(G) \leq \delta(G)$.
- 2** Only need to prove $\kappa(G) \leq \kappa'(G)$.
- 3** If G is disconnected or trivial,

$$\kappa(G) = \kappa'(G) = 0 \leq \delta(G)$$



Theorem

For any graph G with at least two vertices,

$$\kappa(G) \leq \kappa'(G) \leq \delta(G)$$

Proof.

- 1** We have observed : $\kappa'(G) \leq \delta(G)$.
- 2** Only need to prove $\kappa(G) \leq \kappa'(G)$.
- 3** If G is disconnected or trivial,

$$\kappa(G) = \kappa'(G) = 0 \leq \delta(G)$$



Proof.

Now we assume that G is connected.

1 If $|V(G)| = 2$, then $G = K_2$.

2 Then,

$$\kappa(K_2) = 1$$

(makes th graph trivial).

3 $\kappa'(G) = 1$

(Makes the graphs disconnected with two vertices).

4 $\delta(G) = 1$.

5 $\kappa(G) = \kappa'(G) = \delta(G) = 1$.



Proof.

Now we assume that G is connected.

1 If $|V(G)| = 2$, then $G = K_2$.

2 Then,

$$\kappa(K_2) = 1$$

(makes th graph trivial).

3 $\kappa'(G) = 1$

(Makes the graphs disconnected with two vertices).

4 $\delta(G) = 1$.

5 $\kappa(G) = \kappa'(G) = \delta(G) = 1$.



Proof.

Now we assume that G is connected.

1 If $|V(G)| = 2$, then $G = K_2$.

2 Then,

$$\kappa(K_2) = 1$$

(makes th graph trivial).

3 $\kappa'(G) = 1$

(Makes the graphs disconnected with two vertices).

4 $\delta(G) = 1$.

5 $\kappa(G) = \kappa'(G) = \delta(G) = 1$.



Proof.

Now we assume that G is connected.

1 If $|V(G)| = 2$, then $G = K_2$.

2 Then,

$$\kappa(K_2) = 1$$

(makes th graph trivial).

3 $\kappa'(G) = 1$

(Makes the graphs disconnected with two vertices).

4 $\delta(G) = 1$.

5 $\kappa(G) = \kappa'(G) = \delta(G) = 1$.



Proof.

Now we assume that G is connected.

1 If $|V(G)| = 2$, then $G = K_2$.

2 Then,

$$\kappa(K_2) = 1$$

(makes th graph trivial).

3 $\kappa'(G) = 1$

(Makes the graphs disconnected with two vertices).

4 $\delta(G) = 1$.

5 $\kappa(G) = \kappa'(G) = \delta(G) = 1$.



Let G be connected and $|V(G)| \geq 3$. We now discuss two possible cases:

- 1 Case 1 : G is simple.
- 2 Case 2 : G is any graph.

It is enough we prove the case for simple graphs. (Explanation Later).

Let G be connected and $|V(G)| \geq 3$. We now discuss two possible cases:

- 1 Case 1 : G is simple.
- 2 Case 2 : G is any graph.

It is enough we prove the case for simple graphs. (Explanation Later).

Let G be connected and $|V(G)| \geq 3$. We now discuss two possible cases:

- 1 Case 1 : G is simple.
- 2 Case 2 : G is any graph.

It is enough we prove the case for simple graphs. (Explanation Later).

Case 1 : G is simple, connected and $|V(G)| \geq 3$. Need to show $\kappa(G) \leq \kappa'(G)$.

- 1 Let $k = \kappa'(G)$.
- 2 Let $F = \{e_1, e_2, \dots, e_k\}$ be an edge cut of G where $e_i = u_i v_i$, $u_i, v_i \in V$. (Note : u_i 's and v_j 's need not be distinct.)
- 3 Need to construct a vertex cut from F .
- 4 Let $H = G \setminus \{\{u_i : 1 \leq i \leq k-1\} \cup \{u_k, v_k\}\}$
- 5 We retain the edge e_k in H and remove all other edges of F .

Case 1 : G is simple, connected and $|V(G)| \geq 3$. Need to show $\kappa(G) \leq \kappa'(G)$.

1 Let $k = \kappa'(G)$.

2 Let $F = \{e_1, e_2, \dots, e_k\}$ be an edge cut of G where $e_i = u_i v_i$, $u_i, v_i \in V$. (Note : u_i 's and v_j 's need not be distinct.)

3 Need to construct a vertex cut from F .

4 Let $H = G \setminus \{\{u_i : 1 \leq i \leq k-1\} \cup \{u_k, v_k\}\}$

5 We retain the edge e_k in H and remove all other edges of F .

Case 1 : G is simple, connected and $|V(G)| \geq 3$. Need to show $\kappa(G) \leq \kappa'(G)$.

- 1** Let $k = \kappa'(G)$.
- 2** Let $F = \{e_1, e_2, \dots, e_k\}$ be an edge cut of G where $e_i = u_i v_i$, $u_i, v_i \in V$. (Note : u_i 's and v_j 's need not be distinct.)
- 3** Need to construct a vertex cut from F .
- 4** Let $H = G \setminus \{\{u_i : 1 \leq i \leq k-1\} \cup \{u_k, v_k\}\}$
- 5** We retain the edge e_k in H and remove all other edges of F .

Case 1 : G is simple, connected and $|V(G)| \geq 3$. Need to show $\kappa(G) \leq \kappa'(G)$.

- 1** Let $k = \kappa'(G)$.
- 2** Let $F = \{e_1, e_2, \dots, e_k\}$ be an edge cut of G where $e_i = u_i v_i$, $u_i, v_i \in V$. (Note : u_i 's and v_j 's need not be distinct.)
- 3** Need to construct a vertex cut from F .
- 4** Let $H = G \setminus \{\{u_i : 1 \leq i \leq k-1\} \cup \{u_k, v_k\}\}$
- 5** We retain the edge e_k in H and remove all other edges of F .

Case 1 : G is simple, connected and $|V(G)| \geq 3$. Need to show $\kappa(G) \leq \kappa'(G)$.

- 1** Let $k = \kappa'(G)$.
- 2** Let $F = \{e_1, e_2, \dots, e_k\}$ be an edge cut of G where $e_i = u_i v_i$, $u_i, v_i \in V$. (Note : u_i 's and v_j 's need not be distinct.)
- 3** Need to construct a vertex cut from F .
- 4** Let $H = G \setminus \{\{u_i : 1 \leq i \leq k-1\} \cup \{u_k, v_k\}\}$
- 5** We retain the edge e_k in H and remove all other edges of F .

Case 1 : G is simple, connected and $|V(G)| \geq 3$. Need to show $\kappa(G) \leq \kappa'(G)$.

- 1** Let $k = \kappa'(G)$.
- 2** Let $F = \{e_1, e_2, \dots, e_k\}$ be an edge cut of G where $e_i = u_i v_i$, $u_i, v_i \in V$. (Note : u_i 's and v_j 's need not be distinct.)
- 3** Need to construct a vertex cut from F .
- 4** Let $H = G \setminus \{\{u_i : 1 \leq i \leq k-1\} \cup \{u_k, v_k\}\}$
- 5** We retain the edge e_k in H and remove all other edges of F .

- 1 If H is already disconnected, then $\kappa(G) \leq k - 1 < \kappa'(G)$.
- 2 If H is connected, then e_k is a cut-edge of H and hence, removal of one of u_k or v_k from H makes the graph disconnected or trivial (H can be K_2).
- 3 Hence, $\kappa(G) = k = \kappa'(G)$.
- 4 In both cases, $\kappa(G) \leq \kappa'(G)$.

- 1 If H is already disconnected, then $\kappa(G) \leq k - 1 < \kappa'(G)$.
- 2 If H is connected, then e_k is a cut-edge of H and hence, removal of one of u_k or v_k from H makes the graph disconnected or trivial (H can be K_2).
- 3 Hence, $\kappa(G) = k = \kappa'(G)$.
- 4 In both cases, $\kappa(G) \leq \kappa'(G)$.

- 1 If H is already disconnected, then $\kappa(G) \leq k - 1 < \kappa'(G)$.
- 2 If H is connected, then e_k is a cut-edge of H and hence, removal of one of u_k or v_k from H makes the graph disconnected or trivial (H can be K_2).
- 3 Hence, $\kappa(G) = k = \kappa'(G)$.
- 4 In both cases, $\kappa(G) \leq \kappa'(G)$.

- 1 If H is already disconnected, then $\kappa(G) \leq k - 1 < \kappa'(G)$.
- 2 If H is connected, then e_k is a cut-edge of H and hence, removal of one of u_k or v_k from H makes the graph disconnected or trivial (H can be K_2).
- 3 Hence, $\kappa(G) = k = \kappa'(G)$.
- 4 In both cases, $\kappa(G) \leq \kappa'(G)$.

Case 2 : G is any graph.

1 Let H be an underlying simple spanning subgraph of G .

2 Then, $\kappa(H) = \kappa(G)$ and $\kappa'(H) \leq \kappa'(G)$.

3 From Case 1 we know $\kappa(H) \leq \kappa'(H)$.

4 $\implies \kappa(G) = \kappa(H) \leq \kappa'(H) \leq \kappa'(G)$. □

Hence, in all cases $\kappa(G) \leq \kappa'(G) \leq \delta(G)$.

Case 2 : G is any graph.

1 Let H be an underlying simple spanning subgraph of G .

2 Then, $\kappa(H) = \kappa(G)$ and $\kappa'(H) \leq \kappa'(G)$.

3 From Case 1 we know $\kappa(H) \leq \kappa'(H)$.

4 $\implies \kappa(G) = \kappa(H) \leq \kappa'(H) \leq \kappa'(G)$. □

Hence, in all cases $\kappa(G) \leq \kappa'(G) \leq \delta(G)$.

Case 2 : G is any graph.

1 Let H be an underlying simple spanning subgraph of G .

2 Then, $\kappa(H) = \kappa(G)$ and $\kappa'(H) \leq \kappa'(G)$.

3 From Case 1 we know $\kappa(H) \leq \kappa'(H)$.

4 $\implies \kappa(G) = \kappa(H) \leq \kappa'(H) \leq \kappa'(G)$. □

Hence, in all cases $\kappa(G) \leq \kappa'(G) \leq \delta(G)$.

Case 2 : G is any graph.

1 Let H be an underlying simple spanning subgraph of G .

2 Then, $\kappa(H) = \kappa(G)$ and $\kappa'(H) \leq \kappa'(G)$.

3 From Case 1 we know $\kappa(H) \leq \kappa'(H)$.

4 $\implies \kappa(G) = \kappa(H) \leq \kappa'(H) \leq \kappa'(G)$. □

Hence, in all cases $\kappa(G) \leq \kappa'(G) \leq \delta(G)$.

Case 2 : G is any graph.

1 Let H be an underlying simple spanning subgraph of G .

2 Then, $\kappa(H) = \kappa(G)$ and $\kappa'(H) \leq \kappa'(G)$.

3 From Case 1 we know $\kappa(H) \leq \kappa'(H)$.

4 $\implies \kappa(G) = \kappa(H) \leq \kappa'(H) \leq \kappa'(G)$. □

Hence, in all cases $\kappa(G) \leq \kappa'(G) \leq \delta(G)$.

Case 2 : G is any graph.

1 Let H be an underlying simple spanning subgraph of G .

2 Then, $\kappa(H) = \kappa(G)$ and $\kappa'(H) \leq \kappa'(G)$.

3 From Case 1 we know $\kappa(H) \leq \kappa'(H)$.


4 $\implies \kappa(G) = \kappa(H) \leq \kappa'(H) \leq \kappa'(G)$. □

Hence, in all cases $\kappa(G) \leq \kappa'(G) \leq \delta(G)$.

Example

An Example where the above inequality is strict.


$$\kappa(G) = 1 < \kappa'(G) = 2 < \delta(G) = 3$$



Example

An Example where the above inequality is strict.

$$\kappa(G) = 1 < \kappa'(G) = 2 < \delta(G) = 3$$



Definition

A maximal connected subgraph B of a graph G that has a no cut-vertex of its own is called a **block** of G .


maximal in the sense that if H is a connected subgraph of G that has no cut-vertex and $H \subseteq B$, then $H = B$

Definition


A maximal connected subgraph B of a graph G that has a no cut-vertex of its own is called a **block** of G .

maximal in the sense that if H is a connected subgraph of G that has no cut-vertex and $H \subseteq B$, then $H = B$

Example



Example



- 1 A block of a graph does not have a cut-vertex of its own. However, it may contain cut-vertices of the whole graph.
- 2 By definition, G itself is a block if it is connected and it has no cut-vertex.
- 3 Two blocks in a graph share at most one vertex; else, the two blocks together form a block, thus contradicting the maximality of blocks.
- 4 If two distinct blocks of G share a vertex v , then v is a cut-vertex of G .
- 5 Any two distinct blocks are edge disjoint; so the edge sets of blocks partition the edge set of G .
- 6 To establish a property P of a graph G , often it is enough to establish P for each of its blocks, and thereby simplify the proofs.

Observations

- 1 A block of a graph does not have a cut-vertex of its own. However, it may contain cut-vertices of the whole graph.
- 2 By definition, G itself is a block if it is connected and it has no cut-vertex.
- 3 Two blocks in a graph share at most one vertex; else, the two blocks together form a block, thus contradicting the maximality of blocks.
- 4 If two distinct blocks of G share a vertex v , then v is a cut-vertex of G .
- 5 Any two distinct blocks are edge disjoint; so the edge sets of blocks partition the edge set of G .
- 6 To establish a property P of a graph G , often it is enough to establish P for each of its blocks, and thereby simplify the proofs.

Observations

- 1** A block of a graph does not have a cut-vertex of its own. However, it may contain cut-vertices of the whole graph.
- 2** By definition, G itself is a block if it is connected and it has no cut-vertex.
- 3** Two blocks in a graph share at most one vertex; else, the two blocks together form a block, thus contradicting the maximality of blocks.
- 4** If two distinct blocks of G share a vertex v , then v is a cut-vertex of G .
- 5** Any two distinct blocks are edge disjoint; so the edge sets of blocks partition the edge set of G .
- 6** To establish a property P of a graph G , often it is enough to establish P for each of its blocks, and thereby simplify the proofs.

Observations

- 1 A block of a graph does not have a cut-vertex of its own. However, it may contain cut-vertices of the whole graph.
- 2 By definition, G itself is a block if it is connected and it has no cut-vertex.
- 3 Two blocks in a graph share at most one vertex; else, the two blocks together form a block, thus contradicting the maximality of blocks.
- 4 If two distinct blocks of G share a vertex v , then v is a cut-vertex of G .
- 5 Any two distinct blocks are edge disjoint; so the edge sets of blocks partition the edge set of G .
- 6 To establish a property P of a graph G , often it is enough to establish P for each of its blocks, and thereby simplify the proofs.

Observations

- 1 A block of a graph does not have a cut-vertex of its own. However, it may contain cut-vertices of the whole graph.
- 2 By definition, G itself is a block if it is connected and it has no cut-vertex.
- 3 Two blocks in a graph share at most one vertex; else, the two blocks together form a block, thus contradicting the maximality of blocks.
- 4 If two distinct blocks of G share a vertex v , then v is a cut-vertex of G .
- 5 Any two distinct blocks are edge disjoint; so the edge sets of blocks partition the edge set of G .
- 6 To establish a property P of a graph G , often it is enough to establish P for each of its blocks, and thereby simplify the proofs.

- 1 A block of a graph does not have a cut-vertex of its own. However, it may contain cut-vertices of the whole graph.
- 2 By definition, G itself is a block if it is connected and it has no cut-vertex.
- 3 Two blocks in a graph share at most one vertex; else, the two blocks together form a block, thus contradicting the maximality of blocks.
- 4 If two distinct blocks of G share a vertex v , then v is a cut-vertex of G .
- 5 Any two distinct blocks are edge disjoint; so the edge sets of blocks partition the edge set of G .
- 6 To establish a property P of a graph G , often it is enough to establish P for each of its blocks, and thereby simplify the proofs.

Definition

Two paths $P : x - y$ and $Q : x - y$ are said to be internally disjoint if they have no internal vertex common.

Theorem

A graph G is k -vertex connected ($1 \leq k \leq n - 1$) if and only if given any two distinct vertices u and v , there exist k internally disjoint $u - v$ paths (that is, no two of the paths have an internal vertex common).

The following result is a special case of a theorem proved by Menger (1932).

Theorem

Let G be a graph with $|V(G)| \geq 3$. Then G is a block if and only if given any two vertices x and y of G , there exist at least two internally disjoint $x - y$ -paths in G .

" \Rightarrow " Given any two vertices x, y of G , there exist at least two internally disjoint $x - y$ paths. To show that G is a block.

- 1 G is connected.
- 2 So, we have to only show that G has no cut-vertices.
- 3 On the contrary, suppose v is a cut-vertex of G .
- 4 Then by a previous Theorem there exist vertices x, y such that every $x - y$ path passes through v .
- 5 This implies that there do not exist two internally disjoint $x - y$ paths, which is a contradiction to the hypothesis.
- 6 Hence, G must be a block.

" \Rightarrow " Given any two vertices x, y of G , there exist at least two internally disjoint $x - y$ paths. To show that G is a block.

- 1 G is connected.
- 2 So, we have to only show that G has no cut-vertices.
- 3 On the contrary, suppose v is a cut-vertex of G .
- 4 Then by a previous Theorem there exist vertices x, y such that every $x - y$ path passes through v .
- 5 This implies that there do not exist two internally disjoint $x - y$ paths, which is a contradiction to the hypothesis.
- 6 Hence, G must be a block.

" \Rightarrow " Given any two vertices x, y of G , there exist at least two internally disjoint $x - y$ paths. To show that G is a block.

- 1 G is connected.
- 2 So, we have to only show that G has no cut-vertices.
- 3 On the contrary, suppose v is a cut-vertex of G .
- 4 Then by a previous Theorem there exist vertices x, y such that every $x - y$ path passes through v .
- 5 This implies that there do not exist two internally disjoint $x - y$ paths, which is a contradiction to the hypothesis.
- 6 Hence, G must be a block.

" \Rightarrow " Given any two vertices x, y of G , there exist at least two internally disjoint $x - y$ paths. To show that G is a block.

- 1 G is connected.
- 2 So, we have to only show that G has no cut-vertices.
- 3 On the contrary, suppose v is a cut-vertex of G .
- 4 Then by a previous Theorem there exist vertices x, y such that every $x - y$ path passes through v .
- 5 This implies that there do not exist two internally disjoint $x - y$ paths, which is a contradiction to the hypothesis.
- 6 Hence, G must be a block.

" \Rightarrow " Given any two vertices x, y of G , there exist at least two internally disjoint $x - y$ paths. To show that G is a block.

- 1 G is connected.
- 2 So, we have to only show that G has no cut-vertices.
- 3 On the contrary, suppose v is a cut-vertex of G .
- 4 Then by a previous Theorem there exist vertices x, y such that every $x - y$ path passes through v .
- 5 This implies that there do not exist two internally disjoint $x - y$ paths, which is a contradiction to the hypothesis.
- 6 Hence, G must be a block.

" \Rightarrow " Given any two vertices x, y of G , there exist at least two internally disjoint $x - y$ paths. To show that G is a block.

- 1 G is connected.
- 2 So, we have to only show that G has no cut-vertices.
- 3 On the contrary, suppose v is a cut-vertex of G .
- 4 Then by a previous Theorem there exist vertices x, y such that every $x - y$ path passes through v .
- 5 This implies that there do not exist two internally disjoint $x - y$ paths, which is a contradiction to the hypothesis.
- 6 Hence, G must be a block.

" \Rightarrow " Given any two vertices x, y of G , there exist at least two internally disjoint $x - y$ paths. To show that G is a block.

- 1 G is connected.
- 2 So, we have to only show that G has no cut-vertices.
- 3 On the contrary, suppose v is a cut-vertex of G .
- 4 Then by a previous Theorem there exist vertices x, y such that every $x - y$ path passes through v .
- 5 This implies that there do not exist two internally disjoint $x - y$ paths, which is a contradiction to the hypothesis.
- 6 Hence, G must be a block.

" \Leftarrow " Given G is a block. To show that given any two vertices x, y of G , there exist at least two internally disjoint $x - y$ paths.

Proof by induction on $d(x, y)$.

Base Case :

- 1 $d(x, y) = 1 \Rightarrow x$ and y are adjacent.
- 2 Let $e = xy$ the edge between x and y .
- 3 $G - e$ is connected. If not e is a cut edge and one of x or y is a cut-vertex of G .
- 4 A contradiction as G is a block.
- 5 Hence, there exists $P = x - y$ path in $G - e$.
- 6 $P = x - y$ in $G - e$ and $e = xy$ are two internally disjoint $x - y$ paths in G .

" \Leftarrow " Given G is a block. To show that given any two vertices x, y of G , there exist at least two internally disjoint $x - y$ paths.

Proof by induction on $d(x, y)$.

Base Case :

- 1 $d(x, y) = 1 \Rightarrow x$ and y are adjacent.
- 2 Let $e = xy$ the edge between x and y .
- 3 $G - e$ is connected. If not e is a cut edge and one of x or y is a cut-vertex of G .
- 4 A contradiction as G is a block.
- 5 Hence, there exists $P = x - y$ path in $G - e$.
- 6 $P = x - y$ in $G - e$ and $e = xy$ are two internally disjoint $x - y$ paths in G .

" \Leftarrow " Given G is a block. To show that given any two vertices x, y of G , there exist at least two internally disjoint $x - y$ paths.

Proof by induction on $d(x, y)$.

Base Case :

- 1 $d(x, y) = 1 \Rightarrow x$ and y are adjacent.
- 2 Let $e = xy$ the edge between x and y .
- 3 $G - e$ is connected. If not e is a cut edge and one of x or y is a cut-vertex of G .
- 4 A contradiction as G is a block.
- 5 Hence, there exists $P = x - y$ path in $G - e$.
- 6 $P = x - y$ in $G - e$ and $e = xy$ are two internally disjoint $x - y$ paths in G .

" \Leftarrow " Given G is a block. To show that given any two vertices x, y of G , there exist at least two internally disjoint $x - y$ paths.

Proof by induction on $d(x, y)$.

Base Case :

- 1 $d(x, y) = 1 \Rightarrow x$ and y are adjacent.
- 2 Let $e = xy$ the edge between x and y .
- 3 $G - e$ is connected. If not e is a cut edge and one of x or y is a cut-vertex of G .
- 4 A contradiction as G is a block.
- 5 Hence, there exists $P = x - y$ path in $G - e$.
- 6 $P = x - y$ in $G - e$ and $e = xy$ are two internally disjoint $x - y$ paths in G .

" \Leftarrow " Given G is a block. To show that given any two vertices x, y of G , there exist at least two internally disjoint $x - y$ paths.

Proof by induction on $d(x, y)$.

Base Case :

- 1 $d(x, y) = 1 \Rightarrow x$ and y are adjacent.
- 2 Let $e = xy$ the edge between x and y .
- 3 $G - e$ is connected. If not e is a cut edge and one of x or y is a cut-vertex of G .
- 4 A contradiction as G is a block.
- 5 Hence, there exists $P = x - y$ path in $G - e$.
- 6 $P = x - y$ in $G - e$ and $e = xy$ are two internally disjoint $x - y$ paths in G .

" \Leftarrow " Given G is a block. To show that given any two vertices x, y of G , there exist at least two internally disjoint $x - y$ paths.

Proof by induction on $d(x, y)$.

Base Case :

- 1 $d(x, y) = 1 \Rightarrow x$ and y are adjacent.
- 2 Let $e = xy$ the edge between x and y .
- 3 $G - e$ is connected. If not e is a cut edge and one of x or y is a cut-vertex of G .
- 4 A contradiction as G is a block.
- 5 Hence, there exists $P = x - y$ path in $G - e$.
- 6 $P = x - y$ in $G - e$ and $e = xy$ are two internally disjoint $x - y$ paths in G .

" \Leftarrow " Given G is a block. To show that given any two vertices x, y of G , there exist at least two internally disjoint $x - y$ paths.

Proof by induction on $d(x, y)$.

Base Case :

- 1 $d(x, y) = 1 \Rightarrow x$ and y are adjacent.
- 2 Let $e = xy$ the edge between x and y .
- 3 $G - e$ is connected. If not e is a cut edge and one of x or y is a cut-vertex of G .
- 4 A contradiction as G is a block.
- 5 Hence, there exists $P = x - y$ path in $G - e$.
- 6 $P = x - y$ in $G - e$ and $e = xy$ are two internally disjoint $x - y$ paths in G .

" \Leftarrow " Given G is a block. To show that given any two vertices x, y of G , there exist at least two internally disjoint $x - y$ paths.

Proof by induction on $d(x, y)$.

Base Case :

- 1 $d(x, y) = 1 \Rightarrow x$ and y are adjacent.
- 2 Let $e = xy$ the edge between x and y .
- 3 $G - e$ is connected. If not e is a cut edge and one of x or y is a cut-vertex of G .
- 4 A contradiction as G is a block.
- 5 Hence, there exists $P = x - y$ path in $G - e$.
- 6 $P = x - y$ in $G - e$ and $e = xy$ are two internally disjoint $x - y$ paths in G .

Induction Hypothesis : Assume true for all vertices with $d(x, y) \leq k$.

Induction Step : Let $d(x, y) = k + 1$.

- 1 Let $P : x - y$ be a shortest $x - y$ path.
- 2 so length of $P = d(x, y)$.
- 3 Let w be a vertex that precedes y in P .
- 4 So, $d(x, w) = d(x, y) - 1$.
- 5 Hence, by induction hypothesis, there exist two internally disjoint $x - w$ paths, say $Q : x - w$ and $R : x - w$.

Induction Hypothesis : Assume true for all vertices with $d(x, y) \leq k$.

Induction Step : Let $d(x, y) = k + 1$.

- 1** Let $P : x - y$ be a shortest $x - y$ path.
- 2 so length of $P = d(x, y)$.
- 3 Let w be a vertex that precedes y in P .
- 4 So, $d(x, w) = d(x, y) - 1$.
- 5 Hence, by induction hypothesis, there exist two internally disjoint $x - w$ paths, say $Q : x - w$ and $R : x - w$.

Induction Hypothesis : Assume true for all vertices with $d(x, y) \leq k$.

Induction Step : Let $d(x, y) = k + 1$.

- 1** Let $P : x - y$ be a shortest $x - y$ path.
- 2** so length of $P = d(x, y)$.
- 3** Let w be a vertex that precedes y in P .
- 4** So, $d(x, w) = d(x, y) - 1$.
- 5** Hence, by induction hypothesis, there exist two internally disjoint $x - w$ paths, say $Q : x - w$ and $R : x - w$.

Induction Hypothesis : Assume true for all vertices with $d(x, y) \leq k$.

Induction Step : Let $d(x, y) = k + 1$.

- 1** Let $P : x - y$ be a shortest $x - y$ path.
- 2** so length of $P = d(x, y)$.
- 3** Let w be a vertex that precedes y in P .
- 4** So, $d(x, w) = d(x, y) - 1$.
- 5** Hence, by induction hypothesis, there exist two internally disjoint $x - w$ paths, say $Q : x - w$ and $R : x - w$.

Induction Hypothesis : Assume true for all vertices with $d(x, y) \leq k$.

Induction Step : Let $d(x, y) = k + 1$.

- 1** Let $P : x - y$ be a shortest $x - y$ path.
- 2** so length of $P = d(x, y)$.
- 3** Let w be a vertex that precedes y in P .
- 4** So, $d(x, w) = d(x, y) - 1$.
- 5** Hence, by induction hypothesis, there exist two internally disjoint $x - w$ paths, say $Q : x - w$ and $R : x - w$.

Induction Hypothesis : Assume true for all vertices with $d(x, y) \leq k$.


Induction Step : Let $d(x, y) = k + 1$.

- 1** Let $P : x - y$ be a shortest $x - y$ path.
- 2** so length of $P = d(x, y)$.
- 3** Let w be a vertex that precedes y in P .
- 4** So, $d(x, w) = d(x, y) - 1$.
- 5** Hence, by induction hypothesis, there exist two internally disjoint $x - w$ paths, say $Q : x - w$ and $R : x - w$.

Proof Contd

- 1 Since G is a block, w is not a cut-vertex.
- 2 So, there exists a path $S : x - y$ in $G - w$.
- 3 Let z be the last vertex in S that lies in $V(Q) \cap V(R)$.
- 4 For definiteness, let $z \in V(Q)$. Then $Q : x - z$, $S : z - y$ and $R : x - w + wy$ are two internally disjoint $x - y$ paths.


□



Proof Contd

- 1 Since G is a block, w is not a cut-vertex.
- 2 So, there exists a path $S : x - y$ in $G - w$.
- 3 Let z be the last vertex in S that lies in $V(Q) \cap V(R)$.
- 4 For definiteness, let $z \in V(Q)$. Then $Q : x - z$, $S : z - y$ and $R : x - w + wy$ are two internally disjoint $x - y$ paths.


□



Proof Contd


- 1 Since G is a block, w is not a cut-vertex.
- 2 So, there exists a path $S : x - y$ in $G - w$.
- 3 Let z be the last vertex in S that lies in $V(Q) \cap V(R)$.
- 4 For definiteness, let $z \in V(Q)$. Then $Q : x - z$, $S : z - y$ and $R : x - w + wy$ are two internally disjoint $x - y$ paths.

□



Proof Contd

- 1 Since G is a block, w is not a cut-vertex.
- 2 So, there exists a path $S : x - y$ in $G - w$.
- 3 Let z be the last vertex in S that lies in $V(Q) \cap V(R)$.
- 4 For definiteness, let $z \in V(Q)$. Then $Q : x - z$, $S : z - y$ and $R : x - w + wy$ are two internally disjoint $x - y$ paths. □




Theorem

A graph G is k -edge connected if and only if given any two distinct vertices u and v , there exists k edge-disjoint $u - v$ paths (that is, no two paths have an edge common).

There are many sufficient conditions for a graph to be k -vertex-connected or k -edge-connected. (In advance course on GT).

Example




1 $\kappa(G) = 2$ and $\kappa'(G) = 2$

2 The given graph is 2-connected and 2-edge connected.

3 Between any two vertices there exists two internally disjoint paths and two edge disjoint paths.

Example



- 1** $\kappa(G) = 2$ and $\kappa'(G) = 2$
- 2** The given graph is 2-connected and 2-edge connected.
- 3** Between any two vertices there exists two internally disjoint paths and two edge disjoint paths.