

Bayesian statistics with R

3. Analyses by hand

Olivier Gimenez

March 2021

Back to Bayes

A simple example

- Let us take a simple example to fix ideas.
- 120 deer were radio-tracked over winter.
- 61 close to a plant, 59 far from any human activity.
- Question: is there a treatment effect on survival?

	Released	Alive	Dead	Other
treatment	61	19	38	4
control	59	21	38	0

- So, $n = 57$ deer were assigned to the treatment group of which $k = 19$ survived the winter.

- So, $n = 57$ deer were assigned to the treatment group of which $k = 19$ survived the winter.
- Of interest is the probability of over-winter survival, call it θ , for the general population within the treatment area.

- So, $n = 57$ deer were assigned to the treatment group of which $k = 19$ survived the winter.
- Of interest is the probability of over-winter survival, call it θ , for the general population within the treatment area.
- The obvious estimate is simply to take the ratio $k/n = 19/57$.

- So, $n = 57$ deer were assigned to the treatment group of which $k = 19$ survived the winter.
- Of interest is the probability of over-winter survival, call it θ , for the general population within the treatment area.
- The obvious estimate is simply to take the ratio $k/n = 19/57$.
- How would the classical statistician justify this estimate?

- Our model is that we have a Binomial experiment (assuming independent and identically distributed draws from the population).

- Our model is that we have a Binomial experiment (assuming independent and identically distributed draws from the population).
- K the number of alive individuals at the end of the winter, so that $P(K = k) = \binom{n}{k} \theta^k (1 - \theta)^{n-k}$.

- Our model is that we have a Binomial experiment (assuming independent and identically distributed draws from the population).
- K the number of alive individuals at the end of the winter, so that $P(K = k) = \binom{n}{k} \theta^k (1 - \theta)^{n-k}$.
- The classical approach is to maximise the corresponding likelihood with respect to θ to obtain the entirely plausible MLE:

$$\hat{\theta} = k/n = 19/57$$

The Bayesian approach

- The Bayesian starts off with a prior.

The Bayesian approach

- The Bayesian starts off with a prior.
- Now, the one thing we know about θ is that is a continuous random variable and that it lies between zero and one.

The Bayesian approach

- The Bayesian starts off with a prior.
- Now, the one thing we know about θ is that is a continuous random variable and that it lies between zero and one.
- Thus, a suitable prior distribution might be the Beta defined on $[0, 1]$.







The Bayesian approach

- The Bayesian starts off with a prior.
- Now, the one thing we know about θ is that is a continuous random variable and that it lies between zero and one.
- Thus, a suitable prior distribution might be the Beta defined on $[0, 1]$.
- What is the Beta distribution?

What is the Beta distribution?

$$q(\theta \mid \alpha, \beta) = \frac{1}{\text{Beta}(\alpha, \beta)} \theta^{\alpha-1} (1 - \theta)^{\beta-1}$$

with $\text{Beta}(\alpha, \beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha + \beta)}$ and $\Gamma(n) = (n - 1)!$

beta(1,1)**beta(2,1)****beta(1,2)****beta(2,2)****beta(10,10)****beta(0.8,0.8)**

The Bayesian approach

- We assume a priori that $\theta \sim \text{Beta}(a, b)$ so that $\Pr(\theta) = \theta^{a-1}(1 - \theta)^{b-1}$

The Bayesian approach

- We assume a priori that $\theta \sim \text{Beta}(a, b)$ so that $\Pr(\theta) = \theta^{a-1}(1 - \theta)^{b-1}$
- Then we have:

$$\begin{aligned} \Pr(\theta \mid k) &\propto \binom{n}{k} \theta^k (1 - \theta)^{n-k} \theta^{a-1} (1 - \theta)^{b-1} \\ &\propto \theta^{(a+k)-1} (1 - \theta)^{(b+n-k)-1} \end{aligned}$$

The Bayesian approach

- We assume a priori that $\theta \sim \text{Beta}(a, b)$ so that $\Pr(\theta) = \theta^{a-1}(1 - \theta)^{b-1}$
- Then we have:

$$\begin{aligned} \textcolor{red}{Pr}(\theta \mid k) &\propto \binom{\textcolor{blue}{n}}{\textcolor{blue}{k}} \theta^{\textcolor{blue}{k}} (1 - \theta)^{\textcolor{blue}{n} - \textcolor{blue}{k}} \textcolor{green}{\theta^{a-1}} (1 - \theta)^{\textcolor{green}{b-1}} \\ &\propto \theta^{(a+k)-1} (1 - \theta)^{(b+n-k)-1} \end{aligned}$$

- That is:

$$\theta \mid k \sim \text{Beta}(a + k, b + n - k)$$

The Bayesian approach

- We assume a priori that $\theta \sim \text{Beta}(a, b)$ so that $\Pr(\theta) = \theta^{a-1}(1 - \theta)^{b-1}$
- Then we have:

$$\begin{aligned} \Pr(\theta \mid k) &\propto \binom{n}{k} \theta^k (1 - \theta)^{n-k} \theta^{a-1} (1 - \theta)^{b-1} \\ &\propto \theta^{(a+k)-1} (1 - \theta)^{(b+n-k)-1} \end{aligned}$$

- That is:

$$\theta \mid k \sim \text{Beta}(a + k, b + n - k)$$

- Take a Beta prior with a Binomial likelihood, you get a Beta posterior (conjugacy)

Application to the deer example

- Posterior distribution of survival is $\theta \sim \text{Beta}(a + k, b + n - k)$.

Application to the deer example

- Posterior distribution of survival is $\theta \sim \text{Beta}(a + k, b + n - k)$.
- If we take a Uniform prior, i.e. $\text{Beta}(1, 1)$, then we have:

Application to the deer example

- Posterior distribution of survival is $\theta \sim \text{Beta}(a + k, b + n - k)$.
- If we take a Uniform prior, i.e. $\text{Beta}(1, 1)$, then we have:
- $\theta_{\text{treatment}} \sim \text{Beta}(1 + 19, 1 + 57 - 19) = \text{Beta}(20, 39)$

Application to the deer example

- Posterior distribution of survival is $\theta \sim \text{Beta}(a + k, b + n - k)$.
- If we take a Uniform prior, i.e. $\text{Beta}(1, 1)$, then we have:
- $\theta_{\text{treatment}} \sim \text{Beta}(1 + 19, 1 + 57 - 19) = \text{Beta}(20, 39)$
- Note that in this specific situation, the posterior has an explicit expression, easy to manipulate.


Application to the deer example

- Posterior distribution of survival is $\theta \sim \text{Beta}(a + k, b + n - k)$.
- If we take a Uniform prior, i.e. $\text{Beta}(1, 1)$, then we have:
- $\theta_{\text{treatment}} \sim \text{Beta}(1 + 19, 1 + 57 - 19) = \text{Beta}(20, 39)$
- Note that in this specific situation, the posterior has an explicit expression, easy to manipulate.
- In particular, $E(\text{Beta}(a, b)) = \frac{a}{a + b} = 20/59$ to be compared with the MLE $19/57$.


A general result

This is a general result, the Bayesian and frequentist estimates will always agree if there is sufficient data, so long as the likelihood is not explicitly ruled out by the prior.

Prior $Beta(1, 1)$ and posterior survival $Beta(20, 39)$



Prior $Beta(1, 1)$ and posterior survival $Beta(20, 39)$



Our model so far

$$y \sim \text{Binomial}(N, \theta)$$

[likelihood]

$$\theta \sim \text{Beta}(1, 1)$$

[prior for θ]

