


Solution: Backpropagation

## Backpropagation: a simple example

$$f(x, y, z) = (x + y)z$$

## Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$



## Backpropagation: a simple example

$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2$ ,  $y = 5$ ,  $z = -4$




## Backpropagation: a simple example

$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2$ ,  $y = 5$ ,  $z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$




## Backpropagation: a simple example

$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2$ ,  $y = 5$ ,  $z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$



## Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



## Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial f}$$



## Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial f}$$

## Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial z}$$

## Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial z}$$

## Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial q}$$

## Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



# Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



Chain rule:

$$\frac{\partial f}{\partial y} = \frac{\partial f}{\partial q} \frac{\partial q}{\partial y}$$

Upstream  
gradient

Local  
gradient

$$\frac{\partial f}{\partial y}$$

## Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial y}$$

Chain rule:

$$\frac{\partial f}{\partial y} = \frac{\partial f}{\partial q} \frac{\partial q}{\partial y}$$

Upstream  
gradient

Local  
gradient

## Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial x}$$

Chain rule:

$$\frac{\partial f}{\partial x} = \frac{\partial f}{\partial q} \frac{\partial q}{\partial x}$$

Upstream  
gradient

Local  
gradient



## Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$




$$\frac{\partial f}{\partial x}$$


Chain rule:


$$\frac{\partial f}{\partial x} = \frac{\partial f}{\partial q} \frac{\partial q}{\partial x}$$


Upstream  
gradient


Local  
gradient
















Another example:  $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$






Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$




Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



Another example:


$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$f(x) = e^x$	$\rightarrow$	$\frac{df}{dx} = e^x$		$f(x) = \frac{1}{x}$	$\rightarrow$	$\frac{df}{dx} = -1/x^2$
$f_a(x) = ax$	$\rightarrow$	$\frac{df}{dx} = a$		$f_c(x) = c + x$	$\rightarrow$	$\frac{df}{dx} = 1$

Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$$f(x) = e^x$$

→

$$\frac{df}{dx} = e^x$$

$$f_a(x) = ax$$

→

$$\frac{df}{dx} = a$$

$$f(x) = \frac{1}{x}$$

→

$$\frac{df}{dx} = -1/x^2$$


$$f_c(x) = c + x$$

→

$$\frac{df}{dx} = 1$$

Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$$f(x) = e^x$$

→

$$\frac{df}{dx} = e^x$$

$$f_a(x) = ax$$

→

$$\frac{df}{dx} = a$$

$$f(x) = \frac{1}{x}$$

→

$$\frac{df}{dx} = -1/x^2$$


$$f_c(x) = c + x$$

→

$$\frac{df}{dx} = 1$$

Another example:


$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$f(x) = e^x$	$\rightarrow$	$\frac{df}{dx} = e^x$		$f(x) = \frac{1}{x}$	$\rightarrow$	$\frac{df}{dx} = -1/x^2$
$f_a(x) = ax$	$\rightarrow$	$\frac{df}{dx} = a$		$f_c(x) = c + x$	$\rightarrow$	$\frac{df}{dx} = 1$

Another example:


$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$f(x) = e^x$	$\rightarrow$	$\frac{df}{dx} = e^x$		$f(x) = \frac{1}{x}$	$\rightarrow$	$\frac{df}{dx} = -1/x^2$
$f_a(x) = ax$	$\rightarrow$	$\frac{df}{dx} = a$		$f_c(x) = c + x$	$\rightarrow$	$\frac{df}{dx} = 1$

Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$$f(x) = e^x$$

$\rightarrow$

$$\frac{df}{dx} = e^x$$

$$f_a(x) = ax$$

$\rightarrow$

$$\frac{df}{dx} = a$$

$$f(x) = \frac{1}{x}$$

$\rightarrow$

$$\frac{df}{dx} = -1/x^2$$

$$f_c(x) = c + x$$


$\rightarrow$

$$\frac{df}{dx} = 1$$



Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$$f(x) = e^x$$

→

$$\frac{df}{dx} = e^x$$

$$f_a(x) = ax$$

→

$$\frac{df}{dx} = a$$

$$f(x) = \frac{1}{x}$$

→

$$\frac{df}{dx} = -1/x^2$$


$$f_c(x) = c + x$$

→

$$\frac{df}{dx} = 1$$

Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$$f(x) = e^x \rightarrow \frac{df}{dx} = e^x$$


$$f_a(x) = ax \rightarrow \frac{df}{dx} = a$$

$$f(x) = \frac{1}{x} \rightarrow \frac{df}{dx} = -1/x^2$$

$$f_c(x) = c + x \rightarrow \frac{df}{dx} = 1$$

Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$$f(x) = e^x \quad \rightarrow \quad \frac{df}{dx} = e^x$$


$$f_a(x) = ax \quad \rightarrow \quad \frac{df}{dx} = a$$

$$f(x) = \frac{1}{x} \quad \rightarrow \quad \frac{df}{dx} = -1/x^2$$

$$f_c(x) = c + x \quad \rightarrow \quad \frac{df}{dx} = 1$$

Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$$f(x) = e^x$$

→

$$\frac{df}{dx} = e^x$$

$$f_a(x) = ax$$

→

$$\frac{df}{dx} = a$$

$$f(x) = \frac{1}{x}$$

→

$$\frac{df}{dx} = -1/x^2$$


$$f_c(x) = c + x$$

→

$$\frac{df}{dx} = 1$$

Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



[upstream gradient] x [local gradient]

$$[0.2] \times [1] = 0.2$$

$$[0.2] \times [1] = 0.2 \text{ (both inputs!)}$$

$$f(x) = e^x$$

→

$$\frac{df}{dx} = e^x$$

$$f_a(x) = ax$$

→

$$\frac{df}{dx} = a$$

$$f(x) = \frac{1}{x}$$

→

$$\frac{df}{dx} = -1/x^2$$


$$f_c(x) = c + x$$

→

$$\frac{df}{dx} = 1$$

Another example:


$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$f(x) = e^x$	$\rightarrow$	$\frac{df}{dx} = e^x$		$f(x) = \frac{1}{x}$	$\rightarrow$	$\frac{df}{dx} = -1/x^2$
$f_a(x) = ax$	$\rightarrow$	$\frac{df}{dx} = a$		$f_c(x) = c + x$	$\rightarrow$	$\frac{df}{dx} = 1$

Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



[upstream gradient] x [local gradient]

$$w_0: [0.2] \times [-1] = -0.2$$

$$x_0: [0.2] \times [2] = 0.4$$

$$f(x) = e^x$$

→

$$\frac{df}{dx} = e^x$$

$$f_a(x) = ax$$

→

$$\frac{df}{dx} = a$$

$$f(x) = \frac{1}{x}$$

→

$$\frac{df}{dx} = -1/x^2$$

$$f_c(x) = c + x$$


→

$$\frac{df}{dx} = 1$$

Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$

Computational graph representation may not be unique. Choose one where local gradients at each node can be easily expressed!






Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2x_2)}}$$

Computational graph representation may not be unique. Choose one where local gradients at each node can be easily expressed!




Sigmoid local gradient:

$$\frac{d\sigma(x)}{dx} = \frac{e^{-x}}{(1 + e^{-x})^2} = \left( \frac{1 + e^{-x} - 1}{1 + e^{-x}} \right) \left( \frac{1}{1 + e^{-x}} \right) = (1 - \sigma(x)) \sigma(x)$$

Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2x_2)}}$$

Computational graph representation may not be unique. Choose one where local gradients at each node can be easily expressed!



Sigmoid function


$$\sigma(x) = \frac{1}{1 + e^{-x}}$$

[upstream gradient] x [local gradient]  
 $[1.00] \times [(1 - 1/(1+e^{-1})) (1/(1+e^{-1}))] = 0.2$


Sigmoid local gradient:

$$\frac{d\sigma(x)}{dx} = \frac{e^{-x}}{(1 + e^{-x})^2} = \left( \frac{1 + e^{-x} - 1}{1 + e^{-x}} \right) \left( \frac{1}{1 + e^{-x}} \right) = (1 - \sigma(x)) \sigma(x)$$


# Backprop with Vectors



# Backprop with Vectors



# Backprop with Vectors



# Backprop with Vectors

