

COVID-19 Projections for AR

UA Research Team

March 25, 2020

Contents


1	COVID-19 outbreak in AR	1
1.1	Plot by HRR on March 22	2
1.2	Spatial distribution of confirmed cases over county and HRR	2
1.3	eSIR model	2
1.4	HRR-level forecasts	3
1.5	P(Infection) plots for each HRR	4
1.6	Capacity crossover points for the mean case	5
1.7	Capacity crossover points for the worst case	7
1.8	14-days rolling mean for mean case and worst case	8
2	Appendix	10
2.1	P(removal) plots for each HRR	10
2.2	Parameter choices	11
3	References	11

1 COVID-19 outbreak in AR

The main aim of this project is to analyze the dynamics of outbreak of novel coronavirus using a hierarchical epidemiological model to forecast the spread of infection and its impact on hospitalization across the state and hospital referral regions (HRR). This is a short report of our results so far.


1.1 Plot by HRR on March 22

Confirmed Covid-19 cases by HRR



<https://usafacts.org/visualizations/coronavirus-covid-19-spread-map/>

1.2 Spatial distribution of confirmed cases over county and HRR



1.3 eSIR model

The standard SIR model [see 3] takes as input two time series of proportions of infected and removed cases, denoted by Y_t^I and Y_t^R at time t , respectively, where the removed part of the population is a sum of the recovered cases and deaths at time t . The standard SIR assumes a constant rate for both transmission and removal which does not reflect reality where these rates are affected by government imposed isolation measures. We apply an extension to the standard SIR model (eSIR, see [1,2]) by letting a time-varying probability that a susceptible person meets an infected person or vice versa. The dynamic model is represented below:



$$\frac{d\theta_t^S}{dt} = -\beta\pi(t)\theta_t^S\theta_t^I, \quad \frac{d\theta_t^I}{dt} = \beta\pi(t)\theta_t^S\theta_t^I - \gamma\theta_t^I, \quad \frac{d\theta_t^R}{dt} = \gamma\theta_t^I$$

Here θ_t^I and θ_t^R are the respective prevalence of infection and removal at time t , and expected proportions are equal to the prevalence of infection and the probability of removal at time t .

Here $\pi(t)$ is called a transmission modifier as it modifies the probability of a susceptible person meeting an infected person. In particular, Suppose at a time t , $q^S(t)$ is the chance of an at-risk person being in-home isolation, and $q^R(t)$ is the chance of an infected person being in-hospital quarantine, then, we will have:

$$\pi(t) = (1 - q^S(t))(1 - q^R(t)) \in (0, 1).$$

Note that the above extended SIR model assumes primarily that *both population-level chance of being susceptible and population-level chance of being infected remain the same, but the chance of a susceptible person meeting with an infected person is reduced by $\pi(t)$.*

For a real data analysis, $\pi(t)$, the transmission rate modifier needs to be specified according to isolation measures taken in a given region. We can assume the following:

$$\pi(t) = \begin{cases} \pi_{01} & \text{Before March 12, no isolation} \\ \pi_{02} & \text{March 12-19, in-person class stops} \\ \pi_{03} & \text{March 19 onwards, mandatory work-from-home at UA} \end{cases}$$


Obviously, different values of π_{0i} leads to different dynamic behaviours; we chose $\pi_{02} = 0.9$ and $\pi_{03} = 0.8$ as that produces good three-steps ahead forecasts.


1.4 HRR-level forecasts

State-level data from 01/22 to 03/23 pulled from USA facts and map them to the HRRs. The model forecasts the probability of infection and removal, over 200 days from the starting point (1/22), i.e. end date is Aug 10.

The forecasting plots of the prevalence of infected and removed compartments are given in Section 1.3 and in 2.1 (appendix). In each of these plots, the observed proportions are denoted by the black dots left to the blue vertical line, which represents the last date of observations, or time t_0 (the current time up to which we have observed data). The two turning points are the green and purple vertical lines. The first turning point is where the daily increase of $P(I)$ starts decreasing, and the second turning point is where the $P(R)$ overtakes $P(I)$. The 95% credible interval of the predicted proportions of infected and removals are given by the salmon colored area. The relatively large area under these curves denote the large uncertainty in forecasting far ahead in time based on limited available data. The gray and red curves are the posterior mean and median curves for the prevalence of the infected θ_t^I in section 1.3, and prevalence of the removed θ_t^R in section 2.1.

1.5 P(Infection) plots for each HRR





Next, we use a 14-day moving average of the probability of infection to account for the average length of stay for Covid-19 infected patients. Let us denote by $\hat{\theta}_t^I$ as the moving average of the prevalence of infected individuals $\hat{\theta}_t^I$, $t = 1, \dots, 14$. The predicted value of number of affected individuals is going to be the number of susceptible times the prevalence of infected individuals, i.e.

$$\hat{N}^I = N \times \hat{\theta}_t^S \times \hat{\theta}_t^I.$$

The upper credible limit and the mean are very different on the figures above, representing the uncertainty involved in forecasting with limited data. We show the estimated number of infected individuals where $\hat{\theta}_t^I$ is taken to be the posterior mean and the upper credible limit of the posterior distribution of θ_t^I . The following table shows the maximum value of the expected number of infected and the date at which the maximum value is attained.

Table 1: Worst case and Mean case Predictions from the eSIR model by each HRR				
HRR	max total worstcase	max day worstcase	max total meancase	max day meancase
16	48095.62515	7/8/2020	8936.499409	8/8/2020
18	37295.5838	6/25/2020	9103.493193	7/3/2020
19	257313.8573	6/11/2020	64634.96703	6/26/2020
21	81749.43015	7/3/2020	17385.77938	7/27/2020
22	36515.80452	7/5/2020	7932.597184	7/26/2020
270	136931.2093	6/17/2020	31044.57732	7/6/2020
379	258568.0131	6/24/2020	57657.41115	7/11/2020

%


1.6 Capacity crossover points for the mean case




The next set of plots show the prevalence of infection with projected hospitalizations and ICU admissions. Our assumption is 1/5 of infected will be hospitalized and 1/5 of those hospitalized will need ICU facilities, based on the current HGHI proportions.

%

Table 2: Dates when the projected hospitalization and ICU bed demands will cross the available hospital beds and ICU beds per HRR



HRR	mean case hospital cross date	mean case icu cross date	worst case hospital cross date	worst case icu cross date
16	5/31/2020	5/21/2020	5/12/2020	5/6/2020
18	5/11/2020	4/28/2020	4/28/2020	4/18/2020
19	5/4/2020	4/26/2020	4/23/2020	4/17/2020
21	5/19/2020	5/12/2020	5/6/2020	5/1/2020
22	5/21/2020	5/18/2020	5/5/2020	5/3/2020
270	5/5/2020	4/27/2020	4/24/2020	4/17/2020
379	5/10/2020	4/29/2020	4/28/2020	4/20/2020






1.7 Capacity crossover points for the worst case

Same as before but worst case instead of mean.







1.8 14-days rolling mean for mean case and worst case

These plots show how the mean and the worst case differ.


14-Days Rolling Mean (Worst vs. Mean case)




14-Days Rolling Mean (Worst vs. Mean case)




14-Days Rolling Mean (Worst vs. Mean case)




14-Days Rolling Mean (Worst vs. Mean case)




14-Days Rolling Mean (Worst vs. Mean case)



14-Days Rolling Mean (Worst vs. Mean case)








14-Days Rolling Mean (Worst vs. Mean case)



2 Appendix

2.1 P(removal) plots for each HRR





2.2 Parameter choices

The output is dependent on the parameter choices, viz. death_in_R, beta0, gamma0 and R0. Here's what they mean and what we can do about them.

- **death_in_R**: the numeric value of average of cumulative deaths in the removed compartments. (I used 0 as right now we don't have info on this for AR, but can use values from other states and see what changes.)
- **beta0**: the hyperparameter β_0 of average transmission rate, the default is the one estimated from the SARS first-month outbreak (0.2586).
- **gamma0**: the hyperparameter γ_0 of average removed rate, the default is the one estimated from the SARS first-month outbreak (0.0821).
- **R0**: the hyperparameter of the mean reproduction number R_0 . The default is thus the ratio of beta0/gamma0, which can be specified directly. (Our mean R_0 is 3.15). R_0 is highly important, and a different R_0 would give different results.

3 References

1. Song, P. X., Wang, L., Zhou, Y., He, J., Zhu, B., Wang, F., ... & Eisenberg, M. (2020). An epidemiological forecast model and software assessing interventions on COVID-19 epidemic in China.

medRxiv. <https://www.medrxiv.org/content/medrxiv/early/2020/03/03/2020.02.29.20029421.full.pdf>

2. Song Lab at UM. eSIR: Extended state-space SIR models, 2020. R package version 0.1.0.
3. Dave Osthus, Kyle S Hickmann, Petrutia C Caragea, Dave Higdon, and Sara Y Del Valle. Forecasting seasonal influenza with a state-space sir model. *The Annals of Applied Statistics*, 11(1):202, 2017.