lecture 10 - Kleene Theorem III

Closures:- γ_2 FA $\quad$ γ_1^* FA^*.

→ When the final state is encountered it will be combined with Initial.

→ For the first time, the Initial will also be considered as final.



old states	Transition at 'a'	Transition at 'b'
$^+\bar{z}_1 \equiv q_0$	$z_2 \equiv q_1$	$z_3 \equiv q_3$
$z_2 \equiv q_1$	$z_4^+ \equiv (q_2, q_0)$	$z_5^+ \equiv (q_4, q_0)$
$z_3 \equiv q_3$	$z_2 \equiv q_1$	$z_5^+ \equiv (q_4, q_0)$.
$z_4^+ \equiv (q_2, q_0)$.	$z_6^+ \equiv (q_2, q_0, q_1)$	$z_3 \equiv (q_3, \cancel{q_3})$
$z_5^+ \equiv (q_4, q_0)$.	$z_2 \equiv (q_1, \cancel{q_1})$	$z_7^+ \equiv (q_4, q_0, q_3)$.
$z_6^+ \equiv (q_2, q_0, q_1)$.	$z_6^+ \equiv (q_2, q_0, q_1, \cancel{q_2}, \cancel{q_0})$	$z_7^+ \equiv (q_3, \cancel{q_3}, q_4, q_0)$
$z_7^+ \equiv (q_4, q_0, q_3)$	$z_2 \equiv (q_1, \cancel{q_1}, \cancel{q_1})$	$z_7^+ \equiv (q_4, q_0, q_3, \cancel{q_4}, \cancel{q_0})$.



old state	Transition at 'a'	Transition at 'b'.
$^-z_1^+ \equiv q_0$	$z_2 \equiv q_2$	$z_2 \equiv q_1$
$z_2 \equiv q_1$	$z_3^+ \equiv q_0, \cancel{q_0}$	$z_3^+ \equiv q_0, \cancel{q_0}$
$z_2^+ = 0.0$	$z_2 \equiv q_1$	$z_2 \equiv q_1$

$z_2 \equiv q_1$
$z_3^+ \equiv q_0$

$z_3^+ \equiv q_0, q_0$
$z_2 \equiv q_2$

$z_3^+ \equiv q_0, q_0$
$z_2 \equiv q_1$





old State	Transition at 'a'	Transition at b.
$\pm z_1^- \equiv q_0$	$z_2 \equiv q_0$	$z_3 \equiv q_1$
$z_2 \equiv q_0$	$z_2 \equiv q_0$	$z_3 \equiv q_1$
$z_3 \equiv q_1$	$z_2 \equiv q_0$	$z_3 \equiv q_1$





old State	Transition at 'a'	Transition at 'b'
$\pm z_1^+ \equiv q_0$	$z_2^+ \equiv (q_0, q_0)$	$z_2^+ \equiv (q_0, q_0)$
$z_2^+ = q_0$	$z_2^+ \equiv (q_0, q_0)$	$z^+ z \equiv (q_0, q_0)$







NFA To DFA(FA).





To avoid Nondeterminism.




q0
q1
q2
q3
q4
b
b
b
a
a
a
a,b
a,b