





Common causes of death, normalized to COVID. Pre-Covid: 1Q2017 thru 1Q2020, Post-Covid 2Q2020 thru 1Q2021, USA





Relative effect of COVID on various age groups, and compared to cause of death. These are all relative to COVID, which is 1.0 on each of four charts, USA




Texas and Tri-county comparisons





<https://data.cdc.gov/NCHS/Conditions-contributing-to-deaths-involving-corona/hk9y-quqm/data>



	Under 65	Over 65
All	26.0%	74.0%
COVID-19	25.5%	74.5%

Conditions Contributing to COVID-19 Deaths, by State and Age, Provisional 2020-2022

This dataset shows health conditions and contributing causes mentioned in conjunction with deaths involving coronavirus disease 2019 (COVID-19) by age group and jurisdiction of occurrence. 2020-2022 data are provisional.

False Positives Demonstration

Use 0.19% as estimated daily incidence

Prevalence estimated as avg. infected period of 2 weeks X incidence

95% accuracy of test

0.19% X 14 = 2.660%

	Positive	Negative	
test pos	2.527%	4.867%	7.39%
test neg	0.133%	92.473%	92.61%
	2.660%	97.340%	100.00%

Sensitivity
Probability of detection where condition exists
 $\text{True} + / (\text{True} + \& \text{False} -)$
95%

False pos. is more than half of total positives.


TRUE +	2.527%/7.39%	34.2%
FALSE +	4.867%/7.39%	65.8%
Total		100.00%

Specificity
Probability of not detecting where condition doesn't exist
 $\text{True} - / (\text{True} - \& \text{False} +)$
95%



Example only; sensitivity and specificity not necessarily equal.

Average and Excess Deaths:

16-Jul-22



Average and Excess Deaths (cont'd):



USA Excess Deaths, 2020 (from CDC data):

Annualized on 52 weeks	All Cause	All Cause, excl. CV19	CV19
3 yr average before 2020	859:100,000	859:100,000	-
2020	1016:100,000	905:100,000	-
Diff.	157:100,000	46:100,000	111:100,000

29% of All-Cause excess deaths are non-CV19

USA Excess Deaths, 2021 (from CDC data):

Annualized on 52 weeks	All Cause	All Cause, excl. CV19	CV19
3 yr average before 2020	859:100,000	859:100,000	-
2021	1052:100,000	909:100,000	-
Diff.	193:100,000	50:100,000	143:100,000

26% of All-Cause excess deaths are non-CV19

USA Excess Deaths to date (2022, from CDC data):





Week 29	All Cause	All Cause, excl. CV19	CV19
3 yr average before 2020	487:100,000	487:100,000	-
2022	561:100,000	504:100,000	-
Diff.	74:100,000	17:100,000	57:100,000

Linear Year Projection

23% of All-Cause excess deaths not CV19 102:100,000

<https://data.cdc.gov/NCHS/Excess-Deaths-Associated-with-COVID-19/xkxf-xrst/data>




Total, latest update	424:100,000	113:100,000	311:100,000
Annualized	168:100,000	45:100,000	123:100,000



(CDC started updating this again 02 July 2022)

Data in recent weeks are incomplete. Only 60% of death records are submitted to NCHS within 10 days of the date of death, and completeness varies by jurisdiction. Data are not weighted and counts are likely underreported.

Recent exponential growth examples:



Vaccinations and cumulative outcomes:



<https://healthdata.gov/Health/COVID-19-Community-Profile-Report/gqxm-d9w9>

https://github.com/CSSEGISandData/COVID-19/blob/master/csse_covid_19_data/csse_covid_19_daily_reports_us/03-29-2022.csv

(Feb 2021 picked because that's about when vaccines became available)

