




Experimental page: ratios of curve fit deaths to curve fit confirmed cases (CFR)

















Excess deaths as a Z score:





Above based on Z score of two standard deviation from 2017-2019. What follows is cumulative plot of same.



Data in recent weeks are incomplete. Only 60% of death records are submitted to NCHS within 10 days of the date of death, and completeness varies by jurisdiction. Data are not weighted and counts are likely

False Positives Demonstration

Use 0.07% as estimated daily incidence

Prevalence estimated as avg. infected period of 2 weeks X incidence 00% accuracy of tost

Baseline set at Z = 0	excess deaths			Excess normalized to std dev of 2017 - 2019 deaths.		
250			uevoi	2017 - 2019 (ucauis.	
200	000000000000000000000000000000000000000	••••				
150						
N 100			0000000	000		
50 —	,	00000000				
0						
-50						
0 10	20	30 week	40	50	60	
	cumulative 2020	cumulative	2021			

99%	99% accuracy of test			0.07% X 14 = 0.980%
	Positive	Negative		
test pos	0.970%	0.990%	1.96%	
test neg	0.010%	98.030%	<u>98.04%</u>	
	0.980%	99.020%	100.00%	

False pos. is more than half of total positives.

Total		100.00%
FALSE +	0.99%/1.96%	<u>50.5%</u>
TRUE +	0.97%/1.96%	49.5%
•	•	

Counter-act this tendency by increasing test sensitivity. However this may increase false negatives, the recipients of which may be positive, think they're negative, and go spread it around some more.

US mortality vs. India 200:100,000 180:100,000 160:100,000 140:100,000 120:100,000 100:100,000 80:100,000 60:100,000 40:100,000 20:100,000 0:100,000 10-Mar 31-Oct 1-Dec 18-Jun 26-Sep 4-Jan 14-Apr 23-Jul US cum. fit per 100k India cum. fit per 100k

USA Excess Deaths, 2020 (from CDC data):

	Annualized on 52 weeks				
		All Cause	All Cause, excl. CV19	CV19	
3	yr average before 2020	859:100,000	859:100,000	-	
	2020	1016:100,000	905:100,000	-	
	Diff.	157:100,000	46:100,000	111:100,000	

3 yr average	
859:100,000	

29% of All-Cause excess deaths are non-CV19

https://data.cdc.gov/NCHS/Excess-Deaths-Associated-with-COVID-19/xkkf-xrst/data

USA Excess Deaths to date (from CDC data):

	32 weeks	All Cause	All Cause, excl. CV19	CV19
3	yr average before 2020	522:100,000	522:100,000	-
	2021	580:100,000	534:100,000	-
	Diff.	58:100,000	11:100,000	46:100,000

3 yr average 859:100,000

20% of All-Cause excess deaths are non-CV19


https://data.cdc.gov/NCHS/Excess-Deaths-Associated-with-COVID-19/xkkf-xrst/data


K = 0.318 R_o : R:

gamma = 0.171 $R_o = \exp(K/\text{gamma}) = 6.42$ 84%


gamma = 0.286 $R > 1 - 1/R_o = 3.04$ 67%

R is recovered variable.







CDC county data on week indicated new cases, by % fully vaccinated.



Estimates of Cases per 100k



reporting jurisdictions is not a uniform distribution; some data suppressed, for example Texas





https://data.cdc.gov/NCHS/Weekly-Counts-of-Deaths-by-State-and-Select-Causes/muzy-jte6/data