


Common causes of death, normalized to COVID. Pre-Covid: 1Q2017 thru 1Q2020, Post-Covid 2Q2020 thru 1Q2021, USA




Relative effect of COVID on various age groups, and compared to cause of death. These are all relative to COVID, which is 1.0 on each of four charts, USA













Texas and Tri-county comparisons









 $\underline{https://data.cdc.gov/NCHS/Conditions-contributing-to-deaths-involving-corona/hk9y-quqm/data}$





Average and Excess Deaths:











	Under 65	Over 65
All	26.0%	74.0%
COVID-19	25.5%	74.5%

Conditions Contributing to COVID-19 Deaths, by State and Age, Provisional 2020-2022

This dataset shows health conditions and contributing causes mentioned in conjunction with deaths involving coronavirus disease 2019 (COVID-19) by age group and jurisdiction of occurrence. 2020-2022 data are provisional.

False Positives Demonstration

Use 0.19% as estimated daily incidence

Prevalence estimated as avg. infected period of 2 weeks X incidence

95% accuracy of test				0.19% X 14 = 2.660%	
	Positive	Negative		Sensitivity	
test pos	2.527%	4.867%	7.39%	Probability of detection	
test neg	0.133%	92.473%	92.61%	where condition exists	
	2.660%	97.340%	100.00%	True + / (True + & False -)	
			95%		
False pos. is more than half of total positives.			Specificity		
TRUE + 2.527%/7.39%		34.2%	Probability of not detecting where		
FALSE + 4.867%/7.39%		65.8%	condition doesn't exist		
Total			100.00%	True - / (True - & False +)	
				000/	

Example only; sensitivity and specifity not necessarily equal.

USA Excess Deaths, 2020 (from CDC data):

Annualized on 52 weeks			
	All Cause	All Cause, excl. CV19	CV19
3 yr average before 2020	859:100,000	859:100,000	-
2020	1016:100,000	905:100,000	-
Diff.	157:100,000	46:100,000	111:100,000

 $29\%\,$ of All-Cause excess deaths are non-CV19

USA Excess Deaths, 2021 (from CDC data):

Annualized on 52 weeks			
	All Cause	All Cause, excl. CV19	CV19
3 yr average before 2020	859:100,000	859:100,000	-
2021	1052:100,000	909:100,000	-
Diff.	193:100,000	50:100,000	143:100,000

26% of All-Cause excess deaths are non-CV19

USA Excess Deaths to date (2022, from CDC data):

	Con Excess Beating to date (2022, from CBC data).				
	Week 34	All Cause	All Cause, excl. CV19	CV19	
3	yr average before 2020	564:100,000	564:100,000	-	
	2022	660:100,000	597:100,000	-	
	Diff.	96:100,000	33:100.000	62:100.000	


2022 Linear Year Projection


35% of All-Cause excess deaths not CV19


95:100,000


 $\underline{\text{https://data.cdc.gov/NCHS/Excess-Deaths-Associated-with-COVID-19/xkkf-xrst/data}}$

Annualized	170:100,000	49:100,000	121:100,000
Total, latest update	446:100,000	130:100,000	316:100,000












(CDC started updating this again 02 July 2022)

Data in recent weeks are incomplete. Only 60% of death records are submitted to NCHS within 10 days of the date of death, and completeness varies by jurisdiction. Data are not weighted and counts are likely underreported.

Recent exponential growth examples:







Vaccinations and cumulative outcomes:



(Feb 2021 picked because that's about when vaccines became available)



https://healthdata.gov/Health/COVID-19-Community-Profile-Report/gaxm-d9w9
https://github.com/CSSEGISandData/COVID-19/blob/master/csse covid 19 daily reports us/03-29-2022.csv