

Noether $\rightarrow \alpha$ v0.3: Fixing L_ψ , Z and g_5 ab-initio

Draft for UBT Project

August 30, 2025

1 Setup: 5D Action, Geometry, BC

We consider $M^4 \times S_\psi^1$ with coordinates x^μ and $\psi \sim \psi + L_\psi$. Let $\Theta(x, \psi)$ denote the unified (biquaternionic) field and $A_M(x, \psi)$ a $U(1)$ gauge field. The 5D action (natural units $c = \hbar = 1$) is

$$S = \int d^4x \int_0^{L_\psi} d\psi \sqrt{|g_5|} \left[g^{MN} (D_M \Theta)^\dagger (D_N \Theta) - V(\Theta) - \frac{1}{4} g^{MR} g^{NS} F_{MN} F_{RS} \right], \quad (1)$$

with $D_M = \partial_M + ig_5 A_M$, $F_{MN} = \partial_M A_N - \partial_N A_M$. We allow a warped background

$$ds^2 = e^{2A(\psi)} \eta_{\mu\nu} dx^\mu dx^\nu + e^{2B(\psi)} d\psi^2, \quad \sqrt{|g_5|} = e^{4A(\psi) + B(\psi)}. \quad (2)$$

Boundary conditions (BC) along S_ψ^1 may be periodic or twisted (phase δ).

Holonomy (Wilson line). The gauge-invariant holonomy (Hosotani parameter) is

$$\theta_H \equiv g_5 \oint_{S_\psi^1} A_\psi d\psi = g_5 \int_0^{L_\psi} d\psi A_\psi(x, \psi). \quad (3)$$

Large gauge transformations shift $\theta_H \rightarrow \theta_H + 2\pi n$, $n \in \mathbb{Z}$.

2 Reduction and Canonical Normalization

Assume the photon zero-mode $A_\mu^{(0)}(x)$ is independent of ψ and normalized canonically in 4D. Reducing the gauge kinetic term in (1) gives

$$S_{\text{gauge}} \supset -\frac{1}{4} \int d^4x Z F_{\mu\nu}^{(0)} F_{(0)}^{\mu\nu}, \quad Z \equiv \int_0^{L_\psi} d\psi e^{B(\psi)-2A(\psi)}, \quad (4)$$

so the canonically normalized 4D photon is $A_\mu^{(0)} \rightarrow A_\mu^{(0)}/\sqrt{Z}$. The covariant derivative contributes the interaction

$$\int d^4x \int_0^{L_\psi} d\psi \sqrt{|g_5|} J^\mu A_\mu \longrightarrow \int d^4x g_4 J_{(0)}^\mu A_\mu^{(0)}, \quad g_4 = \frac{g_5}{\sqrt{Z}}, \quad (5)$$

where J^μ is the Noether current density and $J_{(0)}^\mu$ its overlap with the photon zero-mode. Therefore the fine-structure constant at a reference scale μ_0 is

$$\boxed{\alpha(\mu_0) = \frac{g_4^2}{4\pi} = \frac{g_5^2}{4\pi Z}}. \quad (6)$$

3 Noether Charge and Current Matching

Global $U(1)$: $\Theta \rightarrow e^{i\lambda}\Theta$ yields the 5D Noether current

$$J^M = i [\Theta^\dagger (D^M \Theta) - (D^M \Theta)^\dagger \Theta] \Big|_{A=0}. \quad (7)$$

We fix the normalization of Θ such that the fundamental charged excitation has unit Noether charge

$$Q \equiv \int d^3x \int_0^{L_\psi} d\psi \sqrt{|g_5|} J^0 = \pm 1. \quad (8)$$

This fixes the overall scale entering the coupling (5); no additional free normalization survives.

4 Wilson Line and Quantization

On the compact ψ -cycle the holonomy (3) is physical. For a field with $U(1)$ charge q and BC phase δ the KK momenta are shifted by

$$p_\psi^{(n)} = \frac{2\pi}{L_\psi} (n + a), \quad a \equiv \frac{q\theta_H}{2\pi} + \delta, \quad n \in \mathbb{Z}. \quad (9)$$

Stationary vacua (Hosotani mechanism) are determined dynamically and may select nontrivial θ_H^* . Large gauge invariance enforces periodicity $\theta_H \sim \theta_H + 2\pi$; stable vacua satisfy $\partial V_{\text{eff}}/\partial\theta_H = 0$.

5 One-Loop Effective Potential $V_{\text{eff}}(L_\psi, \theta_H)$

For each field j with spin-statistics sign $\sigma_j = \pm 1$ (boson $+1$, fermion -1), degeneracy d_j , mass m_j , charge q_j , and twist δ_j , the one-loop contribution is

$$V_j(L_\psi, \theta_H) = \frac{\sigma_j d_j}{2} \int \frac{d^4 p}{(2\pi)^4} \sum_{n \in \mathbb{Z}} \ln \left[p^2 + m_j^2 + \left(\frac{2\pi}{L_\psi} \right)^2 (n + a_j)^2 \right], \quad (10)$$

$$a_j \equiv \frac{q_j \theta_H}{2\pi} + \delta_j. \quad (11)$$

Using standard contour/Matsubara techniques (equivalently Poisson resummation) one obtains an exact resummed form

$$V_j(L_\psi, \theta_H) = \sigma_j d_j \int \frac{d^4 p}{(2\pi)^4} \frac{1}{L_\psi} \ln \left(1 - 2e^{-L_\psi \omega_j(p)} \cos 2\pi a_j + e^{-2L_\psi \omega_j(p)} \right), \quad (12)$$

where $\omega_j(p) = \sqrt{p^2 + m_j^2}$. The total potential is

$$V_{\text{eff}}(L_\psi, \theta_H) = \sum_j V_j(L_\psi, \theta_H). \quad (13)$$

The vacuum (L_ψ^*, θ_H^*) solves

$$\frac{\partial V_{\text{eff}}}{\partial \theta_H} = 0, \quad \frac{\partial V_{\text{eff}}}{\partial L_\psi} = 0. \quad (14)$$

These conditions fix L_ψ and θ_H *dynamically* in terms of the field content and BC, with no fits.

Massless limit and modular dependence. For $m_j L_\psi \ll 1$ one can expand (12) to obtain polylogarithms $\text{Li}_5(e^{\pm i 2\pi a_j})$, reproducing the familiar $1/L_\psi^4$ scaling and periodic dependence on a_j . Massive fields yield exponentially suppressed Bessel-function tails for large $m_j L_\psi$.

6 Determining Z (Warped Case)

If the vacuum back-reacts and generates a nontrivial warp, the 4D gauge kinetic factor is

$$Z = \int_0^{L_\psi} d\psi e^{B(\psi)-2A(\psi)} |\xi_0(\psi)|^2, \quad (15)$$

where $\xi_0(\psi)$ is the photon zero-mode profile (constant in flat space). The warp factors $A(\psi)$, $B(\psi)$ and $\xi_0(\psi)$ follow from the coupled background equations of motion derived from (1) (e.g. via a first-order BPS system if available). In flat space, $\xi_0(\psi) \equiv 1$ and $Z = L_\psi$. In general we write

$$Z = L_\psi f(\tau, \text{BC}), \quad (16)$$

where f encodes the modular parameter τ (complex-time sector) and boundary data.

7 Putting It Together: α from UBT

Combining (6), (14) and (15),

$$\alpha(\mu_0) = \frac{g_5^2}{4\pi Z^\star} = \frac{g_5^2}{4\pi L_\psi^\star f^\star(\tau, \text{BC})}, \quad (L_\psi^\star, \theta_H^\star) \text{ solve } \partial V_{\text{eff}} = 0. \quad (17)$$

Here g_5 is fixed by Noether normalization (8) (unit charge for the fundamental excitation) and current matching (5). Thus, given the field content, BC, and UBT potential $V(\Theta)$, the pair $(L_\psi^\star, \theta_H^\star)$ and the warp A, B are determined and *no free fit survives* in α . Standard QED running then connects $\alpha(\mu_0)$ to experimental scales.

8 Electron Mass Consistency (Sketch)

The same background must produce the lightest charged KK eigenvalue

$$m_e^2 = \lambda_{\min} \left[-e^{-B} \partial_\psi (e^{B-2A} \partial_\psi \cdot) + m_\Theta^2 e^{-2A} + \dots \right], \quad (18)$$

for the appropriate (spinor) sector and BC, including the holonomy shift (9). This provides a nontrivial cross-check: the *same* $(L_\psi^\star, \theta_H^\star, A, B)$ that gives α must also yield m_e .

9 Checklist to Avoid Numerology

- Use (12) (or its Poisson/Bessel form) *exactly*; minimize (14) without inserting data.
- Fix Noether charge to $Q = \pm 1$ via (8); this removes arbitrary rescalings of Θ .
- Compute Z from (4) or (15); do not set $Z = L_\psi$ unless flat space is justified by the background EOM.
- Only after obtaining $\alpha(\mu_0)$ ab-initio apply perturbative running to compare to $\alpha(M_Z)$.