

A Unified Field Theory on the Biquaternionic Manifold \mathbb{B}^4 : Consciousness, Quantum Fields, and Emergent Space-Time

David Jaroš

2025

Abstract

We develop a mathematically consistent unified field theory constructed on a four-dimensional biquaternionic manifold \mathbb{B}^4 , which incorporates complexified spacetime and internal phase dimensions naturally into its geometric structure. The core field $\Theta(q)$ is a tensor-spinor-gauge-valued section over \mathbb{B}^4 , encoding all known fundamental interactions, emergent quantum behavior, and phenomenology of consciousness. The theory eliminates the need for external projections ...

We develop a rigorous unified field theory on a four-dimensional biquaternionic manifold $\mathbb{B}^4 = (\mathbb{C} \otimes \mathbb{H})^4$, where each coordinate is a biquaternion representing complexified internal and external degrees of freedom. The primary field $\Theta(q)$ is a tensor-spinor-valued section over \mathbb{B}^4 , encoding gravitational, gauge, quantum, and cognitive structure. We formulate the covariant dynamics, metric geometry, low-energy quantum limits, emergent conscious par...

1 Biquaternionic Manifold and Field Definition

We define the manifold $\mathcal{M} = \mathbb{B}^4$, where each point is a 4-tuple of biquaternions:

$$q^\mu = x^\mu + iy^\mu + \mathbf{j}z^\mu + i\mathbf{j}w^\mu, \quad \mu = 0, 1, 2, 3,$$

with components $x^\mu, y^\mu, z^\mu, w^\mu \in \mathbb{R}$. The basis elements $\{1, \mathbf{i}, \mathbf{j}, \mathbf{k}\}$ satisfy the quaternionic algebra.

The unified field is defined as:

$$\Theta(q) \in \Gamma \left(T^{(1,1)}(\mathbb{B}^4) \otimes \mathbb{S} \otimes \mathbb{G} \right),$$

where $T^{(1,1)}$ denotes the (1,1) tensor bundle over \mathbb{B}^4 , \mathbb{S} is a spinor bundle, and \mathbb{G} is an internal gauge fiber.

2 Metric Geometry and Covariant Derivatives

We define a complexified metric tensor on \mathbb{B}^4 :

$$G_{\mu\nu}(q) = \langle dq^\mu, dq^\nu \rangle,$$

where $\langle \cdot, \cdot \rangle$ is a biquaternionic inner product. The affine and spin connections $\Gamma_{\mu\nu}^\rho$, Ω_μ are derived accordingly.

Covariant derivative:

$$\mathcal{D}_\mu \Theta = \partial_\mu \Theta + \Omega_\mu \cdot \Theta + ig A_\mu^a T^a \Theta,$$

with gauge coupling and spin connection terms included.

3 General Relativity and Emergent Geometry

The metric tensor $G_{\mu\nu}(q)$ defined over \mathbb{B}^4 generalizes the Lorentzian metric of General Relativity to a biquaternionic setting. The curvature tensors are constructed from the complexified affine connection:

$$R_{\sigma\mu\nu}^\rho = \partial_\mu \Gamma_{\nu\sigma}^\rho - \partial_\nu \Gamma_{\mu\sigma}^\rho + \Gamma_{\mu\lambda}^\rho \Gamma_{\nu\sigma}^\lambda - \Gamma_{\nu\lambda}^\rho \Gamma_{\mu\sigma}^\lambda.$$

The Einstein field equations emerge as a projection of the variation of the geometric Lagrangian:

$$\delta \mathcal{L}_{\text{geom}} = \frac{1}{2\kappa} \left(R_{\mu\nu} - \frac{1}{2} G_{\mu\nu} R \right) \delta G^{\mu\nu}.$$

This shows that classical spacetime curvature is a low-energy limit of the intrinsic geometry of the biquaternionic manifold. The extended degrees of freedom in y^μ, z^μ give rise to higher-order corrections and dynamical compactification in early-universe cosmology.

4 Lagrangian and Field Equations

Total action on \mathbb{B}^4 reads:

$$S = \int_{\mathbb{B}^4} d^4q \sqrt{|\det G|} (\mathcal{L}_\Theta + \mathcal{L}_{\text{geom}} + \mathcal{L}_{\text{gauge}}),$$

where:

$$\mathcal{L}_\Theta = \Re [\bar{\Theta} (i\Gamma^\mu \mathcal{D}_\mu - M(q)) \Theta],$$

with $M(q)$ potentially depending on internal phase coordinates y^μ, z^μ , representing cognitive or entropic symmetry breaking.

5 Tensor Decomposition of Θ

The field can be locally decomposed as:

$$\Theta(q) = \phi(q) + i\psi^\mu(q)\gamma_\mu + \eta(q)\mathbb{I} + i\chi(q)\mathbb{J},$$

where $\chi(q)$ represents internal oscillations related to consciousness and subjective time.

6 Quantum Limits and Classical Reductions

In the real spacetime limit $y^\mu, z^\mu, w^\mu \rightarrow 0$, the action reduces to:

- Dirac equation for spinor sector:

$$(i\gamma^\mu \partial_\mu - m)\psi = 0,$$

- Schrödinger equation in nonrelativistic limit:

$$i\hbar \partial_t \psi = \left(-\frac{\hbar^2}{2m} \nabla^2 + V \right) \psi,$$

- Effective quantum field theory in flat $\mathbb{R}^4 \subset \mathbb{B}^4$.

7 Gauge Theory and Standard Model Embedding

The internal gauge structure of Θ accommodates:

$$\mathbb{G} \cong SU(3) \times SU(2) \times U(1),$$

and the covariant derivative generalizes to the Yang-Mills form. Higgs-like mass terms can arise via internal symmetry oscillations in the z^μ, w^μ directions.

8 Gauge Symmetries: QED, QCD and the Standard Model Embedding

We extend the field $\Theta(q)$ to carry internal gauge indices under the symmetry group $\mathcal{G} = U(1) \times SU(2) \times SU(3)$, corresponding to the Standard Model gauge groups for electromagnetism, weak, and strong interactions.

Let $A_\mu = A_\mu^a T^a$ denote the full gauge connection, with generators T^a acting on $\Theta(q)$ in the appropriate internal representation.

The full covariant derivative becomes:

$$\mathcal{D}_\mu \Theta = \partial_\mu \Theta + \Omega_\mu \cdot \Theta + ig_1 B_\mu Y \Theta + ig_2 W_\mu^i \tau^i \Theta + ig_3 G_\mu^a \lambda^a \Theta,$$

where: - B_μ is the $U(1)_Y$ hypercharge field, - W_μ^i are the weak $SU(2)_L$ fields, - G_μ^a are the gluon fields of $SU(3)_C$, - Y , τ^i , and λ^a are the corresponding generators.

The gauge-invariant kinetic Lagrangian reads:

$$\mathcal{L}_{\text{gauge}} = -\frac{1}{4} \sum_a F_{\mu\nu}^a F^{a\mu\nu},$$

with:

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + f^{abc} A_\mu^b A_\nu^c,$$

where f^{abc} are the structure constants of the respective Lie algebras.

The interaction term in the unified Lagrangian:

$$\mathcal{L}_{\Theta,\text{int}} = \Re [\bar{\Theta} i \Gamma^\mu \mathcal{D}_\mu \Theta],$$

ensures that the dynamics of Θ is fully coupled to the standard model gauge fields.

:

9 Low-Energy Limits: Dirac and Schrödinger Equations

To demonstrate compatibility with established quantum mechanics, we derive the Dirac and Schrödinger equations as limiting cases of the unified biquaternionic field equation for $\Theta(q)$.

Starting from the unified covariant derivative:

$$\mathcal{D}_A \Theta = \partial_A \Theta + \Omega_A \cdot \Theta + ig A_A^a T^a \Theta,$$

we consider the case where curvature effects are small, gauge fields are static or slowly varying, and the manifold \mathbb{C}^5 can be approximated by a local inertial frame. In this approximation, the equation of motion simplifies to:

$$\Gamma^A \partial_A \Theta = m \Theta,$$

which reduces to the ****Dirac equation**** in the limit where only 4 spacetime dimensions are active and Γ^A matrices correspond to standard Dirac matrices γ^μ . This confirms that the spinor nature of Θ is consistent with relativistic quantum field theory.

In the non-relativistic limit ($v \ll c$), a standard Foldy–Wouthuysen decomposition leads to the Schrödinger equation:

$$i\hbar \frac{\partial \psi}{\partial t} = \left(-\frac{\hbar^2}{2m} \nabla^2 + V \right) \psi,$$

where ψ is a projection of Θ onto low-energy modes.

Thus, our unified theory contains both quantum mechanics and relativistic field theory as natural limits, while offering a richer structure due to its biquaternionic and spin-tensor form defined on \mathbb{C}^5 .

10 Conscious Oscillations and Emergent Mind

Internal modes $\chi(q) \sim \sin(\omega \cdot y^\mu)$ describe periodic subjective phase. Collapse of these oscillations projects classical reality. We interpret eigenmodes:

$$\Theta_n(q) \sim e^{in\psi} \Psi_n(x),$$

as discrete conscious quanta (psychons), with transitions corresponding to awareness or memory shifts.

11 Gauge Symmetries: QED, QCD and the Standard Model Embedding

We extend the field $\Theta(q)$ to carry internal gauge indices under the symmetry group $\mathcal{G} = U(1) \times SU(2) \times SU(3)$, corresponding to the Standard Model gauge groups for electromagnetism, weak, and strong interactions.

Let $A_\mu = A_\mu^a T^a$ denote the full gauge connection, with generators T^a acting on $\Theta(q)$ in the appropriate internal representation.

The full covariant derivative becomes:

$$\mathcal{D}_\mu \Theta = \partial_\mu \Theta + \Omega_\mu \cdot \Theta + ig_1 B_\mu Y \Theta + ig_2 W_\mu^i \tau^i \Theta + ig_3 G_\mu^a \lambda^a \Theta,$$

where:

- B_μ is the $U(1)_Y$ hypercharge field,
- W_μ^i are the weak $SU(2)_L$ fields,
- G_μ^a are the gluon fields of $SU(3)_C$,
- Y , τ^i , and λ^a are the corresponding generators.

The gauge-invariant kinetic Lagrangian reads:

$$\mathcal{L}_{\text{gauge}} = -\frac{1}{4} \sum_a F_{\mu\nu}^a F^{a\mu\nu},$$

with:

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + f^{abc} A_\mu^b A_\nu^c,$$

where f^{abc} are the structure constants of the respective Lie algebras.

The interaction term in the unified Lagrangian:

$$\mathcal{L}_{\Theta, \text{int}} = \Re [\bar{\Theta} i \Gamma^\mu \mathcal{D}_\mu \Theta],$$

ensures that the dynamics of Θ is fully coupled to the standard model gauge fields.

12 Free Energy Principle and Fokker–Planck Flow

We apply the FEP:

$$\frac{\partial P}{\partial \psi} = -\nabla_q \cdot (DP) + \frac{1}{2} \nabla_q^2 (D^2 P),$$

modeling informational entropy flow across \mathbb{B}^4 . Drift terms represent prediction error; diffusion encodes uncertainty and updating.

13 Cosmological Aspects

The internal phase coordinate plays a role similar to an inflaton or entropy-gradient field. Toroidal compactification of internal directions leads to inflationary and cyclic cosmological models.

14 Conclusion

We have proposed a unified framework on \mathbb{B}^4 where geometry, gauge theory, quantum physics, and subjective consciousness are aspects of a single field $\Theta(q)$. This approach explains both standard physics and phenomenological features of cognition, predicting new psychon-like excitations and quantum collapse via internal oscillations.

15 Quantum Electrodynamics and Quantum Chromodynamics

The field $\Theta(q)$ contains internal symmetry structures corresponding to the gauge groups of the Standard Model. In particular, the internal gauge fiber \mathbb{G} carries representations of:

$$\mathbb{G} \cong SU(3)_{\text{color}} \times SU(2)_L \times U(1)_Y,$$

embedded within the matrix structure of Θ .

15.1 QED Sector

The Abelian part of the gauge group $U(1)$ governs the electromagnetic interaction. The corresponding gauge field A_μ enters the covariant derivative as:

$$\mathcal{D}_\mu \Theta = \partial_\mu \Theta + \Omega_\mu \cdot \Theta + ieA_\mu \Theta,$$

with Maxwell-type field strength:

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu.$$

15.2 QCD Sector

The non-Abelian color interaction is embedded via the gluon fields G_μ^a , $a = 1, \dots, 8$, and SU(3) generators T^a :

$$\mathcal{D}_\mu \Theta = \dots + ig_s G_\mu^a T^a \Theta.$$

The gluonic field strength is:

$$G_{\mu\nu}^a = \partial_\mu G_\nu^a - \partial_\nu G_\mu^a + g_s f^{abc} G_\mu^b G_\nu^c.$$

15.3 Symmetry Breaking and Higgs-Like Mechanism

The internal phase coordinates y^μ, z^μ provide a natural mechanism for spontaneous symmetry breaking:

$$\langle \chi(q) \rangle \neq 0 \quad \Rightarrow \quad m \neq 0,$$

where χ acts as a Higgs-like internal mode within Θ . Mass hierarchies emerge from oscillatory patterns in the biquaternionic manifold.

16 Discussion of Particle Spectrum

Excitations of the field Θ include not only standard fermions and bosons but also additional modes due to its internal spinor-gauge structure. We predict:

- Standard particle spectrum (quarks, leptons, gauge bosons) as tensor-spinor projections,
- Scalar internal oscillations as Higgs-like or axion-like particles,
- Psychon modes — coherent phase oscillations corresponding to discrete conscious states,
- Potential graviton-like curvature modes in the geometric component.

17 Summary and Outlook

This framework combines geometry, quantum field theory, and information dynamics in a single elegant model defined on \mathbb{B}^4 . It naturally unifies:

- General relativity as geometric dynamics of $G_{\mu\nu}(q)$,
- Quantum field theory via tensor-spinor structure of $\Theta(q)$,
- Gauge interactions through internal fiber \mathbb{G} ,
- Consciousness as internal oscillations and collapse of $\chi(q)$.

Future work will explore quantization in curved biquaternionic space, phenomenology of psychon transitions, and links to holographic principles and modular topologies.

18 Limit: General Relativity

In the classical limit where oscillatory and imaginary components vanish, the biquaternionic metric $G_{\mu\nu}(q)$ reduces to a standard pseudo-Riemannian real-valued tensor on a 4D Lorentzian manifold.

