

Covariant Biquaternionic Tetrads in UBT: Metric Rank, Connection Reconstruction, and Integrability

Ing. David Jaroš

19 July 2026

Abstract

We formulate the local classical geometry of Unified Biquaternion Theory (UBT) directly from the covariant first derivative of its single fundamental biquaternionic field. Defining $E_\mu = \mathcal{N}_0^{-1/2} D_\mu \Theta$, the classical Lorentz sector is the real four-dimensional slice $E_\mu = ie_\mu^0 \mathbf{1} + e_\mu^k \mathbf{e}_k$. Quaternion conjugation gives the exact central identity

$$\frac{1}{2}(E_\mu^\# E_\nu + E_\nu^\# E_\mu) = g_{\mu\nu} \mathbf{1},$$

so the metric is not defined by a trace, real-part map, phase projection, or imaginary-time fiber average. The tetrad-to-metric map has rank ten at every nondegenerate tetrad and a six-dimensional kernel equal to local Lorentz frame freedom. We then reconstruct the connection. For every tetrad and specified torsion, the metric-compatible Lorentz connection is uniquely $\omega = \hat{\omega}(e) + K(T)$; the torsion-free GR branch is the special case $K = 0$. Hence the remaining full-UBT question is the dynamics of torsion, not a kinematically arbitrary Ω_μ . We give an explicit affine field Θ_{SR} generating the Minkowski tetrad and solving the standard flat source-free second-order master operator. Finally, we prove that a naive one-sided connection with invertible Θ has a flatness obstruction under the stated torsion-free compatibility assumptions, while a two-sided biquaternionic derivative obeys $[D_\mu, D_\nu]\Theta = F_{\mu\nu}^A \Theta - \Theta F_{\mu\nu}^B$ and permits nonzero intertwined left/right curvatures. Lorentz-slice and metric compatibility reduce its pure no-new-field representative to $A_\mu = \Omega_\mu$, $B_\mu = -\Omega_\mu^\dagger$, modulo a cancelling central term. We then prove that this pure pairing makes every nondegenerate torsion-free tetrad generated by one Θ admit a concurrent vector $\nabla_\mu V^\nu = \delta_\mu^\nu$; it therefore excludes Schwarzschild with nonzero mass and generic Einstein geometries. The remaining curved route must derive a nontrivial relative bimodule action as a composite or auxiliary nonpropagating variable. Curved-space existence, torsion dynamics, and Einstein dynamics from the canonical UBT action remain open.

Contents

1	Scope and proof status	2
2	Relation to standard tetrad geometry and earlier biquaternionic work	3
3	Biquaternionic algebra and the Lorentz slice	4

4	The covariant UBT tetrad	4
5	Projection-Free Metric and Algebraic Bivector	5
6	The local rank theorem	5
7	Connection and Christoffel symbols	6
8	Curvature and special relativity	7
9	Integrability and representation selection	7
9.1	One-sided flatness obstruction	7
9.2	Two-sided biquaternionic derivative	8
9.3	Pure Lorentz-compatible pair and no-new-field reduction	8
9.4	Exact prescribed-connection integrability criterion	9
10	Torsion dynamics in the minimal first-order branch	9
11	Lorentz-slice and imaginary-time stability	10
12	Conditional infrared endpoint of the metric dynamics	10
13	Complex time and regularity	11
14	Relation to earlier compact-fiber closure	11
15	Schwarzschild and perturbations: honest current status	11
16	Revised gap ledger	11
17	Conclusion	13

1 Scope and proof status

The purpose of this paper is to establish a minimal local geometric structure consistent with the original biquaternionic intuition of UBT and to identify exactly what remains to be proved. We do not claim an unconditional derivation of the Einstein equations.

Proved in this draft

- a projection-free Lorentzian metric from a central biquaternionic anticommutator;
- local representation of every Lorentz metric and rank ten of the tetrad-to-metric differential, with six Lorentz gauge directions;
- unique reconstruction of every metric-compatible frame connection from the tetrad and specified torsion;
- the torsion-free Levi–Civita spin connection as the unique classical GR connection;
- an explicit affine Θ representer of every constant Lorentz tetrad, including Minkowski spacetime;
- a no-go theorem for the naive one-sided invertible curved branch and the exact curvature identity for a two-sided biquaternionic derivative;
- exact reduction of the pure Lorentz-compatible pair to one spin connection, with no independent fields A_μ, B_μ ;
- a concurrent-vector no-go theorem showing that this pure pair cannot generate generic curved GR or Schwarzschild with nonzero mass.

Open

- deriving vacuum torsion or spin-sourced torsion from the canonical UBT action;
- deriving from the canonical action a nontrivial relative left/right action beyond the pure spin lift, and proving that it is composite or auxiliary rather than propagating;
- proving local and global existence for the implicit curved-space system $D_\mu \Theta = \sqrt{\mathcal{N}_0} E_\mu$;
- proving that the complete dynamics preserves the real Lorentz slice;
- deriving Einstein dynamics and the perturbation bridge from the original UBT master dynamics;
- canonical on-shell generation of Schwarzschild, Kerr, FRW, and wave sectors.

2 Relation to standard tetrad geometry and earlier biquaternionic work

The tetrad postulate, Levi–Civita spin connection, torsion, and contorsion used below are standard Cartan/tetrad geometry [1, 2]; they are not claimed as new theorems of UBT. The UBT-specific content is the identification $E_\mu = \mathcal{N}_0^{-1/2} D_\mu \Theta$, the central biquaternionic

anticommutator metric on the Lorentz slice, its rank interpretation, the affine single-field Minkowski representer, the one-sided obstruction/two-sided intertwiner analysis, and the exact pure-pair reduction with its concurrent-vector obstruction. Related complex and biquaternionic gravity programmes include Refs. [4, 5]; the present paper makes no claim that standard frame geometry originates here.

3 Biquaternionic algebra and the Lorentz slice

Let $\mathbb{B} = \mathbb{C} \otimes \mathbb{H}$, with commuting complex unit i and quaternion units

$$\mathbf{e}_j \mathbf{e}_k = -\delta_{jk} \mathbf{1} + \epsilon_{jkl} \mathbf{e}_l.$$

Quaternion conjugation $\sharp$ reverses the quaternion units without conjugating the commuting complex coefficients:

$$(z^0 \mathbf{1} + z^k \mathbf{e}_k)^\sharp = z^0 \mathbf{1} - z^k \mathbf{e}_k.$$

Definition 3.1 (Classical Lorentz module).

$$W_L = \{ix^0 \mathbf{1} + x^k \mathbf{e}_k \mid x^a \in \mathbb{R}\}.$$

For $X = ix^0 \mathbf{1} + x^k \mathbf{e}_k$ and $Y = iy^0 \mathbf{1} + y^k \mathbf{e}_k$, let

$$\eta(X, Y) = -x^0 y^0 + x^k y^k.$$

Theorem 3.2 (Central anticommutator identity). *For all $X, Y \in W_L$,*

$$\boxed{\frac{1}{2}(X^\sharp Y + Y^\sharp X) = \eta(X, Y) \mathbf{1}.} \quad (1)$$

Proof. In scalar–vector notation, quaternion multiplication is $(a, \mathbf{u})(b, \mathbf{v}) = (ab - \mathbf{u} \cdot \mathbf{v}, a\mathbf{v} + b\mathbf{u} + \mathbf{u} \times \mathbf{v})$. For $X = (ix^0, \mathbf{x})$, $X^\sharp = (ix^0, -\mathbf{x})$. The scalar part of $X^\sharp Y$ is $-x^0 y^0 + \mathbf{x} \cdot \mathbf{y}$. The vector parts of $X^\sharp Y$ and $Y^\sharp X$ are opposites, including the cross products, so they cancel in the half-sum. $\square$

4 The covariant UBT tetrad

The fundamental field remains

$$\Theta(q, \tau) \in \mathbb{B}, \quad \tau = t + i\psi.$$

On physical spacetime it is assumed smooth. Define

$$\boxed{E_\mu := \mathcal{N}_0^{-1/2} D_\mu \Theta,} \quad (2)$$

where $\mathcal{N}_0 > 0$ is fixed globally and D_μ is a covariant derivative. The connection is kept representation-independent at this stage:

$$D_\mu \Theta = \partial_\mu \Theta + \rho_*(\Omega_\mu) \Theta. \quad (3)$$

The map ρ_* must be specified in each concrete representation; Eq. (3) does not assume left, right, or two-sided action prematurely.

Definition 4.1 (Admissible classical tetrad). *A configuration belongs to the classical tetrad sector if*

$$E_\mu = ie_\mu^0 \mathbf{1} + e_\mu^k \mathbf{e}_k \in W_L$$

for all μ and the matrix e_μ^a is invertible.

5 Projection-Free Metric and Algebraic Bivector

By Theorem 1,

$$\boxed{\frac{1}{2}(E_\mu^\sharp E_\nu + E_\nu^\sharp E_\mu) = g_{\mu\nu} \mathbf{1}}, \quad (4)$$

where

$$g_{\mu\nu} = e_\mu^a e_\nu^b \eta_{ab}, \quad \eta_{ab} = \text{diag}(-1, 1, 1, 1). \quad (5)$$

The complete left-hand side is already a central real element. Hence Eq. (4) defines the metric without discarding or projecting noncentral components. The fixed constant $\mathcal{N}_0$ only sets units.

The complementary antisymmetric product is

$$\Sigma_{\mu\nu} := \frac{1}{2}(E_\mu^\sharp E_\nu - E_\nu^\sharp E_\mu), \quad \Sigma_{\nu\mu} = -\Sigma_{\mu\nu}. \quad (6)$$

Thus one product contains both the symmetric central geometry and an oriented bivector sector. No gauge-field identification of $\Sigma_{\mu\nu}$ is made without a separate transformation and dynamical analysis.

6 The local rank theorem

Define $\Phi(e) = e\eta e^\top$.

Theorem 6.1 (Full local metric rank). *At every invertible tetrad, the differential $d\Phi_e : \mathbb{R}^{4 \times 4} \rightarrow \text{Sym}_4(\mathbb{R})$ is surjective and has rank ten. Its kernel is the six-dimensional local Lorentz algebra.*

Proof. The first variation is

$$\delta g_{\mu\nu} = \eta_{ab}(\delta e_\mu^a e_\nu^b + e_\mu^a \delta e_\nu^b).$$

For an arbitrary symmetric $h_{\mu\nu}$ choose

$$\delta e_\mu^a = \frac{1}{2} h_{\mu\rho} e^{\rho a}.$$

Then $\delta g_{\mu\nu} = h_{\mu\nu}$. Hence the differential is surjective and has rank ten. The domain has dimension sixteen, leaving a six-dimensional kernel. The transformations $\delta e = e\lambda$ with $\lambda^\top \eta + \eta \lambda = 0$ lie in the kernel and constitute the six-dimensional Lorentz Lie algebra. $\square$

Corollary 6.2. *The local comparison “a biquaternion has eight real components, whereas a metric has ten” is not the relevant rank count. The metric depends on four covariant first derivatives, which form a sixteen-component real tetrad before local Lorentz redundancy is removed.*

This theorem closes the local algebraic rank question. It does not prove that all tetrad variations arise from admissible variations of one on-shell Θ and its derived connection.

7 Connection and Christoffel symbols

The Christoffel symbols and the frame connection describe different indices:

- $\Gamma_{\mu\nu}^\rho$ transports coordinate indices;
- $\omega_\mu^a{}_b$, and its spin/biquaternionic lift Ω_μ , transport the local Lorentz frame.

They are related by

$$\partial_\mu e_\nu^a - \Gamma_{\mu\nu}^\rho e_\rho^a + \omega_\mu^a{}_b e_\nu^b = 0. \quad (7)$$

It is generally incorrect to identify Γ with a real component of Ω .

Theorem 7.1 (Unique torsion-free classical frame connection). *For every nondegenerate C^2 tetrad, metric compatibility and vanishing torsion determine a unique Lorentz connection,*

$$\boxed{\dot{\omega}_\mu^a{}_b = e_b{}^\nu \left(\dot{\Gamma}_{\mu\nu}^\rho(g) e_\rho^a - \partial_\mu e_\nu^a \right)}, \quad (8)$$

with $\dot{\omega}_{\mu ab} = -\dot{\omega}_{\mu ba}$. Its spin lift is unique in the chosen representation, up to ordinary local Lorentz gauge.

Proof. The Levi-Civita connection $\dot{\Gamma}(g)$ is uniquely fixed by $\dot{\nabla}g = 0$ and $\dot{\Gamma}_{[\mu\nu]}^\rho = 0$. Solving Eq. (7) gives Eq. (8). If two Lorentz connections have the same zero torsion, their difference is antisymmetric in its first two frame indices and symmetric in the indices forced by equality of torsion; cycling these symmetries gives zero. $\square$

The torsion-free result extends to arbitrary specified torsion. Write

$$T^a = de^a + \omega^a{}_b \wedge e^b = \frac{1}{2} T^a{}_{bc} e^b \wedge e^c. \quad (9)$$

Theorem 7.2 (Tetrad-torsion reconstruction). *For every nondegenerate tetrad and specified torsion, there is exactly one metric-compatible Lorentz connection,*

$$\boxed{\omega^a{}_b = \dot{\omega}^a{}_b(e) + K^a{}_b(T)}, \quad (10)$$

where, with the convention above,

$$\boxed{K_{abc} = \frac{1}{2} (T_{cab} - T_{abc} - T_{bca})}. \quad (11)$$

Proof. Metric compatibility makes K_{abc} antisymmetric in a, b . Since the Levi-Civita part has zero torsion, $T_{abc} = K_{acb} - K_{abc}$. Combining this equation with its cyclic permutations yields Eq. (11), proving existence and uniqueness. $\square$

Thus Ω_μ is not an additional arbitrary classical matter field once the tetrad and torsion are fixed. The full-UBT problem is to derive T from the action. In the GR vacuum branch $T = 0$, so $K = 0$. Because every metric-compatible ω_μ lies in $\mathfrak{so}(1, 3)$, its parallel transport preserves η_{ab} and the Lorentz slice, including in branches with nonzero contorsion.

The detailed proof and exact symbolic checks are in `canonical/gr_closure/gap_10omega_connection_elimination.tex` and `tools/verify_gap_10omega_connection.py`.

8 Curvature and special relativity

For a one-sided representation, the curvature is

$$\mathcal{R}_{\mu\nu} = \partial_\mu \Omega_\nu - \partial_\nu \Omega_\mu + [\Omega_\mu, \Omega_\nu]. \quad (12)$$

It must be distinguished from the algebraic tetrad bivector $\Sigma_{\mu\nu}$.

In a flat simply connected region, zero curvature permits a local frame with $\Omega_\mu = 0$. In inertial Cartesian coordinates one can also set $\Gamma^\rho_{\mu\nu} = 0$. The flat limit has an explicit single-field representer.

Theorem 8.1 (Affine representer of a constant Lorentz tetrad). *Let $\bar{E}_\mu \in W_L$ be constant and linearly independent. Then*

$$\Theta_{\text{aff}}(x) = \Theta_0 + \sqrt{\mathcal{N}_0} \bar{E}_\mu x^\mu \quad (13)$$

in the inertial gauge satisfies $\mathcal{N}_0^{-1/2} \partial_\mu \Theta_{\text{aff}} = \bar{E}_\mu$ and induces the constant metric determined by the central anticommutator.

For

$$\bar{E}_0 = i\mathbf{1}, \quad \bar{E}_k = \mathbf{e}_k, \quad (14)$$

Eq. (13) becomes

$$\Theta_{\text{SR}} = \Theta_0 + \sqrt{\mathcal{N}_0} (ix^0 \mathbf{1} + x^k \mathbf{e}_k), \quad (15)$$

which generates $g_{\mu\nu} = \eta_{\mu\nu}$. Every second spacetime derivative vanishes, so the standard flat source-free constant-coefficient second-order master operator annihilates Θ_{SR} . Nonzero connection coefficients in curvilinear coordinates do not by themselves imply curvature.

9 Integrability and representation selection

The tetrad is not independent in UBT; it must satisfy $E_\mu = \mathcal{N}_0^{-1/2} D_\mu \Theta$. Once the classical connection is reconstructed from E and torsion, this becomes an implicit nonlinear first-order system in which the unknown tetrad occurs on both sides.

9.1 One-sided flatness obstruction

Suppose

$$D_\mu^L \Theta = \partial_\mu \Theta + A_\mu \Theta, \quad [D_\mu^L, D_\nu^L] \Theta = F_{\mu\nu}^L \Theta. \quad (16)$$

Theorem 9.1 (One-sided invertible-field no-go). *Assume $E_\nu = \mathcal{N}_0^{-1/2} D_\nu^L \Theta$, torsion-free antisymmetric tetrad compatibility, the same induced one-sided connection on $D_\nu \Theta$, and invertible Θ . Then $F_{\mu\nu}^L = 0$.*

Proof. Torsion-free compatibility makes the antisymmetrised covariant Hessian vanish, so $F_{\mu\nu}^L \Theta = 0$. Right multiplication by Θ^{-1} gives $F_{\mu\nu}^L = 0$. $\square$

Thus the naive one-sided regular representation cannot be the generic curved GR route for invertible Θ . A noninvertible stabilizer sector is a possible exceptional branch, not a representation of arbitrary geometries.

9.2 Two-sided biquaternionic derivative

The algebra $\mathbb{B}$ is naturally a bimodule over itself. Consider

$$\boxed{D_\mu \Theta = \partial_\mu \Theta + A_\mu \Theta - \Theta B_\mu.} \quad (17)$$

A direct expansion gives

$$\boxed{[D_\mu, D_\nu] \Theta = F_{\mu\nu}^A \Theta - \Theta F_{\mu\nu}^B.} \quad (18)$$

Torsion-free antisymmetric compatibility therefore reduces to

$$F_{\mu\nu}^A \Theta = \Theta F_{\mu\nu}^B. \quad (19)$$

For invertible Θ ,

$$\boxed{F_{\mu\nu}^A = \Theta F_{\mu\nu}^B \Theta^{-1}.} \quad (20)$$

The left and right curvatures may be nonzero. The field Θ acts as an intertwiner between them. This does not prove existence for an arbitrary curved tetrad; it only exhibits an algebra-native escape from the stated one-sided flatness obstruction.

9.3 Pure Lorentz-compatible pair and no-new-field reduction

Let $\dagger$ denote Hermitian conjugation and $W_L = \{X \in \mathbb{B} \mid X^\dagger = -X\}$. Requiring Eq. (17) to preserve W_L for every $X \in W_L$ is equivalent to

$$B_\mu = -A_\mu^\dagger + c_\mu \mathbf{1}, \quad c_\mu \in \mathbb{R}. \quad (21)$$

If the same transport preserves the Lorentz quadratic form, the real dilation vanishes. Splitting off the trace then gives, modulo a common central one-form that cancels identically,

$$\boxed{A_\mu = \Omega_\mu, \quad B_\mu = -\Omega_\mu^\dagger,} \quad (22)$$

so that

$$D_\mu X = \partial_\mu X + \Omega_\mu X + X \Omega_\mu^\dagger. \quad (23)$$

Thus A_μ and B_μ are not two new gravitational fields in the pure Lorentz branch. They are the left and right module actions of the one spin connection reconstructed from tetrad and torsion.

Theorem 9.2 (Concurrent-vector obstruction). *Assume Eq. (23), zero torsion, and a nondegenerate tetrad satisfying $D_\mu \Theta = \sqrt{\mathcal{N}_0} E_\mu$. Then the Lorentz-real projection of Θ defines a spacetime vector V such that*

$$\boxed{\nabla_\mu V^\nu = \delta_\mu{}^\nu.} \quad (24)$$

Consequently $\mathcal{L}_V g = 2g$ and $R^\rho{}_{\sigma\mu\nu} V^\sigma = 0$.

Proof. The derivative in Eq. (23) commutes with the Lorentz involution $\mathcal{J}X = -X^\dagger$. Hence $X = (\Theta + \mathcal{J}\Theta)/2 \in W_L$ obeys $D_\mu X = \sqrt{\mathcal{N}_0} E_\mu$. Writing $X = X^a \mathbf{u}_a$, $E_\mu = e_\mu{}^a \mathbf{u}_a$ and defining $V^\nu = \mathcal{N}_0^{-1/2} e_a{}^\nu X^a$, the inverse tetrad postulate gives

$$\nabla_\mu V^\nu = \mathcal{N}_0^{-1/2} e_a{}^\nu D_\mu X^a = e_a{}^\nu e_\mu{}^a = \delta_\mu{}^\nu.$$

The homothety and curvature-kernel identities follow by symmetrising and antisymmetrising covariant derivatives. $\square$

For Schwarzschild $ds^2 = -f dt^2 + f^{-1} dr^2 + r^2 d\Omega^2$ with $f = 1 - 2M/r$, rotational averaging reduces any putative homothetic vector to $V = T(t, r)\partial_t + R(t, r)\partial_r$. The angular equation gives $R = r$, while the rr equation gives $f'(r) = 0$, contradicting $f' = 2M/r^2$ for $M \neq 0$. Therefore the pure pair is closed as a kinematic no-new-field reduction but closed as a no-go for generic curved GR.

The unresolved generic branch may be parameterised as

$$A_\mu = \Omega_\mu + P_\mu, \quad B_\mu = -\Omega_\mu^\dagger + Q_\mu. \quad (25)$$

The relative action $P_\mu X - X Q_\mu$ must be derived from the canonical action and shown composite or auxiliary rather than propagating. A common central constant or one-form cannot remove the obstruction because it cancels from the derivative.

The resulting equation is schematically

$$E_\mu = \mathcal{N}_0^{-1/2} [\partial_\mu \Theta + A_\mu [E, T] \Theta - \Theta B_\mu [E, T]]. \quad (26)$$

It is an implicit nonlinear PDE/fixed-point system. If Θ is restricted to a Jacobi-theta function class, the complete system may additionally be transcendental; implicitness and transcendental functional dependence are separate mathematical properties.

The one-sided/two-sided integrability proof is in `canonical/gr_closure/gap_10i_integrability_selection.tex`. The pure-pair reduction, concurrent-vector theorem, Schwarzschild exclusion, and symbolic verifier are in `canonical/gr_closure/gap_10i_paired_connection_audit.tex` and `tools/verify_gap_10i_paired_connection.py`.

9.4 Exact prescribed-connection integrability criterion

For specified smooth (E_μ, A_μ, B_μ) , the inhomogeneous system can be converted into a homogeneous parallel-section problem. Let $\mathcal{C}_\mu X = A_\mu X - X B_\mu$, $s = \sqrt{\mathcal{N}_0}$, and

$$Y = \begin{pmatrix} \Theta \\ 1 \end{pmatrix}, \quad \hat{\mathcal{C}}_\mu = \begin{pmatrix} \mathcal{C}_\mu & -sE_\mu \\ 0 & 0 \end{pmatrix}. \quad (27)$$

Then $D_\mu \Theta = sE_\mu$ is exactly $(\partial_\mu + \hat{\mathcal{C}}_\mu)Y = 0$. Its curvature is

$$\hat{\mathcal{F}}_{\mu\nu} = \begin{pmatrix} \mathcal{F}_{\mu\nu} & -s(\mathcal{D}_\mu E_\nu - \mathcal{D}_\nu E_\mu) \\ 0 & 0 \end{pmatrix}, \quad (28)$$

where $\mathcal{F}_{\mu\nu} X = F_{\mu\nu}^A X - X F_{\mu\nu}^B$. For an initial value $Y_0 = (\Theta_0, 1)^T$, a single-valued solution exists on a connected domain exactly when the augmented holonomy fixes Y_0 ; on a simply connected domain, flat augmented curvature is a sufficient condition for every initial value. Thus the prescribed-coefficient integrability problem is closed. What remains in GAP-10I-CURVED is the UBT-specific self-consistent selection of (E, A, B, T) , regularity and global continuation. The detailed proof is in `canonical/gr_closure/gap_10i_augmented_holonomy.tex`.

10 Torsion dynamics in the minimal first-order branch

Consider the standard Hilbert–Palatini action with independent tetrad and Lorentz connection,

$$S_{\text{HP}} = \frac{1}{4\kappa} \int \epsilon_{abcd} e^a \wedge e^b \wedge R^{cd}(\omega) + S_{\text{m}}[e, \omega, \Theta], \quad (29)$$

and define $\delta_\omega S_m = \frac{1}{2} \int \tau_{ab} \wedge \delta\omega^{ab}$. Connection variation gives the algebraic Cartan equation

$$\boxed{\epsilon_{abcd} T^a \wedge e^b = \kappa \tau_{cd}.} \quad (30)$$

For a nondegenerate tetrad the 24-component map $T^a \mapsto \epsilon_{abcd} T^a \wedge e^b$ has rank 24. Hence zero spin current forces $T = 0$, while a specified spin current determines a unique contorsion. This closes GAP-10T-PALATINI conditionally. It does not derive the Hilbert–Palatini term, the exact paired Θ spin current, its normalization, or the absence of extra torsion invariants from canonical UBT; therefore the overall GAP-10T-DYN is narrowed rather than fully closed. The proof and exact rank verifier are in `canonical/gr_closure/gap_10t_palatini_torsion_dynamics.tex` and `tools/verify_remaining_gr_subclosures.py`.

11 Lorentz-slice and imaginary-time stability

The real Lorentz slice has an intrinsic fixed-set characterization. Define

$$\mathcal{J}X = -\overline{X^\sharp}. \quad (31)$$

Then $\mathcal{J}$ is an anti-linear involution and $\text{Fix}(\mathcal{J}) = W_L$. Consequently, every unique Cauchy evolution equivariant under $\mathcal{J}$, with $\mathcal{J}$ -real sources and initial data, preserves the Lorentz slice by uniqueness. This closes GAP-10L-SYM conditionally; the canonical equations must still be shown to satisfy the equivariance and well-posedness hypotheses.

For imaginary time, a tetrad flow

$$\partial_\psi e_\mu{}^a = \Lambda_\psi{}^a{}_b e_\mu{}^b, \quad \Lambda_{\psi ab} = -\Lambda_{\psi ba}, \quad (32)$$

is pure local Lorentz gauge and obeys $\partial_\psi g_{\mu\nu} = 0$ exactly. Strictly ψ -independent data are also preserved by every unique ψ -translation-invariant evolution. These results close GAP-10 ψ -KIN and GAP-10 ψ -SYM conditionally, while the overall physical stability problem remains narrowed. Details are in `canonical/gr_closure/gap_10l_psi_symmetry_propagation.tex`.

12 Conditional infrared endpoint of the metric dynamics

Variation of the minimal tetrad–Palatini action gives the Cartan equation and, in the zero-spin-current branch, Einstein’s equation

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \kappa T_{\mu\nu}. \quad (33)$$

Moreover, Lovelock uniqueness implies that a four-dimensional local natural symmetric divergence-free second-order metric equation with no additional light geometric fields must have the form $aG_{\mu\nu} + bg_{\mu\nu} = \kappa T_{\mu\nu}$. Thus the conditional low-energy endpoint is no longer ambiguous: under those assumptions it is Einstein– Λ . GAP-10D-PALATINI and GAP-10D-UNIQUENESS are therefore closed conditionally. GAP-10D itself remains narrowed because canonical UBT must still derive the low-energy assumptions, coefficients and matter action. The precise scope is stated in `canonical/gr_closure/gap_10d_low_energy_uniqueness.tex`.

13 Complex time and regularity

The field may depend on $\tau = t + i\psi$. The metric definition Eq. (4) is pointwise; it does not integrate over ψ . A classical four-dimensional sector may require

$$\partial_\psi g_{\mu\nu} = 0,$$

but this must follow from the dynamics or a clearly stated classical-sector condition.

Smoothness of the physical spacetime field and intrinsic biquaternionic regularity in the argument q are different properties. The latter has not yet been uniquely fixed: Cauchy–Fueter, slice-regular, and UBT-specific Jacobi structures must be compared before a regularity axiom is locked.

14 Relation to earlier compact-fiber closure

Earlier work defined a metric by averaging a quadratic form over the compact imaginary-time coordinate and used the resulting function space to repair a rank count. Those constructions remain mathematically auditable, but they are not required by the central anticommutator theorem and are no longer the canonical local metric definition. They are retained in the repository as an exploratory candidate-completion branch and are documented in `canonical/gr_closure/HISTORICAL_FIBER_ROUTE_STATUS.md`. The branch is rejected as canonical because of weak selection and large representer redundancy, not because its mathematics was disproved. In particular, the earlier fiber-average constructions are not used as a canonical derivation of the metric, rank closure, connection, or Einstein dynamics.

The conceptual distinction is:

- the old route enlarged the effective target by retaining a whole ψ profile;
- the present route uses the ordinary tetrad content of the covariant first jet and obtains $16 - 6 = 10$ locally.

15 Schwarzschild and perturbations: honest current status

The repository contains an explicit and numerically verified construction of the spatial isotropic Schwarzschild metric. The static vacuum Einstein relations determine the lapse uniquely once the spatial branch and boundary conditions are assumed. What remains open is the selection of the complete Schwarzschild tetrad by the canonical UBT master dynamics and connection.

The relation $\partial_\psi \Theta = i\Phi(r)\Theta$ is not used as a canonical derivation of the lapse, and a generic Maxwell or $U(1)_\psi$ field does not generate vacuum Schwarzschild. The perturbation bridge must be rederived in the covariant connection/tetrad formulation; earlier fiber-linearisation results remain conditional results of the exploratory branch.

16 Revised gap ledger

Gap	Status
GAP-10K	CLOSED locally: tetrad-to-metric differential has rank ten.
GAP-10 Ω -KIN	CLOSED: specified tetrad and torsion uniquely determine the metric-compatible frame connection.
GAP-10 Ω -GR	CLOSED: the torsion-free branch is the unique Levi-Civita spin connection, up to Lorentz gauge.
GAP-10T-PALATINI	CLOSED CONDITIONALLY: minimal first-order variation gives an invertible algebraic Cartan torsion equation.
GAP-10T-DYN	NARROWED: derive the minimal branch, exact UBT spin current, normalization, and any extra torsion terms.
GAP-10L-CONN	CLOSED: metric-compatible Lorentz transport preserves η and the Lorentz slice.
GAP-10L-SYM	CLOSED CONDITIONALLY: unique $\mathcal{J}$ -equivariant evolution preserves the Lorentz fixed set.
GAP-10L-DYN	NARROWED: verify equivariance and well-posed uniqueness for the complete UBT dynamics.
GAP-10I-SR	CLOSED: explicit affine Θ representer for every constant Lorentz tetrad.
GAP-10I-1S	CLOSED AS NO-GO: naive one-sided invertible torsion-free branch forces zero curvature.
GAP-10I-PAIR-KIN	CLOSED: Lorentz and metric compatibility reduce the pure pair to one spin connection modulo a cancelling central term.
GAP-10I-PAIR-GR	CLOSED AS NO-GO: the pure pair implies a concurrent vector and excludes generic curved GR and Schwarzschild with $M \neq 0$.
GAP-10I-2S	NARROWED: derive a nontrivial relative bimodule action as composite or auxiliary, with no new propagating field.
GAP-10I-PRESCRIBED	CLOSED: exact augmented-holonomy criterion for specified (E, A, B) .
GAP-10I-CURVED	NARROWED: self-consistent on-shell selection, regularity, and global continuation remain.
GAP-10D-PALATINI / UNIQUENESS	CLOSED CONDITIONALLY: minimal first-order dynamics and Lovelock infrared uniqueness.
GAP-10D	NARROWED: derive those low-energy assumptions, couplings, and matter sector from canonical UBT.

Gap	Status
GAP-10 ψ -KIN / SYM	CLOSED / CLOSED CONDITIONALLY: Lorentz-gauge flow or translation symmetry protects the metric.
GAP-10 ψ	NARROWED: show which mechanism is selected and exclude unstable physical ψ modes.
GAP-B-MASTER	OPEN: perturbation bridge from the original master dynamics.
GAP-U2 Θ	OPEN: canonical dynamical generation of the full Schwarzschild tetrad/lapse.

17 Conclusion

The clean local UBT geometry is

$$\Theta \longrightarrow E_\mu = D_\mu \Theta / \sqrt{\mathcal{N}_0} \longrightarrow \begin{cases} \frac{1}{2}\{E_\mu, E_\nu\}_\# = g_{\mu\nu}\mathbf{1}, \\ \frac{1}{2}[E_\mu, E_\nu]_\# = \Sigma_{\mu\nu}, \\ [D_\mu, D_\nu]\Theta = F_{\mu\nu}^A \Theta - \Theta F_{\mu\nu}^B \end{cases} \text{ in the general two-sided branch.}$$

This structure resolves the local rank objection without auxiliary maps, metric projections, or fiber averaging. The connection is also no longer an arbitrary addition: for specified tetrad and torsion it is uniquely reconstructed, and the torsion-free GR branch is Levi-Civita. Special relativity has an explicit affine single- Θ representer. The naive one-sided curved route is ruled out for generic invertible Θ . The two-sided derivative supplies a nonflat intertwiner condition, while the pure Lorentz-compatible pair is reduced to the existing spin connection and then ruled out as the generic curved route by the concurrent-vector theorem. The decisive remaining work is therefore sharply localized: derive from the single UBT action a nontrivial relative bimodule action and its algebraic elimination, derive the minimal first-order/low-energy assumptions and spin current, solve the self-consistent augmented-holonomy problem for curved tetrads, establish well-posed symmetry propagation, and generate observed geometries on shell.

References

- [1] R. M. Wald, *General Relativity*, University of Chicago Press (1984).
- [2] F. W. Hehl, P. von der Heyde, G. D. Kerlick, and J. M. Nester, “General Relativity with Spin and Torsion: Foundations and Prospects,” *Rev. Mod. Phys.* **48**, 393–416 (1976).
- [3] D. Lovelock, “The Einstein Tensor and Its Generalizations,” *J. Math. Phys.* **12**, 498–501 (1971).
- [4] A. H. Chamseddine, “Complexified Gravity in Noncommutative Spaces,” *Commun. Math. Phys.* **218**, 283–292 (2001).
- [5] E. P. J. de Haas, “Biquaternion Formulation of Relativistic Tensor Dynamics,” arXiv:1401.4470 (2014).